



# wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 01:33 AM GMT

PDB ID : 3MWP  
Title : Nucleoprotein structure of lassa fever virus  
Authors : Qi, X.; Lan, S.; Wang, W.; Schelde, L.M.; Dong, H.; Wallat, G.; Liang, Y.;  
Ly, H.; Dong, C.; Scottish Structural Proteomics Facility (SSPF)  
Deposited on : 2010-05-06  
Resolution : 1.79 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	577	
1	B	577	
1	C	577	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	ZN	B	570	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12477 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	1	0
			4011	2521	697	766	27			
1	B	515	Total	C	N	O	S	0	0	0
			4027	2528	701	771	27			
1	C	506	Total	C	N	O	S	0	0	0
			3961	2488	688	758	27			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP P13699
A	-6	ALA	-	EXPRESSION TAG	UNP P13699
A	-5	MET	-	EXPRESSION TAG	UNP P13699
A	-4	ASP	-	EXPRESSION TAG	UNP P13699
A	-3	HIS	-	EXPRESSION TAG	UNP P13699
A	-2	VAL	-	EXPRESSION TAG	UNP P13699
A	-1	GLU	-	EXPRESSION TAG	UNP P13699
A	0	PHE	-	EXPRESSION TAG	UNP P13699
B	-7	GLY	-	EXPRESSION TAG	UNP P13699
B	-6	ALA	-	EXPRESSION TAG	UNP P13699
B	-5	MET	-	EXPRESSION TAG	UNP P13699
B	-4	ASP	-	EXPRESSION TAG	UNP P13699
B	-3	HIS	-	EXPRESSION TAG	UNP P13699
B	-2	VAL	-	EXPRESSION TAG	UNP P13699
B	-1	GLU	-	EXPRESSION TAG	UNP P13699
B	0	PHE	-	EXPRESSION TAG	UNP P13699
C	-7	GLY	-	EXPRESSION TAG	UNP P13699
C	-6	ALA	-	EXPRESSION TAG	UNP P13699
C	-5	MET	-	EXPRESSION TAG	UNP P13699
C	-4	ASP	-	EXPRESSION TAG	UNP P13699
C	-3	HIS	-	EXPRESSION TAG	UNP P13699
C	-2	VAL	-	EXPRESSION TAG	UNP P13699
C	-1	GLU	-	EXPRESSION TAG	UNP P13699

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	PHE	-	EXPRESSION TAG	UNP P13699

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	175	Total	O	0	0
			175	175		
3	B	192	Total	O	0	0
			192	192		
3	C	108	Total	O	0	0
			108	108		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Nucleoprotein

Chain A:




#### • Molecule 1: Nucleoprotein

Chain B:



#### • Molecule 1: Nucleoprotein

Chain C:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	176.82Å 176.82Å 56.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	153.18 – 1.79 47.59 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (153.18-1.79) 99.9 (47.59-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.6.0081	Depositor
R, $R_{free}$	0.178 , 0.196 0.199 , 0.216	Depositor DCC
$R_{free}$ test set	9262 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.6	Xtriage
Anisotropy	0.560	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 41.6	EDS
Estimated twinning fraction	0.348 for H, K, L 0.652 for K, H, -L 0.008 for -h,-k,l 0.021 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
Reported twinning fraction	0.348 for H, K, L 0.652 for K, H, -L	Depositor
L-test for twinning	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 184177 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12477	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	1/4070 (0.0%)	0.71	1/5493 (0.0%)
1	B	0.64	0/4084	0.76	4/5513 (0.1%)
1	C	0.57	0/4015	0.68	1/5418 (0.0%)
All	All	0.61	1/12169 (0.0%)	0.72	6/16424 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	461	CYS	CB-SG	-8.33	1.68	1.82

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	377	MET	CG-SD-CE	-13.53	78.55	100.20
1	B	115	ARG	NE-CZ-NH1	6.53	123.56	120.30
1	B	115	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	461	CYS	CB-CA-C	-5.73	98.95	110.40
1	B	500	ASP	CB-CG-OD1	5.07	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4011	0	0	4	0
1	B	4027	0	0	3	0
1	C	3961	0	0	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	175	0	0	2	0
3	B	192	0	0	1	0
3	C	108	0	0	2	0
All	All	12477	0	0	13	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

The worst 5 of 13 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:533:ASP:OD2	3:C:655:HOH:O	2.01	0.78
1:B:502:TYR:CE1	1:B:550:LEU:CD1	2.79	0.66
1:A:533:ASP:OD2	3:A:728:HOH:O	2.13	0.65
1:C:389:ASP:OD1	3:C:655:HOH:O	2.18	0.56
1:A:76:ASN:ND2	3:A:744:HOH:O	2.38	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/577 (87%)	496 (99%)	7 (1%)	0	100	100
1	B	507/577 (88%)	502 (99%)	5 (1%)	0	100	100
1	C	492/577 (85%)	483 (98%)	8 (2%)	1 (0%)	56	38
All	All	1502/1731 (87%)	1481 (99%)	20 (1%)	1 (0%)	59	41

All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	C	550	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	450/498 (90%)	449 (100%)	1 (0%)	96	95
1	B	452/498 (91%)	448 (99%)	4 (1%)	87	83
1	C	444/498 (89%)	436 (98%)	8 (2%)	71	58
All	All	1346/1494 (90%)	1333 (99%)	13 (1%)	85	80

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	11	LEU
1	C	212	LYS
1	C	464	SER
1	B	484	ILE
1	C	263	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	512/577 (88%)	0.36	15 (2%)	49 41	31, 33, 45, 66	0
1	B	515/577 (89%)	0.52	16 (3%)	47 39	31, 33, 45, 58	0
1	C	506/577 (87%)	0.46	29 (5%)	23 17	32, 36, 49, 65	0
All	All	1533/1731 (88%)	0.45	60 (3%)	36 30	31, 34, 47, 66	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	ILE	5.0
1	C	545	LEU	4.9
1	B	338	LEU	4.7
1	C	556	ARG	4.4
1	C	516	LYS	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	B	570	1/1	0.55	20.34	103,103,103,103	0
2	ZN	C	570	1/1	0.09	-0.02	44,44,44,44	0
2	ZN	A	570	1/1	0.09	-1.94	33,33,33,33	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.