



Full wwPDB X-ray Structure Validation Report i

Feb 26, 2014 – 03:58 PM GMT

PDB ID : 3MZW
Title : HER2 extracellular region with affinity matured 3-helix affibody ZHER2:342
Authors : Eigenbrot, C.; Ultsch, M.H.
Deposited on : 2010-05-13
Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	624	
2	B	58	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NAG	A	1046	-	X
3	NAG	A	1549	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4890 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine kinase-type cell surface receptor HER2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	569	4395	2731	790	823	51	0	1	0


- Molecule 2 is a protein called Immunoglobulin G-binding protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	53	430	268	79	82	1	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	VAL	ALA	ENGINEERED MUTATION	UNP P38507
B	9	MET	GLN	ENGINEERED MUTATION	UNP P38507
B	10	ARG	GLN	ENGINEERED MUTATION	UNP P38507
B	13	TYR	PHE	ENGINEERED MUTATION	UNP P38507
B	14	TRP	TYR	ENGINEERED MUTATION	UNP P38507
B	17	ALA	LEU	ENGINEERED MUTATION	UNP P38507
B	18	LEU	HIS	ENGINEERED MUTATION	UNP P38507
B	24	ASN	GLU	ENGINEERED MUTATION	UNP P38507
B	25	GLN	GLU	ENGINEERED MUTATION	UNP P38507
B	27	LYS	ARG	ENGINEERED MUTATION	UNP P38507
B	28	ARG	ASN	ENGINEERED MUTATION	UNP P38507
B	29	ALA	GLY	ENGINEERED MUTATION	UNP P38507
B	32	ARG	GLN	ENGINEERED MUTATION	UNP P38507
B	35	TYR	LYS	ENGINEERED MUTATION	UNP P38507

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	2	28	16	2	10	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	9	9	9	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine kinase-type cell surface receptor HER2

Chain A: 



- Molecule 2: Immunoglobulin G-binding protein A

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.73Å 97.84Å 148.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.90 38.99 – 2.90	Depositor EDS
% Data completeness (in resolution range)	96.7 (40.00-2.90) 96.7 (38.99-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.204 , 0.278 0.197 , 0.268	Depositor DCC
R_{free} test set	1333 reflections (6.15%)	DCC
Wilson B-factor (Å ²)	58.6	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.7	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 23006 reflections	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4890	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/4497	0.65	1/6117 (0.0%)
2	B	0.44	0/437	0.61	0/588
All	All	0.47	0/4934	0.64	1/6705 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	LEU	CA-CB-CG	5.53	128.03	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4395	0	0	20	0
2	B	430	0	0	0	0
3	A	28	0	26	0	0
4	A	28	0	0	0	0
5	A	9	0	0	0	0
All	All	4890	0	26	20	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (20) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:447:HIS:CD2	1:A:448:HIS:CD2	2.81	0.69
1:A:156:GLN:NE2	1:A:195:ARG:NH2	2.42	0.67
1:A:379:GLU:O	1:A:405:ASN:ND2	2.31	0.64
1:A:511:GLN:O	1:A:536:ARG:NH1	2.36	0.58
1:A:155:ASN:O	1:A:156:GLN:C	2.50	0.48
1:A:424:GLN:NE2	1:A:447:HIS:CE1	2.82	0.48
1:A:373:GLU:OE2	1:A:373:GLU:N	2.47	0.47
1:A:528:LEU:CB	1:A:529:PRO:CD	2.92	0.47
1:A:125:GLU:OE2	1:A:204:ARG:NH2	2.49	0.46
1:A:155:ASN:O	1:A:158:ALA:N	2.50	0.45
1:A:383:GLU:OE1	1:A:410:ARG:NE	2.51	0.43
1:A:265:GLY:O	1:A:266:ARG:NH1	2.53	0.42
1:A:360:ASP:O	1:A:361:GLY:O	2.36	0.42
1:A:154:ASN:N	1:A:154:ASN:OD1	2.52	0.42
1:A:480:ASP:C	1:A:480:ASP:OD1	2.58	0.42
1:A:221:GLY:O	1:A:231:LEU:N	2.53	0.42
1:A:219:ALA:N	1:A:232:ALA:O	2.53	0.41
1:A:514:ARG:NH1	1:A:531:GLU:OE2	2.53	0.41
1:A:280:ASN:ND2	1:A:410:ARG:O	2.54	0.41
1:A:8:ASP:OD1	1:A:10:LYS:NZ	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/624 (91%)	516 (91%)	44 (8%)	6 (1%)	21	60
2	B	51/58 (88%)	51 (100%)	0	0	100	100
All	All	617/682 (90%)	567 (92%)	44 (7%)	6 (1%)	22	63

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	361	GLY
1	A	366	ASN
1	A	491	GLN
1	A	528	LEU
1	A	256	THR
1	A	305	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	491/538 (91%)	447 (91%)	44 (9%)	14	39
2	B	45/50 (90%)	39 (87%)	6 (13%)	6	16
All	All	536/588 (91%)	486 (91%)	50 (9%)	13	37

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	9[A]	MET
1	A	9[B]	MET
1	A	13	LEU
1	A	21	LEU
1	A	24	LEU
1	A	27	LEU
1	A	37	ASN
1	A	68	GLN
1	A	69	VAL
1	A	71	GLN
1	A	77	LEU
1	A	80	VAL
1	A	101	LEU
1	A	144	THR
1	A	146	LEU
1	A	154	ASN
1	A	155	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	156	GLN
1	A	157	LEU
1	A	159	LEU
1	A	164	THR
1	A	202	CYS
1	A	254	THR
1	A	256	THR
1	A	260	MET
1	A	274	VAL
1	A	283	SER
1	A	284	THR
1	A	306	THR
1	A	318	ARG
1	A	319	VAL
1	A	326	GLU
1	A	355	LEU
1	A	360	ASP
1	A	366	ASN
1	A	479	GLU
1	A	480	ASP
1	A	485	GLU
1	A	510	SER
1	A	546	GLN
1	A	560	ASP
1	A	561	GLN
1	A	577	ARG
2	B	9	MET
2	B	22	LEU
2	B	25	GLN
2	B	33	SER
2	B	45	LEU
2	B	58	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1237	1,4	12,14,15	0.56	0	15,19,21	1.40	2 (13%)
4	NAG	A	1238	4	12,14,15	0.56	0	15,19,21	1.16	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1237	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	1238	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1238	NAG	O5-C5-C6	3.41	110.56	106.98
4	A	1237	NAG	O5-C5-C6	3.28	110.42	106.98
4	A	1237	NAG	C3-C4-C5	-2.84	105.13	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1046	1	12,14,15	0.67	0	15,19,21	1.67	2 (13%)
3	NAG	A	1549	1	12,14,15	0.57	0	15,19,21	2.10	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1046	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1549	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1549	NAG	O5-C5-C6	7.46	114.81	106.98
3	A	1046	NAG	O5-C5-C6	4.87	112.09	106.98
3	A	1046	NAG	C2-N2-C7	2.50	127.30	123.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	569/624 (91%)	-0.20	19 (3%) 44 53	19, 51, 111, 175	3 (0%)
2	B	53/58 (91%)	-0.42	1 (1%) 64 72	35, 56, 101, 114	0
All	All	622/682 (91%)	-0.22	20 (3%) 45 54	19, 51, 110, 175	3 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	LEU	7.0
1	A	303	GLU	4.5
1	A	314	LYS	4.4
1	A	313	SER	3.6
1	A	304	ASP	3.5
2	B	58	LYS	3.3
1	A	315	PRO	3.2
1	A	572	PRO	3.1
1	A	201	GLY	3.0
1	A	566	ALA	2.9
1	A	298	GLN	2.8
1	A	294	PRO	2.8
1	A	571	PRO	2.8
1	A	528	LEU	2.5
1	A	1	THR	2.4
1	A	568	TYR	2.3
1	A	318	ARG	2.3
1	A	100	PRO	2.3
1	A	286	VAL	2.2
1	A	285	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q < 0.9
4	NAG	A	1238	14/15	0.23	5.67	87,107,112,118	0
4	NAG	A	1237	14/15	0.17	1.21	63,80,91,96	0

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q < 0.9
3	NAG	A	1046	14/15	0.39	3.51	94,100,103,104	0
3	NAG	A	1549	14/15	0.36	2.84	95,113,118,118	0

6.5 Other polymers (i)

There are no such residues in this entry.