



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 08:36 AM GMT

PDB ID : 1N0L  
Title : Crystal structure of the PapD chaperone (C-terminally 6x histidine-tagged) bound to the PapE pilus subunit (N-terminal-deleted) from uropathogenic E. coli  
Authors : Sauer, F.G.; Pinkner, J.S.; Waksman, G.; Hultgren, S.J.  
Deposited on : 2002-10-14  
Resolution : 2.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	224	
1	C	224	
2	B	138	
2	D	138	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5226 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperone protein PapD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	Se	0	0	0
			1702	1076	293	328	2	3			
1	C	215	Total	C	N	O	S	Se	0	0	0
			1696	1073	290	328	2	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	MSE	MET	MODIFIED RESIDUE	UNP P15319
A	66	MSE	MET	MODIFIED RESIDUE	UNP P15319
A	172	MSE	MET	MODIFIED RESIDUE	UNP P15319
A	219	HIS	-	EXPRESSION TAG	UNP P15319
A	220	HIS	-	EXPRESSION TAG	UNP P15319
A	221	HIS	-	EXPRESSION TAG	UNP P15319
A	222	HIS	-	EXPRESSION TAG	UNP P15319
A	223	HIS	-	EXPRESSION TAG	UNP P15319
A	224	HIS	-	EXPRESSION TAG	UNP P15319
C	18	MSE	MET	MODIFIED RESIDUE	UNP P15319
C	66	MSE	MET	MODIFIED RESIDUE	UNP P15319
C	172	MSE	MET	MODIFIED RESIDUE	UNP P15319
C	219	HIS	-	EXPRESSION TAG	UNP P15319
C	220	HIS	-	EXPRESSION TAG	UNP P15319
C	221	HIS	-	EXPRESSION TAG	UNP P15319
C	222	HIS	-	EXPRESSION TAG	UNP P15319
C	223	HIS	-	EXPRESSION TAG	UNP P15319
C	224	HIS	-	EXPRESSION TAG	UNP P15319

- Molecule 2 is a protein called mature Fimbrial protein PapE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	118	Total	C	N	O	S	Se	0	0	0
			877	550	146	177	2	2			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	107	Total	C	N	O	S	Se	0	0	0
			787	499	129	155	2	2			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASP	DELETION	UNP P08407
B	?	-	ASN	DELETION	UNP P08407
B	?	-	LEU	DELETION	UNP P08407
B	?	-	THR	DELETION	UNP P08407
B	?	-	PHE	DELETION	UNP P08407
B	?	-	ARG	DELETION	UNP P08407
B	?	-	GLY	DELETION	UNP P08407
B	?	-	LYS	DELETION	UNP P08407
B	?	-	LEU	DELETION	UNP P08407
B	?	-	ILE	DELETION	UNP P08407
B	?	-	ILE	DELETION	UNP P08407
B	46	MSE	MET	MODIFIED RESIDUE	UNP P08407
B	55	MSE	MET	MODIFIED RESIDUE	UNP P08407
B	130	MSE	MET	MODIFIED RESIDUE	UNP P08407
D	?	-	ASP	DELETION	UNP P08407
D	?	-	ASN	DELETION	UNP P08407
D	?	-	LEU	DELETION	UNP P08407
D	?	-	THR	DELETION	UNP P08407
D	?	-	PHE	DELETION	UNP P08407
D	?	-	ARG	DELETION	UNP P08407
D	?	-	GLY	DELETION	UNP P08407
D	?	-	LYS	DELETION	UNP P08407
D	?	-	LEU	DELETION	UNP P08407
D	?	-	ILE	DELETION	UNP P08407
D	?	-	ILE	DELETION	UNP P08407
D	46	MSE	MET	MODIFIED RESIDUE	UNP P08407
D	55	MSE	MET	MODIFIED RESIDUE	UNP P08407
D	130	MSE	MET	MODIFIED RESIDUE	UNP P08407

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	54	Total	O	0	0
			54	54		
3	B	21	Total	O	0	0
			21	21		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	73	Total 73	O 73	0	0
3	D	16	Total 16	O 16	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Chaperone protein PapD

Chain A:




#### • Molecule 1: Chaperone protein PapD

Chain C:




#### • Molecule 2: mature Fimbrial protein PapE


Chain B:



#### • Molecule 2: mature Fimbrial protein PapE

Chain D:





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.07Å 56.86Å 60.93Å 108.57° 89.26° 104.14°	Depositor
Resolution (Å)	24.73 – 2.30 28.81 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.9 (24.73-2.30) 91.0 (28.81-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.88 (at 2.31Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.226 , 0.279 0.225 , 0.279	Depositor DCC
$R_{free}$ test set	2020 reflections (7.81%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 31.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 29105 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/1734	0.64	0/2349
1	C	0.37	0/1728	0.64	0/2342
2	B	0.34	0/889	0.61	0/1208
2	D	0.31	0/797	0.60	0/1083
All	All	0.35	0/5148	0.63	0/6982

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1702	0	1711	63	0
1	C	1696	0	1700	67	0
2	B	877	0	852	61	0
2	D	787	0	780	57	0
3	A	54	0	0	3	0
3	B	21	0	0	3	0
3	C	73	0	0	3	0
3	D	16	0	0	1	0
All	All	5226	0	5043	237	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 23.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:215:LYS:H	1:A:215:LYS:HD2	1.17	1.04
1:A:32:LEU:HB2	1:A:93:ILE:HB	1.38	1.00
1:A:89:ASN:HD21	1:A:110:LYS:HE2	1.27	0.99
1:C:44:LYS:HE2	1:C:46:ILE:HD11	1.44	0.97
2:B:71:VAL:HG12	2:B:72:GLN:H	1.31	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/224 (95%)	203 (95%)	10 (5%)	0	100	100
1	C	213/224 (95%)	201 (94%)	12 (6%)	0	100	100
2	B	110/138 (80%)	95 (86%)	12 (11%)	3 (3%)	8	5
2	D	99/138 (72%)	89 (90%)	5 (5%)	5 (5%)	3	1
All	All	635/724 (88%)	588 (93%)	39 (6%)	8 (1%)	18	17

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	55	MSE
2	D	94	ILE
2	D	40	LYS
2	D	90	ASN
2	D	91	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/197 (97%)	185 (97%)	6 (3%)	52	68
1	C	190/197 (96%)	182 (96%)	8 (4%)	40	53
2	B	97/113 (86%)	97 (100%)	0	100	100
2	D	86/113 (76%)	82 (95%)	4 (5%)	36	47
All	All	564/620 (91%)	546 (97%)	18 (3%)	51	67

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	96	ARG
1	C	136	LYS
2	D	24	TRP
1	C	26	ASN
1	C	53	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	112	ASN
1	C	26	ASN
1	C	179	GLN
2	B	63	ASN
1	C	185	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/224 (95%)	-0.02	4 (1%) 64 73	13, 34, 54, 74	0
1	C	215/224 (95%)	-0.03	3 (1%) 72 80	12, 34, 53, 65	0
2	B	118/138 (85%)	0.90	19 (16%) 2 4	17, 53, 74, 80	0
2	D	107/138 (77%)	0.84	15 (14%) 3 6	18, 54, 75, 83	0
All	All	655/724 (90%)	0.28	41 (6%) 19 28	12, 38, 69, 83	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	110	THR	6.4
2	B	79	SER	4.8
2	D	89	SER	4.4
2	D	92	GLY	4.4
2	B	27	VAL	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.