



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:42 AM GMT

PDB ID : 3N1Q
Title : Crystal Structure of DhhN bound to CDOFn3
Authors : Kavran, J.M.; Leahy, D.J.
Deposited on : 2010-05-16
Resolution : 2.89 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	170	
1	B	170	
1	E	170	
2	C	102	
2	D	102	
2	F	102	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5767 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Desert hedgehog protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	151	Total	C	N	O	S	0	0	0
			1208	753	224	227	4			
1	A	151	Total	C	N	O	S	0	0	0
			1208	753	224	227	4			
1	E	150	Total	C	N	O	S	0	0	0
			1199	748	222	225	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	20	GLY	-	EXPRESSION TAG	UNP O43323
B	21	SER	-	EXPRESSION TAG	UNP O43323
B	22	GLY	-	EXPRESSION TAG	UNP O43323
B	23	PRO	-	EXPRESSION TAG	UNP O43323
A	20	GLY	-	EXPRESSION TAG	UNP O43323
A	21	SER	-	EXPRESSION TAG	UNP O43323
A	22	GLY	-	EXPRESSION TAG	UNP O43323
A	23	PRO	-	EXPRESSION TAG	UNP O43323
E	20	GLY	-	EXPRESSION TAG	UNP O43323
E	21	SER	-	EXPRESSION TAG	UNP O43323
E	22	GLY	-	EXPRESSION TAG	UNP O43323
E	23	PRO	-	EXPRESSION TAG	UNP O43323

- Molecule 2 is a protein called Cell adhesion molecule-related/down-regulatedby oncogenes.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	87	Total	C	N	O	S	0	0	0
			714	450	114	144	6			
2	D	87	Total	C	N	O	S	0	0	0
			714	450	114	144	6			
2	F	87	Total	C	N	O	S	0	0	0
			714	450	114	144	6			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	823	GLY	-	EXPRESSION TAG	UNP Q4KMG0
C	824	SER	-	EXPRESSION TAG	UNP Q4KMG0
C	825	THR	-	EXPRESSION TAG	UNP Q4KMG0
D	823	GLY	-	EXPRESSION TAG	UNP Q4KMG0
D	824	SER	-	EXPRESSION TAG	UNP Q4KMG0
D	825	THR	-	EXPRESSION TAG	UNP Q4KMG0
F	823	GLY	-	EXPRESSION TAG	UNP Q4KMG0
F	824	SER	-	EXPRESSION TAG	UNP Q4KMG0
F	825	THR	-	EXPRESSION TAG	UNP Q4KMG0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	E	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		
4	A	2	Total	Ca	0	0
			2	2		
4	E	2	Total	Ca	0	0
			2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Desert hedgehog protein

Chain B: 




- Molecule 1: Desert hedgehog protein

Chain A: 




- Molecule 1: Desert hedgehog protein

Chain E: 




- Molecule 2: Cell adhesion molecule-related/down-regulatedby oncogenes

Chain C: 




- Molecule 2: Cell adhesion molecule-related/down-regulatedby oncogenes

Chain D: 



- Molecule 2: Cell adhesion molecule-related/down-regulatedby oncogenes

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.72Å 100.27Å 97.26Å 90.00° 98.98° 90.00°	Depositor
Resolution (Å)	22.73 – 2.89 22.73 – 2.89	Depositor EDS
% Data completeness (in resolution range)	89.6 (22.73-2.89) 99.5 (22.73-2.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.89Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.237 , 0.288 0.239 , 0.285	Depositor DCC
R_{free} test set	988 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , -4.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 19342 reflections	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	5767	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/1235	0.38	0/1673
1	B	0.25	0/1235	0.39	0/1673
1	E	0.25	0/1226	0.43	1/1661 (0.1%)
2	C	0.22	0/732	0.39	0/987
2	D	0.21	0/732	0.37	0/987
2	F	0.22	0/732	0.37	0/987
All	All	0.23	0/5892	0.39	1/7968 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	46	LYS	C-N-CA	5.39	135.18	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	117	MET	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1208	0	0	5	0
1	B	1208	0	0	0	0
1	E	1199	0	0	2	0
2	C	714	0	0	0	0
2	D	714	0	0	0	0
2	F	714	0	0	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	E	1	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	E	2	0	0	0	0
5	E	1	0	0	0	0
All	All	5767	0	0	13	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (13) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:F:861:GLY:CA	2:F:883:GLU:O	2.29	0.81
2:F:882:VAL:CG1	2:F:883:GLU:N	2.45	0.78
1:A:55:ARG:NH2	1:A:138:ASP:O	2.35	0.59
2:F:886:LYS:NZ	2:F:888:TRP:CH2	2.79	0.51
1:A:55:ARG:NH2	1:A:138:ASP:C	2.65	0.49
2:F:886:LYS:NZ	2:F:888:TRP:CZ2	2.84	0.46
1:A:137:GLN:O	1:A:138:ASP:CB	2.63	0.45
2:F:882:VAL:CG1	2:F:886:LYS:CD	2.95	0.43
1:A:55:ARG:NH1	1:A:138:ASP:CB	2.82	0.43
1:A:121:VAL:CG1	1:A:122:ARG:N	2.82	0.42
2:F:886:LYS:CD	2:F:888:TRP:O	2.68	0.41
1:E:172:ASP:OD2	1:E:187:LYS:N	2.54	0.41
1:E:141:HIS:NE2	1:E:183:HIS:CE1	2.89	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/170 (88%)	143 (96%)	6 (4%)	0	100	100
1	B	149/170 (88%)	143 (96%)	6 (4%)	0	100	100
1	E	148/170 (87%)	139 (94%)	9 (6%)	0	100	100
2	C	83/102 (81%)	83 (100%)	0	0	100	100
2	D	83/102 (81%)	80 (96%)	3 (4%)	0	100	100
2	F	83/102 (81%)	76 (92%)	5 (6%)	2 (2%)	9	35
All	All	695/816 (85%)	664 (96%)	29 (4%)	2 (0%)	50	85

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	882	VAL
2	F	830	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/138 (91%)	120 (95%)	6 (5%)	35	74
1	B	126/138 (91%)	123 (98%)	3 (2%)	61	91
1	E	125/138 (91%)	121 (97%)	4 (3%)	51	88
2	C	79/93 (85%)	78 (99%)	1 (1%)	80	96
2	D	79/93 (85%)	78 (99%)	1 (1%)	80	96
2	F	79/93 (85%)	78 (99%)	1 (1%)	80	96
All	All	614/693 (89%)	598 (97%)	16 (3%)	59	90

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	106	ARG
1	B	156	ARG
1	B	176	TYR
2	C	873	ASN
1	A	80	ASN
1	A	132	ASP
1	A	161	LEU
1	A	176	TYR
1	A	184	VAL
1	A	189	ASP
2	D	895	GLN
1	E	80	ASN
1	E	116	ASN
1	E	122	ARG
1	E	176	TYR
2	F	885	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/170 (88%)	0.08	0 100 100	20, 35, 57, 74	0
1	B	151/170 (88%)	-0.15	2 (1%) 74 82	14, 25, 49, 71	0
1	E	150/170 (88%)	0.20	2 (1%) 74 82	19, 38, 67, 92	0
2	C	87/102 (85%)	0.08	4 (4%) 31 38	12, 33, 65, 86	0
2	D	87/102 (85%)	0.28	5 (5%) 23 28	21, 40, 71, 89	0
2	F	87/102 (85%)	0.25	8 (9%) 9 11	20, 33, 79, 95	0
All	All	713/816 (87%)	0.10	21 (2%) 49 58	12, 33, 68, 95	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	910	GLY	4.2
2	D	909	GLU	3.3
1	E	75	ARG	3.3
2	D	859	ILE	3.2
2	F	883	GLU	3.0
2	C	910	GLY	2.9
1	B	70	GLY	2.8
2	F	882	VAL	2.8
1	E	138	ASP	2.7
2	D	885	SER	2.6
2	C	911	GLY	2.5
2	D	910	GLY	2.5
2	C	849	TYR	2.5
2	F	909	GLU	2.4
2	F	860	GLN	2.3
2	F	829	GLY	2.3
2	D	911	GLY	2.3
2	F	908	ASN	2.3
1	B	39	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	884	GLY	2.1
2	C	909	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CA	E	191	1/1	0.23	1.64	47,47,47,47	0
4	CA	B	191	1/1	0.18	0.21	33,33,33,33	0
4	CA	B	192	1/1	0.17	-0.32	23,23,23,23	0
4	CA	A	192	1/1	0.17	-0.39	22,22,22,22	0
3	ZN	E	190	1/1	0.14	-1.60	66,66,66,66	0
3	ZN	B	190	1/1	0.08	-2.03	46,46,46,46	0
4	CA	A	191	1/1	0.13	-2.22	25,25,25,25	0
4	CA	E	192	1/1	0.12	-2.38	19,19,19,19	0
3	ZN	A	190	1/1	0.10	-3.39	50,50,50,50	0

6.5 Other polymers ⓘ

There are no such residues in this entry.