



wwPDB X-ray Structure Validation Summary Report

Sep 29, 2014 – 10:59 AM EDT

PDB ID : 4N1H
Title : Structure of a single-domain camelid antibody fragment cAb-F11N in complex with the BlaP beta-lactamase from *Bacillus licheniformis*
Authors : Pain, C.; Kerff, F.; Herman, R.; Sauvage, E.; Preumont, S.; Charlier, P.; Dumoulin, M.
Deposited on : 2013-10-04
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23828
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23828

1 Overall quality at a glance





The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	273	
1	C	273	
2	B	133	
2	D	133	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5962 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2006	1259	347	397	3			
1	C	258	Total	C	N	O	S	0	0	0
			2014	1263	348	400	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	197A	PRO	-	INSERTION	UNP P00808
A	197B	GLY	-	INSERTION	UNP P00808
A	296	GLY	-	EXPRESSION TAG	UNP P00808
A	297	PRO	-	EXPRESSION TAG	UNP P00808
A	298	HIS	-	EXPRESSION TAG	UNP P00808
A	299	HIS	-	EXPRESSION TAG	UNP P00808
A	300	HIS	-	EXPRESSION TAG	UNP P00808
A	301	HIS	-	EXPRESSION TAG	UNP P00808
A	302	HIS	-	EXPRESSION TAG	UNP P00808
C	197A	PRO	-	INSERTION	UNP P00808
C	197B	GLY	-	INSERTION	UNP P00808
C	296	GLY	-	EXPRESSION TAG	UNP P00808
C	297	PRO	-	EXPRESSION TAG	UNP P00808
C	298	HIS	-	EXPRESSION TAG	UNP P00808
C	299	HIS	-	EXPRESSION TAG	UNP P00808
C	300	HIS	-	EXPRESSION TAG	UNP P00808
C	301	HIS	-	EXPRESSION TAG	UNP P00808
C	302	HIS	-	EXPRESSION TAG	UNP P00808

- Molecule 2 is a protein called Camelid heavy-chain antibody variable fragment cAb-F11N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	126	Total	C	N	O	S	0	0	0
			962	598	168	192	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	126	Total	C	N	O	S	0	0	0
			962	598	168	192	4			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	8	Total	O	0	0
			8	8		
3	C	10	Total	O	0	0
			10	10		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-lactamase

Chain A:




• Molecule 1: Beta-lactamase

Chain C:



• Molecule 2: Camelid heavy-chain antibody variable fragment cAb-F11N


Chain B:



• Molecule 2: Camelid heavy-chain antibody variable fragment cAb-F11N

Chain D:





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.09Å 89.82Å 75.75Å 90.00° 95.17° 90.00°	Depositor
Resolution (Å)	75.44 – 3.00 48.57 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (75.44-3.00) 99.5 (48.57-3.00)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.208 , 0.284 0.211 , 0.288	Depositor DCC
R_{free} test set	794 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	47.5	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 17.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 15835 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5962	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	1/2036 (0.0%)	0.66	0/2755
1	C	0.46	1/2045 (0.0%)	0.63	0/2770
2	B	0.53	1/981 (0.1%)	0.63	0/1326
2	D	0.55	1/981 (0.1%)	0.62	0/1326
All	All	0.49	4/6043 (0.1%)	0.64	0/8177

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	117	TRP	CD2-CE2	5.24	1.47	1.41
1	A	229	TRP	CD2-CE2	5.16	1.47	1.41
1	C	251	TRP	CD2-CE2	5.01	1.47	1.41
2	D	80	TRP	CD2-CE2	5.00	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2006	0	2022	55	0
1	C	2014	0	2026	65	0
2	B	962	0	918	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	962	0	918	21	0
3	A	8	0	0	2	0
3	C	10	0	0	0	0
All	All	5962	0	5884	170	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 170 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:52:SER:HB2	2:D:57:ASP:HB2	1.30	1.12
1:C:106:ASN:HB3	1:C:109:THR:HG22	1.32	1.09
1:C:109:THR:HG21	1:C:133:ALA:HB3	1.34	1.04
2:B:34:MET:HG2	2:B:79:VAL:HG21	1.45	0.97
2:B:34:MET:HE1	2:B:98:ALA:HB2	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/273 (93%)	241 (95%)	11 (4%)	1 (0%)	43	87
1	C	256/273 (94%)	241 (94%)	13 (5%)	2 (1%)	27	76
2	B	124/133 (93%)	114 (92%)	9 (7%)	1 (1%)	27	76
2	D	124/133 (93%)	110 (89%)	12 (10%)	2 (2%)	14	56
All	All	757/812 (93%)	706 (93%)	45 (6%)	6 (1%)	27	76

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	45	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	103	VAL
1	C	289	ALA
2	D	112	HIS
1	C	103	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/230 (94%)	196 (91%)	20 (9%)	13	45
1	C	217/230 (94%)	199 (92%)	18 (8%)	16	52
2	B	99/106 (93%)	86 (87%)	13 (13%)	6	25
2	D	99/106 (93%)	90 (91%)	9 (9%)	14	46
All	All	631/672 (94%)	571 (90%)	60 (10%)	12	44

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	85	SER
1	C	68	PHE
2	D	104	LEU
2	B	121	THR
1	C	106	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	77	ASN
2	B	112	HIS
1	C	83	GLN
2	B	39	GLN
1	C	54	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/273 (94%)	-0.31	0 100 100	21, 38, 62, 75	0
1	C	258/273 (94%)	-0.34	0 100 100	21, 38, 64, 76	0
2	B	126/133 (94%)	-0.06	0 100 100	31, 49, 82, 110	0
2	D	126/133 (94%)	-0.03	0 100 100	35, 50, 80, 94	0
All	All	767/812 (94%)	-0.23	0 100 100	21, 42, 71, 110	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.