



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 11:48 PM GMT

PDB ID : 4N2S  
Title : Crystal Structure of THA8 in complex with Zm1a-6 RNA  
Authors : Ke, J.; Chen, R.Z.; Ban, T.; Zhou, X.E.; Gu, X.; Brunzelle, J.S.; Zhu, J.K.;  
Melcher, K.; Xu, H.E.  
Deposited on : 2013-10-06  
Resolution : 3.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)
RNA backbone	1838	1070 (3.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	257	
2	B	12	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1596 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THA8 RNA binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	0	0
			1491	933	272	281	5			

- Molecule 2 is a RNA chain called Zm1a-6 RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	4	Total	C	N	O	P	0	0	0
			90	40	20	26	4			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total	O	0	0
			12	12		
3	B	3	Total	O	0	0
			3	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: THA8 RNA binding protein

Chain A:



- Molecule 2: Zm1a-6 RNA

Chain B:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.25Å 88.25Å 78.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 39.47 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.00) 100.0 (39.47-3.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.205 , 0.232 0.208 , 0.233	Depositor DCC
$R_{free}$ test set	315 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	83.0	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 6644 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	1596	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.62	0/1520	0.65	0/2064
2	B	0.44	0/101	0.63	0/156
All	All	0.61	0/1621	0.65	0/2220

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1491	0	1474	51	0
2	B	90	0	45	6	0
3	A	12	0	0	1	0
3	B	3	0	0	0	0
All	All	1596	0	1519	57	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 18.

All (57) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:154:ALA:O	1:A:158:SER:HB2	1.59	1.02

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:158:SER:O	1:A:160:GLU:HA	1.59	1.01
1:A:66:PRO:HG2	1:A:67:PRO:HD3	1.60	0.83
1:A:115:ARG:HG2	1:A:115:ARG:HH11	1.46	0.79
1:A:160:GLU:HG2	1:A:160:GLU:O	1.88	0.72
1:A:150:LYS:O	1:A:162:VAL:HG11	1.90	0.71
1:A:232:ASP:O	1:A:236:ARG:HG3	1.93	0.69
1:A:66:PRO:CG	1:A:67:PRO:HD3	2.27	0.65
2:B:3:G:H21	2:B:5:A:H4'	1.60	0.65
1:A:234:GLU:O	1:A:238:LEU:HB2	1.99	0.62
1:A:234:GLU:OE1	1:A:238:LEU:HD13	2.00	0.62
1:A:50:ILE:HG13	1:A:51:HIS:N	2.14	0.61
1:A:45:LEU:HD11	1:A:91:GLU:HG2	1.81	0.61
1:A:158:SER:C	1:A:160:GLU:HA	2.20	0.60
1:A:47:THR:HA	1:A:50:ILE:HG12	1.84	0.59
1:A:192:GLU:HG2	1:A:197:LYS:HE2	1.87	0.57
1:A:139:VAL:O	1:A:143:VAL:HG23	2.06	0.55
1:A:208:ARG:HB3	1:A:208:ARG:NH1	2.23	0.54
1:A:54:GLN:O	1:A:58:ARG:HG3	2.08	0.54
1:A:149:GLU:HA	1:A:152:ARG:NH1	2.24	0.53
1:A:67:PRO:HA	3:A:311:HOH:O	2.08	0.52
1:A:219:LEU:HB3	1:A:222:GLU:OE1	2.09	0.52
1:A:140:ASP:O	1:A:144:GLU:HG2	2.10	0.52
2:B:3:G:H5''	2:B:5:A:H1'	1.92	0.51
2:B:3:G:H2'	2:B:4:G:H8	1.74	0.51
2:B:3:G:N2	2:B:5:A:H4'	2.25	0.51
1:A:115:ARG:CG	1:A:115:ARG:HH11	2.21	0.51
1:A:116:PRO:HG3	1:A:149:GLU:HB3	1.92	0.51
1:A:115:ARG:HG2	1:A:115:ARG:NH1	2.20	0.50
1:A:79:ARG:HH11	1:A:79:ARG:HG2	1.78	0.49
1:A:105:LEU:HD22	1:A:124:PHE:CE2	2.47	0.49
1:A:140:ASP:OD1	1:A:186:ALA:HB2	2.15	0.47
1:A:239:PRO:O	1:A:240:ASP:OD1	2.32	0.47
1:A:50:ILE:HG13	1:A:51:HIS:H	1.78	0.47
1:A:105:LEU:HB2	1:A:124:PHE:CZ	2.50	0.47
1:A:189:ARG:HB3	1:A:189:ARG:HE	1.46	0.46
1:A:82:ARG:HG3	1:A:113:TRP:CE3	2.50	0.46
1:A:82:ARG:HG3	1:A:113:TRP:CD2	2.51	0.46
1:A:53:VAL:O	1:A:57:LYS:HG3	2.16	0.46
1:A:221:GLU:O	1:A:225:GLU:HG3	2.16	0.46
1:A:119:GLU:O	1:A:123:THR:HG23	2.15	0.45
1:A:151:GLU:O	1:A:155:ALA:HB3	2.17	0.45
1:A:54:GLN:HB3	1:A:58:ARG:HH11	1.82	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:184:ALA:HB1	1:A:219:LEU:CD1	2.48	0.44
1:A:108:ALA:O	1:A:114:TYR:HB2	2.18	0.44
1:A:47:THR:HA	1:A:50:ILE:CG1	2.48	0.44
1:A:51:HIS:HD2	1:A:54:GLN:NE2	2.17	0.43
1:A:159:SER:HA	1:A:160:GLU:CB	2.48	0.43
1:A:114:TYR:O	1:A:115:ARG:HG2	2.18	0.43
1:A:111:GLU:HB3	1:A:112:PRO:HD2	2.01	0.43
2:B:3:G:H2'	2:B:4:G:C8	2.54	0.43
1:A:115:ARG:NH1	1:A:115:ARG:CG	2.80	0.42
2:B:5:A:H3'	2:B:5:A:N3	2.33	0.42
1:A:45:LEU:HD23	1:A:87:ALA:HB1	2.02	0.42
1:A:225:GLU:O	1:A:228:ALA:HB3	2.21	0.41
1:A:135:ALA:O	1:A:139:VAL:HG23	2.20	0.41
1:A:238:LEU:HG	1:A:239:PRO:HD2	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	195/257 (76%)	185 (95%)	7 (4%)	3 (2%)	15 58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	239	PRO
1	A	153	GLY
1	A	157	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/192 (73%)	127 (90%)	14 (10%)	11	40

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	ARG
1	A	86	LEU
1	A	115	ARG
1	A	116	PRO
1	A	132	ASP
1	A	142	LEU
1	A	159	SER
1	A	160	GLU
1	A	162	VAL
1	A	205	TYR
1	A	208	ARG
1	A	236	ARG
1	A	238	LEU
1	A	240	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	HIS

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	3/12 (25%)	2 (66%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	4	G
2	B	6	A

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	197/257 (76%)	-0.21	0 100 100	51, 72, 106, 169	0
2	B	4/12 (33%)	0.46	0 100 100	100, 141, 165, 166	0
All	All	201/269 (74%)	-0.19	0 100 100	51, 72, 116, 169	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.