



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:48 PM GMT

PDB ID : 3N4F
Title : CRYSTAL STRUCTURE OF Mandelate racemase/muconate lactonizing protein from *Geobacillus* sp. Y412MC10
Authors : Malashkevich, V.N.; Toro, R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-05-21
Resolution : 1.88 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	5260 (1.90-1.86)
Clashscore	79885	6268 (1.90-1.86)
Ramachandran outliers	78287	6195 (1.90-1.86)
Sidechain outliers	78261	6196 (1.90-1.86)
RSRZ outliers	66119	5262 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	392	
1	B	392	
1	C	392	
1	D	392	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13659 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mandelate racemase/muconate lactonizing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	Se	0	5	0
			3111	1970	550	581	3	7			
1	B	386	Total	C	N	O	S	Se	0	4	0
			3069	1941	540	577	3	8			
1	C	386	Total	C	N	O	S	Se	0	3	0
			3061	1940	537	574	3	7			
1	D	376	Total	C	N	O	S	Se	0	4	0
			3001	1905	525	561	3	7			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP D3EID5
A	2	SER	-	expression tag	UNP D3EID5
A	3	LEU	-	expression tag	UNP D3EID5
A	385	GLU	-	expression tag	UNP D3EID5
A	386	GLY	-	expression tag	UNP D3EID5
A	387	HIS	-	expression tag	UNP D3EID5
A	388	HIS	-	expression tag	UNP D3EID5
A	389	HIS	-	expression tag	UNP D3EID5
A	390	HIS	-	expression tag	UNP D3EID5
A	391	HIS	-	expression tag	UNP D3EID5
A	392	HIS	-	expression tag	UNP D3EID5
B	1	MSE	-	expression tag	UNP D3EID5
B	2	SER	-	expression tag	UNP D3EID5
B	3	LEU	-	expression tag	UNP D3EID5
B	385	GLU	-	expression tag	UNP D3EID5
B	386	GLY	-	expression tag	UNP D3EID5
B	387	HIS	-	expression tag	UNP D3EID5
B	388	HIS	-	expression tag	UNP D3EID5
B	389	HIS	-	expression tag	UNP D3EID5
B	390	HIS	-	expression tag	UNP D3EID5
B	391	HIS	-	expression tag	UNP D3EID5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	392	HIS	-	expression tag	UNP D3EID5
C	1	MSE	-	expression tag	UNP D3EID5
C	2	SER	-	expression tag	UNP D3EID5
C	3	LEU	-	expression tag	UNP D3EID5
C	385	GLU	-	expression tag	UNP D3EID5
C	386	GLY	-	expression tag	UNP D3EID5
C	387	HIS	-	expression tag	UNP D3EID5
C	388	HIS	-	expression tag	UNP D3EID5
C	389	HIS	-	expression tag	UNP D3EID5
C	390	HIS	-	expression tag	UNP D3EID5
C	391	HIS	-	expression tag	UNP D3EID5
C	392	HIS	-	expression tag	UNP D3EID5
D	1	MSE	-	expression tag	UNP D3EID5
D	2	SER	-	expression tag	UNP D3EID5
D	3	LEU	-	expression tag	UNP D3EID5
D	385	GLU	-	expression tag	UNP D3EID5
D	386	GLY	-	expression tag	UNP D3EID5
D	387	HIS	-	expression tag	UNP D3EID5
D	388	HIS	-	expression tag	UNP D3EID5
D	389	HIS	-	expression tag	UNP D3EID5
D	390	HIS	-	expression tag	UNP D3EID5
D	391	HIS	-	expression tag	UNP D3EID5
D	392	HIS	-	expression tag	UNP D3EID5

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Mg 2 2	0	0
2	A	2	Total Mg 2 2	0	0
2	D	2	Total Mg 2 2	0	0
2	C	2	Total Mg 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	394	Total O 394 394	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	328	Total 328	O 328	0	0
3	C	450	Total 450	O 450	0	0
3	D	237	Total 237	O 237	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mandelate racemase/muconate lactonizing protein

Chain A: 



- Molecule 1: Mandelate racemase/muconate lactonizing protein

Chain B: 




- Molecule 1: Mandelate racemase/muconate lactonizing protein

Chain C: 



- Molecule 1: Mandelate racemase/muconate lactonizing protein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.39Å 66.10Å 151.54Å 90.00° 94.33° 90.00°	Depositor
Resolution (Å)	8.00 – 1.88 8.00 – 1.88	Depositor EDS
% Data completeness (in resolution range)	93.1 (8.00-1.88) 93.1 (8.00-1.88)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	19.12 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.185 , 0.242 0.195 , 0.252	Depositor DCC
R_{free} test set	5793 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.51 , 39.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 114670 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13659	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 66.71 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.7857e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	0/3194	0.64	1/4317 (0.0%)
1	B	0.54	1/3147 (0.0%)	0.64	2/4253 (0.0%)
1	C	0.59	0/3136	0.65	0/4240
1	D	0.44	0/3079	0.57	0/4162
All	All	0.54	1/12556 (0.0%)	0.63	3/16972 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	330	ARG	CZ-NH2	-5.51	1.25	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	330	ARG	NE-CZ-NH1	9.88	125.24	120.30
1	B	330	ARG	NE-CZ-NH2	-9.71	115.44	120.30
1	A	389	HIS	N-CA-C	5.46	125.73	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3111	0	0	14	1
1	B	3069	0	0	11	0
1	C	3061	0	0	10	0
1	D	3001	0	0	10	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	394	0	0	7	0
3	B	328	0	0	5	0
3	C	450	0	0	7	0
3	D	237	0	0	3	0
All	All	13659	0	0	44	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (44) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:345:MSE:SE	3:C:1613:HOH:O	2.23	1.05
1:B:345:MSE:SE	3:B:1560:HOH:O	2.26	1.02
1:A:390:HIS:ND1	1:A:391:HIS:CD2	2.30	0.99
1:A:345:MSE:SE	3:A:494:HOH:O	2.35	0.93
1:D:345:MSE:SE	3:D:1554:HOH:O	2.42	0.87
1:A:178:MSE:SE	3:A:1570:HOH:O	2.56	0.74
1:C:45:VAL:CG1	3:C:1613:HOH:O	2.48	0.60
1:C:271:ILE:CD1	1:C:282[B]:LEU:CD2	2.80	0.59
1:A:386:GLY:O	1:A:387:HIS:CB	2.54	0.55
1:A:253[A]:ARG:NE	3:A:517:HOH:O	2.41	0.54
1:D:102:TRP:CH2	3:D:1512:HOH:O	2.53	0.53
1:B:330:ARG:NH2	3:B:687:HOH:O	2.40	0.53
1:A:390:HIS:ND1	1:A:391:HIS:NE2	2.57	0.52
1:B:16:LYS:CD	1:B:388:HIS:O	2.59	0.51
1:A:250:LEU:CD1	3:A:1603:HOH:O	2.59	0.51
1:B:388:HIS:NE2	1:D:353:GLU:OE1	2.45	0.48
1:D:162:LYS:NZ	1:D:339:ASP:OD2	2.46	0.48
1:B:230:TYR:OH	1:B:333:GLU:OE1	2.31	0.48
1:D:45:VAL:CG1	3:D:1554:HOH:O	2.62	0.47
1:C:7:GLY:N	3:C:1572:HOH:O	2.47	0.47
1:A:389:HIS:O	1:A:390:HIS:CB	2.64	0.46
1:C:387:HIS:O	1:C:388:HIS:CB	2.64	0.46
1:C:326:SER:O	3:C:1419:HOH:O	2.21	0.45
1:A:157:MSE:CG	3:A:1182:HOH:O	2.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:228:ASN:O	3:C:997:HOH:O	2.21	0.44
1:B:192[B]:ARG:NH1	3:B:870:HOH:O	2.51	0.44
1:D:226:ASP:OD2	1:D:226:ASP:N	2.51	0.44
1:A:390:HIS:CG	1:A:391:HIS:CD2	3.05	0.43
1:C:192:ARG:NH1	3:C:510:HOH:O	2.52	0.43
1:C:58:TYR:OH	1:C:371:ASP:OD1	2.38	0.42
1:D:272:GLU:OE2	1:D:276:ARG:NH2	2.53	0.42
1:A:330:ARG:NH2	3:A:624:HOH:O	2.52	0.42
1:D:53:ASP:OD2	1:D:106:ARG:NH2	2.53	0.42
1:B:41:LYS:NZ	3:B:1386:HOH:O	2.52	0.41
1:B:389:HIS:C	1:B:389:HIS:CD2	2.94	0.41
1:A:82:ASP:OD1	1:A:83:ASP:N	2.53	0.41
1:D:58:TYR:OH	1:D:371:ASP:OD1	2.38	0.41
1:A:185:LYS:NZ	3:A:474:HOH:O	2.54	0.41
1:B:45:VAL:CG1	3:B:1560:HOH:O	2.67	0.41
1:C:330:ARG:NH2	3:C:581:HOH:O	2.52	0.41
1:D:119:HIS:CE1	1:D:327:ALA:O	2.74	0.41
1:B:253:ARG:NH2	1:B:255:GLN:OE1	2.54	0.41
1:B:350:TYR:OH	1:B:365:PHE:O	2.38	0.40
1:A:13:LYS:NZ	1:A:389:HIS:CD2	2.89	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:376:TYR:OH	1:A:376:TYR:OH[2.555]	1.80	0.40

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	393/392 (100%)	382 (97%)	9 (2%)	2 (0%)	38 22
1	B	388/392 (99%)	379 (98%)	9 (2%)	0	100 100
1	C	387/392 (99%)	380 (98%)	6 (2%)	1 (0%)	50 36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	378/392 (96%)	372 (98%)	6 (2%)	0	100	100
All	All	1546/1568 (99%)	1513 (98%)	30 (2%)	3 (0%)	56	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	387	HIS
1	C	388	HIS
1	A	390	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/308 (104%)	313 (98%)	6 (2%)	69	61
1	B	314/308 (102%)	307 (98%)	7 (2%)	64	54
1	C	313/308 (102%)	309 (99%)	4 (1%)	80	76
1	D	305/308 (99%)	298 (98%)	7 (2%)	63	52
All	All	1251/1232 (102%)	1227 (98%)	24 (2%)	69	61

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	LEU
1	A	122	THR
1	A	196	GLU
1	A	231	TRP
1	A	265	LEU
1	A	387	HIS
1	B	113	ASP
1	B	114	LEU
1	B	154	GLU
1	B	197	VAL
1	B	225	SER
1	B	231	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	389	HIS
1	C	4	ASN
1	C	5	ILE
1	C	231	TRP
1	C	389	HIS
1	D	78	LEU
1	D	87	LEU
1	D	106	ARG
1	D	120	LEU
1	D	226	ASP
1	D	231	TRP
1	D	295	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/392 (99%)	0.16	18 (4%) 31 31	6, 14, 36, 58	0
1	B	386/392 (98%)	0.29	19 (4%) 28 28	8, 17, 36, 51	0
1	C	386/392 (98%)	0.05	13 (3%) 43 43	6, 12, 29, 49	0
1	D	376/392 (95%)	0.50	27 (7%) 15 14	11, 25, 50, 61	0
All	All	1538/1568 (98%)	0.25	77 (5%) 28 27	6, 17, 41, 61	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	HIS	6.0
1	B	181	TRP	5.4
1	B	123	GLY	5.0
1	A	123	GLY	4.7
1	C	7	GLY	4.7
1	B	117	GLY	4.6
1	C	389	HIS	4.4
1	D	121	GLU	4.4
1	D	157	MSE	4.1
1	D	122	THR	3.9
1	A	83	ASP	3.9
1	B	6	THR	3.8
1	D	82	ASP	3.7
1	D	181[A]	TRP	3.7
1	C	123	GLY	3.7
1	D	79	ASP	3.6
1	A	387	HIS	3.6
1	D	123	GLY	3.6
1	A	124	ALA	3.5
1	B	388	HIS	3.5
1	A	2	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	84	ARG	3.4
1	B	389	HIS	3.4
1	B	226	ASP	3.3
1	D	124	ALA	3.2
1	C	5	ILE	3.2
1	C	83	ASP	3.2
1	A	122	THR	3.2
1	A	389	HIS	3.1
1	D	372	GLU	3.0
1	C	226	ASP	3.0
1	C	256	ASN	2.9
1	D	12	TRP	2.8
1	A	391	HIS	2.8
1	D	376[A]	TYR	2.8
1	D	314	GLY	2.8
1	D	343	GLU	2.8
1	A	157	MSE	2.7
1	B	122	THR	2.7
1	B	83	ASP	2.7
1	D	118	ALA	2.7
1	A	181	TRP	2.7
1	B	71	ASP	2.6
1	D	69	ALA	2.6
1	D	71	ASP	2.6
1	D	83	ASP	2.6
1	B	256	ASN	2.6
1	B	121	GLU	2.6
1	B	143	ALA	2.6
1	C	124	ALA	2.5
1	D	84	ARG	2.4
1	D	89	GLU	2.4
1	D	76	ARG	2.4
1	D	73	ILE	2.4
1	A	288	TRP	2.3
1	B	9	GLN	2.3
1	B	228	ASN	2.3
1	C	121	GLU	2.2
1	D	144	ASP	2.2
1	A	11	ASP	2.2
1	B	124	ALA	2.2
1	A	226	ASP	2.2
1	C	9	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	ASN	2.2
1	D	54	GLY	2.2
1	B	89	GLU	2.1
1	D	113	ASP	2.1
1	D	386	GLY	2.1
1	C	71	ASP	2.1
1	C	252	GLN	2.1
1	A	390	HIS	2.1
1	D	120	LEU	2.1
1	B	252	GLN	2.1
1	D	13	LYS	2.0
1	C	387	HIS	2.0
1	A	252	GLN	2.0
1	B	157[A]	MSE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MG	C	501	1/1	0.09	0.37	9,9,9,9	0
2	MG	B	501	1/1	0.09	-0.19	13,13,13,13	0
2	MG	D	501	1/1	0.08	-0.88	13,13,13,13	0
2	MG	A	501	1/1	0.05	-2.60	8,8,8,8	0
2	MG	B	502	1/1	0.06	-3.13	15,15,15,15	0
2	MG	C	502	1/1	0.04	-3.40	8,8,8,8	0
2	MG	A	502	1/1	0.04	-4.16	10,10,10,10	0
2	MG	D	502	1/1	0.02	-6.76	14,14,14,14	0

6.5 Other polymers ⓘ

There are no such residues in this entry.