



Full wwPDB X-ray Structure Validation Report

Sep 29, 2014 – 09:05 PM EDT

PDB ID : 4N4Z
Title : Trypanosoma brucei procathepsin B structure solved by Serial Microcrystallography using synchrotron radiation
Authors : Gati, C.; Bourenkov, G.; Klinge, M.; Rehders, D.; Stellato, F.; Oberthuer, D.; White, T.A.; Yevanov, O.; Sommer, B.P.; Mogk, S.; Duszenko, M.; Betzel, C.; Schneider, T.R.; Chapman, H.N.; Redecke, L.
Deposited on : 2013-10-08
Resolution : 3.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23828
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23828

1 Overall quality at a glance


The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1341 (3.40-3.20)
Clashscore	79885	1696 (3.40-3.20)
Ramachandran outliers	78287	1664 (3.40-3.20)
Sidechain outliers	78261	1662 (3.40-3.20)
RSRZ outliers	66119	1342 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	340	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2459 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine peptidase C (CPC).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2392	1504	413	456	19			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			39	22	2	15		

4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	124.42Å 124.42Å 53.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.10 – 3.30 87.98 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (88.10-3.30) 99.8 (87.98-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.199 , 0.251 0.203 , 0.251	Depositor DCC
R_{free} test set	305 reflections (4.74%)	DCC
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 13.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 6744 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2459	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/2464	0.69	3/3353 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	304	TRP	N-CA-C	6.08	127.42	111.00
1	A	107	CYS	CA-CB-SG	-5.39	104.29	114.00
1	A	304	TRP	CB-CA-C	-5.10	100.21	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2208	71	1
2	A	28	0	25	1	0
3	A	39	0	34	0	0
All	All	2459	0	2267	71	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 15.

All (71) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:39:ARG:O	1:A:43:LEU:HD22	1.85	0.76
1:A:83:ARG:HD3	1:A:235:TYR:CE1	2.24	0.73
1:A:195:HIS:HD2	1:A:197:LYS:H	1.33	0.72
1:A:304:TRP:O	1:A:305:ASN:HB2	1.90	0.70
1:A:57:GLN:HE21	2:A:401:NAG:HN2	1.39	0.70
1:A:144:VAL:HG13	1:A:144:VAL:O	1.95	0.67
1:A:52:TYR:O	1:A:57:GLN:HB3	1.96	0.66
1:A:68:ASN:HD21	1:A:259:VAL:HB	1.61	0.64
1:A:60:THR:HG22	1:A:62:ARG:H	1.61	0.64
1:A:110:ILE:HB	1:A:111:PRO:HD3	1.79	0.64
1:A:60:THR:HB	1:A:63:GLU:HG3	1.80	0.62
1:A:228:ASN:HD22	1:A:230:ARG:NH2	1.96	0.62
1:A:60:THR:HG22	1:A:62:ARG:N	2.15	0.62
1:A:175:SER:HA	1:A:231:SER:O	2.02	0.60
1:A:200:ASN:HB2	1:A:202:TYR:CE2	2.36	0.60
1:A:174:PHE:CE1	1:A:231:SER:HB3	2.36	0.60
1:A:56:MET:O	1:A:59:ILE:HG22	2.02	0.60
1:A:257:PHE:HB3	1:A:325:ILE:HD13	1.83	0.59
1:A:193:SER:O	1:A:204:PRO:HA	2.04	0.58
1:A:304:TRP:O	1:A:305:ASN:CB	2.49	0.58
1:A:250:ARG:NH1	1:A:333:ILE:HD11	2.19	0.57
1:A:237:LEU:HD21	1:A:246:GLU:HG2	1.86	0.57
1:A:316:ILE:HG12	1:A:317:ARG:H	1.70	0.56
1:A:119:CYS:HB3	1:A:121:SER:OG	2.05	0.55
1:A:118:ALA:HB3	1:A:210:PHE:CD2	2.42	0.55
1:A:195:HIS:HD2	1:A:197:LYS:N	2.04	0.55
1:A:191:HIS:HB3	1:A:202:TYR:CD1	2.42	0.54
1:A:68:ASN:ND2	1:A:259:VAL:HB	2.22	0.54
1:A:238:GLN:O	1:A:242:ASP:HB2	2.09	0.53
1:A:33:SER:HB2	1:A:36:PHE:H	1.74	0.53
1:A:290:GLY:O	1:A:318:ARG:NH2	2.43	0.52
1:A:113:ILE:HG21	1:A:305:ASN:O	2.10	0.52
1:A:39:ARG:HG3	1:A:43:LEU:HD21	1.93	0.51
1:A:145:HIS:O	1:A:182:ASP:N	2.41	0.51
1:A:44:ASN:ND2	1:A:48:TRP:O	2.43	0.51
1:A:174:PHE:CD1	1:A:231:SER:HB3	2.47	0.50
1:A:28:ASP:HA	1:A:60:THR:HG23	1.94	0.50
1:A:39:ARG:HG3	1:A:43:LEU:CD2	2.43	0.48
1:A:204:PRO:HD2	1:A:207:GLN:HE21	1.78	0.48
1:A:160:ASP:OD1	1:A:163:ASN:ND2	2.38	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:237:LEU:HD21	1:A:246:GLU:CB	2.44	0.47
1:A:144:VAL:O	1:A:144:VAL:CG1	2.62	0.47
1:A:134:ARG:NH1	1:A:331:ALA:HB1	2.30	0.46
1:A:189:PHE:CD1	1:A:210:PHE:CD1	3.04	0.46
1:A:320:SER:O	1:A:321:SER:HB3	2.15	0.46
1:A:123:TRP:CG	1:A:165:GLY:HA3	2.51	0.45
1:A:237:LEU:HD21	1:A:246:GLU:CG	2.46	0.45
1:A:308:TRP:HZ3	1:A:314:PHE:HB3	1.81	0.45
1:A:126:ALA:HB3	1:A:254:GLU:OE1	2.16	0.44
1:A:195:HIS:CD2	1:A:197:LYS:H	2.24	0.44
1:A:135:PHE:CZ	1:A:229:TYR:HB3	2.53	0.44
1:A:193:SER:HB2	1:A:202:TYR:HB2	2.00	0.44
1:A:208:PHE:HB3	1:A:210:PHE:CE2	2.52	0.44
1:A:278:TYR:CZ	1:A:280:GLY:HA2	2.53	0.43
1:A:160:ASP:HB3	1:A:163:ASN:HB2	1.99	0.43
1:A:69:GLY:O	1:A:165:GLY:N	2.44	0.43
1:A:187:TYR:HA	1:A:188:PRO:HD3	1.83	0.43
1:A:63:GLU:O	1:A:66:ARG:HB3	2.18	0.43
1:A:92:ALA:HA	1:A:93:PRO:HD3	1.89	0.43
1:A:118:ALA:HB3	1:A:210:PHE:HD2	1.82	0.43
1:A:71:ILE:HG13	1:A:71:ILE:O	2.19	0.43
1:A:191:HIS:CB	1:A:202:TYR:CD1	3.02	0.42
1:A:316:ILE:CG1	1:A:317:ARG:H	2.31	0.42
1:A:308:TRP:CZ3	1:A:314:PHE:HB3	2.53	0.42
1:A:135:PHE:HA	1:A:139:GLY:HA3	2.02	0.42
1:A:90:ALA:HA	1:A:245:ARG:HD3	2.03	0.41
1:A:107:CYS:HA	1:A:108:PRO:HD3	1.68	0.41
1:A:257:PHE:HB3	1:A:325:ILE:CD1	2.50	0.41
1:A:153:ALA:HB2	1:A:213:PRO:HG2	2.02	0.41
1:A:106:ASN:O	1:A:142:GLN:NE2	2.37	0.41
1:A:228:ASN:HD22	1:A:230:ARG:HH21	1.67	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:28:ASP:O	1:A:82:LYS:NZ[4_545]	1.77	0.43

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/340 (90%)	278 (91%)	27 (9%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/276 (92%)	248 (98%)	5 (2%)	68	92

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	VAL
1	A	119	CYS
1	A	157	ASP
1	A	293	ASN
1	A	310	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	106	ASN
1	A	195	HIS
1	A	228	ASN
1	A	282	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	401	1,2	12,14,15	1.06	2 (16%)	15,19,21	1.38	2 (13%)
2	NAG	A	402	2	12,14,15	1.10	2 (16%)	15,19,21	1.56	4 (26%)
3	NAG	A	403	1,3	12,14,15	1.37	1 (8%)	15,19,21	1.45	3 (20%)
3	NAG	A	404	3	12,14,15	0.89	1 (8%)	15,19,21	1.15	1 (6%)
3	BMA	A	405	3	10,11,12	0.62	0	11,15,17	1.30	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	402	2	-	0/6/23/26	0/1/1/1
3	NAG	A	403	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	404	3	-	0/6/23/26	0/1/1/1
3	BMA	A	405	3	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	NAG	O5-C5	-3.57	1.39	1.45
3	A	404	NAG	O5-C5	-2.76	1.41	1.45
2	A	401	NAG	O5-C5	-2.57	1.41	1.45
2	A	402	NAG	O5-C5	-2.47	1.41	1.45
2	A	401	NAG	C2-N2	-2.33	1.43	1.46
2	A	402	NAG	C3-C2	2.26	1.56	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	NAG	O5-C5-C6	-3.55	103.25	106.98
2	A	401	NAG	O5-C5-C6	3.25	110.39	106.98
2	A	402	NAG	C4-C3-C2	3.09	117.23	110.74
2	A	401	NAG	C3-C2-N2	-3.09	106.99	111.62
3	A	405	BMA	C3-C4-C5	2.83	115.25	110.17
2	A	402	NAG	C2-N2-C7	2.47	126.38	123.39
2	A	402	NAG	O5-C5-C6	2.45	109.56	106.98
3	A	403	NAG	C3-C4-C5	2.44	114.55	110.17
3	A	404	NAG	O5-C5-C6	2.37	109.47	106.98
2	A	402	NAG	O5-C5-C4	-2.24	107.81	110.65
3	A	405	BMA	O5-C5-C6	2.17	109.26	106.98
3	A	403	NAG	O4-C4-C3	-2.16	105.52	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/340 (90%)	0.07	0 100 100	22, 35, 57, 79	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	401	14/15	0.35	0.93	54,61,66,70	0
2	NAG	A	402	14/15	0.39	0.88	46,57,66,67	9
3	NAG	A	404	14/15	0.30	0.45	42,49,53,55	0
3	NAG	A	403	14/15	0.24	-0.07	40,42,47,50	0
3	BMA	A	405	11/12	0.32	-	49,53,56,59	0

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.