



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 02:37 PM GMT

PDB ID : 1NGJ
Title : STRUCTURAL BASIS OF THE 70-KILODALTON HEAT SHOCK COGNATE PROTEIN ATP HYDROLYTIC ACTIVITY, II. STRUCTURE OF THE ACTIVE SITE WITH ADP OR ATP BOUND TO WILD TYPE AND MUTANT ATPASE FRAGMENT
Authors : Flaherty, K.M.; Wilbanks, S.M.; Deluca-Flaherty, C.; Mckay, D.B.
Deposited on : 1994-05-17
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	386	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3153 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEAT-SHOCK COGNATE 70 kD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	0	0
			2958	1857	520	573	8			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONICACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	163	Total 163	O 163	0	0

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	145.30Å 65.00Å 46.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.194 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3153	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/3004	0.62	0/4055

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2958	0	2966	50	0
2	A	1	0	0	0	0
3	A	31	0	13	3	0
4	A	163	0	0	3	0
All	All	3153	0	2979	50	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (50) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:310:PHE:O	1:A:313:THR:HB	1.93	0.69
1:A:244:GLU:OE2	1:A:295:THR:HG21	1.94	0.67
1:A:46:ASP:O	1:A:108:LYS:HG2	1.95	0.66
1:A:60:ALA:HA	1:A:261:ARG:HD3	1.81	0.62
1:A:331:ILE:HD11	1:A:334:ILE:CG1	2.30	0.62
1:A:11:LEU:HG	1:A:124:LEU:HD11	1.81	0.61
1:A:151:ASN:H	1:A:154:GLN:HE21	1.47	0.61
1:A:21:PHE:CZ	1:A:24:GLY:HA2	2.36	0.61
1:A:250:LYS:NZ	1:A:250:LYS:HB3	2.19	0.57
1:A:9:ILE:HG12	1:A:18:VAL:HG13	1.87	0.56
1:A:269:ARG:HH11	1:A:269:ARG:HG2	1.71	0.55
1:A:269:ARG:HG3	4:A:598:HOH:O	2.07	0.55
1:A:283:GLU:HG2	1:A:294:TYR:CE1	2.42	0.55
1:A:257:LYS:HD2	1:A:257:LYS:H	1.73	0.54
1:A:118:GLU:HA	1:A:164:ILE:HD11	1.90	0.53
1:A:201:GLY:HA3	3:A:486:ANP:HNB1	1.74	0.53
1:A:11:LEU:HG	1:A:124:LEU:CD1	2.39	0.53
1:A:331:ILE:HD11	1:A:334:ILE:HG13	1.91	0.52
1:A:77:ARG:O	1:A:80:ASP:HB2	2.09	0.52
1:A:146:VAL:HG22	1:A:150:PHE:CD2	2.44	0.51
1:A:202:GLY:H	3:A:486:ANP:HNB1	1.57	0.51
1:A:246:LYS:O	1:A:250:LYS:HA	2.11	0.50
1:A:379:ILE:HG13	1:A:380:LEU:N	2.26	0.50
1:A:160:ASP:O	1:A:164:ILE:HG23	2.12	0.49
1:A:178:ALA:O	1:A:372:GLY:HA3	2.12	0.49
1:A:193:ARG:NH2	4:A:653:HOH:O	2.40	0.48
1:A:331:ILE:HD11	1:A:334:ILE:HG12	1.95	0.48
1:A:297:ILE:HD12	1:A:297:ILE:H	1.78	0.48
1:A:87:MET:HA	1:A:90:TRP:CE3	2.49	0.48
1:A:190:GLY:O	1:A:191:ALA:HB3	2.14	0.48
1:A:107:TYR:HB3	1:A:112:LYS:HD2	1.96	0.47
1:A:175:GLU:N	1:A:176:PRO:HD2	2.30	0.46
1:A:284:ILE:HG21	1:A:287:LEU:HD13	1.97	0.45
1:A:235:ASN:OD1	1:A:264:ARG:NH1	2.50	0.45
1:A:300:ALA:O	1:A:304:GLU:HB3	2.16	0.45
1:A:202:GLY:HA2	3:A:486:ANP:H4'	1.99	0.45
1:A:29:ILE:HD13	1:A:131:ALA:HA	1.99	0.44
1:A:384:LYS:O	1:A:384:LYS:HG2	2.18	0.43
1:A:269:ARG:NH1	1:A:269:ARG:HG2	2.33	0.43
1:A:310:PHE:CD1	1:A:346:ILE:HD11	2.53	0.43
1:A:192:GLU:HG2	1:A:213:GLU:HB2	2.00	0.42
1:A:315:ASP:HB2	1:A:316:PRO:HD3	2.00	0.42
1:A:236:ARG:HG3	1:A:306:ASN:ND2	2.33	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:364:ASN:HA	1:A:365:PRO:HD3	1.93	0.41
1:A:137:LYS:HD2	1:A:138:THR:H	1.84	0.41
1:A:209:ILE:HD11	1:A:321:LEU:HD13	2.02	0.41
1:A:200:LEU:HG	1:A:340:SER:HB2	2.02	0.40
1:A:76:ARG:NH1	4:A:623:HOH:O	2.54	0.40
1:A:103:VAL:O	1:A:113:SER:HA	2.22	0.40
1:A:205:PHE:O	1:A:224:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	380/386 (98%)	369 (97%)	10 (3%)	1 (0%)	50 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	316/320 (99%)	281 (89%)	35 (11%)	9 5

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	11	LEU
1	A	18	VAL
1	A	32	ASP
1	A	38	THR
1	A	50	LEU
1	A	58	GLN
1	A	73	LEU
1	A	79	ASP
1	A	80	ASP
1	A	124	LEU
1	A	146	VAL
1	A	156	GLN
1	A	164	ILE
1	A	167	LEU
1	A	170	LEU
1	A	171	ARG
1	A	214	ASP
1	A	228	LEU
1	A	250	LYS
1	A	255	GLU
1	A	274	LEU
1	A	279	GLN
1	A	283	GLU
1	A	297	ILE
1	A	301	ARG
1	A	304	GLU
1	A	309	LEU
1	A	313	THR
1	A	321	LEU
1	A	331	ILE
1	A	350	LEU
1	A	358	GLU
1	A	359	LEU
1	A	380	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	57	ASN
1	A	58	GLN
1	A	141	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	154	GLN
1	A	194	ASN
1	A	227	HIS
1	A	306	ASN
1	A	355	ASN
1	A	376	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ANP	A	486	2	33,33,33	2.65	7 (21%)	51,52,52	2.04	12 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	486	2	-	1/18/38/38	0/1/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	486	ANP	PB-N3B	-11.45	1.54	1.64
3	A	486	ANP	PG-N3B	-6.57	1.58	1.64
3	A	486	ANP	C8-N9	3.51	1.41	1.36
3	A	486	ANP	PA-O3A	-2.55	1.55	1.59
3	A	486	ANP	PG-O2G	-2.26	1.48	1.55
3	A	486	ANP	O4'-C1'	2.23	1.44	1.41
3	A	486	ANP	PB-O2B	-2.20	1.48	1.55

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	486	ANP	O1G-PG-N3B	-6.20	102.46	111.83
3	A	486	ANP	C8-N9-C1'	4.89	136.01	126.38
3	A	486	ANP	O3A-PB-N3B	-4.75	93.42	106.59
3	A	486	ANP	C1'-N9-C4	-4.36	119.11	126.64
3	A	486	ANP	O2B-PB-O1B	4.13	119.41	109.89
3	A	486	ANP	O1B-PB-N3B	-3.56	106.45	111.83
3	A	486	ANP	O3G-PG-O2G	3.26	117.00	107.66
3	A	486	ANP	C8-N9-C4	-2.94	104.66	106.90
3	A	486	ANP	PA-O3A-PB	2.76	141.24	131.81
3	A	486	ANP	O3'-C3'-C4'	-2.58	103.48	111.08
3	A	486	ANP	O4'-C4'-C5'	-2.55	100.25	109.36
3	A	486	ANP	O4'-C1'-N9	2.54	110.80	108.44

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	486	ANP	O1B-PB-N3B-PG

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.