



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 03:54 PM GMT

PDB ID : 3NGT
Title : Structure of Leishmania NDKb complexed with AMP.
Authors : Trindade, D.M.; Sousa, T.A.C.B.; Tonoli, C.C.C.; Santos, C.R.; Arni, R.K.;
Ward, R.J.; Oliveira, A.H.C.; Murakami, M.T.
Deposited on : 2010-06-13
Resolution : 2.57 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1891 (2.60-2.56)
Clashscore	79885	2358 (2.60-2.56)
Ramachandran outliers	78287	2316 (2.60-2.56)
Sidechain outliers	78261	2316 (2.60-2.56)
RSRZ outliers	66119	1891 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	151	
1	B	151	
1	C	151	
1	D	151	
1	E	151	
1	F	151	
1	G	151	
1	H	151	
1	I	151	
1	J	151	
1	K	151	
1	L	151	
1	M	151	
1	N	151	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	AMP	A	152	-	X
2	AMP	E	152	-	X
2	AMP	M	152	-	X

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 16728 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	150	Total	C	N	O	S	27	1	0
			1171	744	203	219	5			
1	B	148	Total	C	N	O	S	24	2	0
			1159	739	200	214	6			
1	C	148	Total	C	N	O	S	26	1	0
			1156	737	200	214	5			
1	D	150	Total	C	N	O	S	15	0	0
			1165	741	202	218	4			
1	E	150	Total	C	N	O	S	33	1	0
			1168	743	202	218	5			
1	F	150	Total	C	N	O	S	31	1	0
			1168	743	202	218	5			
1	G	149	Total	C	N	O	S	44	0	0
			1159	738	201	216	4			
1	H	147	Total	C	N	O	S	51	1	0
			1148	731	199	214	4			
1	I	150	Total	C	N	O	S	42	1	0
			1168	743	202	218	5			
1	J	148	Total	C	N	O	S	15	0	0
			1149	733	200	212	4			
1	K	149	Total	C	N	O	S	15	1	0
			1165	741	202	217	5			
1	L	125	Total	C	N	O	S	35	0	0
			960	608	170	179	3			
1	M	149	Total	C	N	O	S	21	1	0
			1165	741	202	217	5			
1	N	150	Total	C	N	O	S	20	1	0
			1171	744	203	219	5			

- Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	G	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	H	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	I	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	J	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	K	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	M	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	N	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total 28	O 28	0	0
3	B	39	Total 39	O 39	0	0
3	C	30	Total 30	O 30	0	0
3	D	26	Total 26	O 26	0	0
3	E	23	Total 23	O 23	0	0
3	F	30	Total 30	O 30	0	0
3	G	22	Total 22	O 22	0	0
3	H	18	Total 18	O 18	0	0
3	I	16	Total 16	O 16	0	0
3	J	22	Total 22	O 22	0	0
3	K	25	Total 25	O 25	0	0
3	L	10	Total 10	O 10	0	0
3	M	30	Total 30	O 30	0	0
3	N	38	Total 38	O 38	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleoside diphosphate kinase

Chain A:




- Molecule 1: Nucleoside diphosphate kinase

Chain B:




- Molecule 1: Nucleoside diphosphate kinase

Chain C:




- Molecule 1: Nucleoside diphosphate kinase

Chain D:




- Molecule 1: Nucleoside diphosphate kinase

Chain E:




- Molecule 1: Nucleoside diphosphate kinase

Chain F:




- Molecule 1: Nucleoside diphosphate kinase

Chain G:




- Molecule 1: Nucleoside diphosphate kinase

Chain H:




- Molecule 1: Nucleoside diphosphate kinase

Chain I:




- Molecule 1: Nucleoside diphosphate kinase

Chain J: 




- Molecule 1: Nucleoside diphosphate kinase

Chain K: 




- Molecule 1: Nucleoside diphosphate kinase

Chain L: 




- Molecule 1: Nucleoside diphosphate kinase

Chain M: 



- Molecule 1: Nucleoside diphosphate kinase

Chain N: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	233.95Å 233.95Å 265.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	161.00 – 2.57 37.26 – 2.57	Depositor EDS
% Data completeness (in resolution range)	97.4 (161.00-2.57) 97.4 (37.26-2.57)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.237 , 0.314 0.227 , 0.292	Depositor DCC
R_{free} test set	4338 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	42.6	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	3 of 86167 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16728	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.05 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.4726e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	4/1198 (0.3%)	1.04	5/1619 (0.3%)
1	B	1.10	4/1192 (0.3%)	0.98	5/1611 (0.3%)
1	C	0.97	3/1186 (0.3%)	0.92	3/1603 (0.2%)
1	D	0.95	1/1192 (0.1%)	0.90	7/1611 (0.4%)
1	E	1.12	4/1198 (0.3%)	0.86	4/1619 (0.2%)
1	F	1.02	5/1198 (0.4%)	1.08	10/1619 (0.6%)
1	G	1.16	8/1186 (0.7%)	0.91	4/1603 (0.2%)
1	H	0.93	4/1177 (0.3%)	1.04	7/1589 (0.4%)
1	I	0.98	6/1198 (0.5%)	0.90	2/1619 (0.1%)
1	J	0.86	1/1176 (0.1%)	0.86	2/1591 (0.1%)
1	K	1.05	3/1192 (0.3%)	0.90	3/1611 (0.2%)
1	L	0.73	1/979 (0.1%)	0.84	3/1322 (0.2%)
1	M	0.95	1/1192 (0.1%)	0.94	3/1611 (0.2%)
1	N	0.99	4/1198 (0.3%)	0.85	1/1619 (0.1%)
All	All	0.99	49/16462 (0.3%)	0.93	59/22247 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1
1	H	1	1
1	K	1	0
All	All	2	2

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	126	GLU	CB-CG	20.25	1.90	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	45	GLU	CB-CG	-19.65	1.14	1.52
1	K	45	GLU	CA-CB	-18.73	1.12	1.53
1	E	99	GLN	CG-CD	-18.49	1.08	1.51
1	I	3	SER	CA-CB	-18.00	1.25	1.52

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	ASP	CB-CG-OD1	-15.46	104.38	118.30
1	H	127	ARG	CD-NE-CZ	15.32	145.06	123.60
1	A	136	ASP	CB-CG-OD2	15.27	132.04	118.30
1	I	2	SER	N-CA-CB	13.15	130.22	110.50
1	K	45	GLU	N-CA-CB	11.87	131.97	110.60

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	H	126	GLU	CA
1	K	45	GLU	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	61	PRO	Peptide
1	H	56	SER	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1171	0	1156	39	0
1	B	1159	0	1152	21	0
1	C	1156	0	1147	37	0
1	D	1165	0	1152	26	0
1	E	1168	0	1157	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1168	0	1157	24	0
1	G	1159	0	1147	37	0
1	H	1148	0	1135	29	0
1	I	1168	0	1157	34	0
1	J	1149	0	1141	33	0
1	K	1165	0	1151	32	0
1	L	960	0	955	27	0
1	M	1165	0	1151	25	0
1	N	1171	0	1156	26	0
2	A	23	0	12	3	0
2	B	23	0	12	2	0
2	C	23	0	12	4	0
2	D	23	0	12	1	0
2	E	23	0	12	3	0
2	F	23	0	12	1	0
2	G	23	0	12	2	0
2	H	23	0	12	0	0
2	I	23	0	12	2	0
2	J	23	0	12	2	0
2	K	23	0	12	2	0
2	M	23	0	12	4	0
2	N	23	0	12	2	0
3	A	28	0	0	4	0
3	B	39	0	0	1	0
3	C	30	0	0	4	0
3	D	26	0	0	1	0
3	E	23	0	0	3	0
3	F	30	0	0	1	0
3	G	22	0	0	3	0
3	H	18	0	0	1	0
3	I	16	0	0	1	0
3	J	22	0	0	0	0
3	K	25	0	0	3	0
3	L	10	0	0	1	0
3	M	30	0	0	0	0
3	N	38	0	0	2	0
All	All	16728	0	16070	375	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

The worst 5 of 375 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:M:90:LEU:HD12	1:M:103:ILE:HD12	1.33	1.08
1:D:2:SER:N	1:D:3:SER:HA	1.65	1.07
1:L:90:LEU:HG	1:L:116:CYS:SG	1.96	1.04
1:J:55:CYS:O	1:J:55:CYS:SG	2.19	1.01
1:C:38:LYS:HD2	3:C:355:HOH:O	1.56	1.01

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/151 (99%)	144 (97%)	4 (3%)	1 (1%)	30	56
1	B	148/151 (98%)	137 (93%)	8 (5%)	3 (2%)	11	21
1	C	147/151 (97%)	135 (92%)	10 (7%)	2 (1%)	16	31
1	D	148/151 (98%)	137 (93%)	11 (7%)	0	100	100
1	E	149/151 (99%)	142 (95%)	6 (4%)	1 (1%)	30	56
1	F	149/151 (99%)	134 (90%)	12 (8%)	3 (2%)	11	21
1	G	147/151 (97%)	135 (92%)	10 (7%)	2 (1%)	16	31
1	H	144/151 (95%)	127 (88%)	12 (8%)	5 (4%)	6	8
1	I	149/151 (99%)	141 (95%)	6 (4%)	2 (1%)	18	34
1	J	146/151 (97%)	140 (96%)	2 (1%)	4 (3%)	8	12
1	K	148/151 (98%)	142 (96%)	6 (4%)	0	100	100
1	L	121/151 (80%)	109 (90%)	7 (6%)	5 (4%)	4	5
1	M	148/151 (98%)	142 (96%)	6 (4%)	0	100	100
1	N	149/151 (99%)	143 (96%)	5 (3%)	1 (1%)	30	56
All	All	2042/2114 (97%)	1908 (93%)	105 (5%)	29 (1%)	16	31

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	123	GLU
1	F	3	SER
1	F	137	GLU
1	H	4	GLU
1	H	51	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/126 (100%)	114 (90%)	12 (10%)	12	22
1	B	125/126 (99%)	110 (88%)	15 (12%)	7	13
1	C	124/126 (98%)	111 (90%)	13 (10%)	10	17
1	D	125/126 (99%)	116 (93%)	9 (7%)	21	39
1	E	126/126 (100%)	116 (92%)	10 (8%)	18	33
1	F	126/126 (100%)	112 (89%)	14 (11%)	9	15
1	G	124/126 (98%)	111 (90%)	13 (10%)	10	17
1	H	123/126 (98%)	105 (85%)	18 (15%)	5	8
1	I	126/126 (100%)	110 (87%)	16 (13%)	6	12
1	J	123/126 (98%)	108 (88%)	15 (12%)	7	13
1	K	125/126 (99%)	118 (94%)	7 (6%)	30	54
1	L	103/126 (82%)	88 (85%)	15 (15%)	5	8
1	M	125/126 (99%)	112 (90%)	13 (10%)	10	18
1	N	126/126 (100%)	112 (89%)	14 (11%)	9	15
All	All	1727/1764 (98%)	1543 (89%)	184 (11%)	10	17

5 of 184 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	119	SER
1	H	134	LYS
1	N	34	LEU
1	G	147	SER
1	H	65	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	50	HIS
1	I	41	GLN
1	L	41	GLN
1	G	50	HIS
1	K	94	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	AMP	A	152	-	25,25,25	1.38	3 (12%)	38,38,38	2.32	9 (23%)
2	AMP	B	152	-	25,25,25	1.24	3 (12%)	38,38,38	3.14	10 (26%)
2	AMP	C	152	-	25,25,25	1.21	2 (8%)	38,38,38	2.03	9 (23%)
2	AMP	D	152	-	25,25,25	1.22	2 (8%)	38,38,38	2.31	9 (23%)
2	AMP	E	152	-	25,25,25	1.27	3 (12%)	38,38,38	2.14	12 (31%)
2	AMP	F	152	-	25,25,25	1.28	3 (12%)	38,38,38	2.01	10 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AMP	G	152	-	25,25,25	1.20	2 (8%)	38,38,38	2.10	10 (26%)
2	AMP	H	152	-	25,25,25	1.17	2 (8%)	38,38,38	1.91	7 (18%)
2	AMP	I	152	-	25,25,25	1.21	3 (12%)	38,38,38	2.01	11 (28%)
2	AMP	J	152	-	25,25,25	1.27	2 (8%)	38,38,38	2.08	8 (21%)
2	AMP	K	152	-	25,25,25	1.20	2 (8%)	38,38,38	2.06	9 (23%)
2	AMP	M	152	-	25,25,25	1.42	4 (16%)	38,38,38	1.92	8 (21%)
2	AMP	N	152	-	25,25,25	1.12	2 (8%)	38,38,38	2.11	8 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	A	152	-	-	0/10/26/26	0/1/3/3
2	AMP	B	152	-	-	0/10/26/26	0/1/3/3
2	AMP	C	152	-	-	0/10/26/26	0/1/3/3
2	AMP	D	152	-	-	0/10/26/26	0/1/3/3
2	AMP	E	152	-	-	0/10/26/26	0/1/3/3
2	AMP	F	152	-	-	0/10/26/26	0/1/3/3
2	AMP	G	152	-	-	0/10/26/26	0/1/3/3
2	AMP	H	152	-	-	0/10/26/26	0/1/3/3
2	AMP	I	152	-	-	0/10/26/26	0/1/3/3
2	AMP	J	152	-	-	0/10/26/26	0/1/3/3
2	AMP	K	152	-	-	0/10/26/26	0/1/3/3
2	AMP	M	152	-	-	0/10/26/26	0/1/3/3
2	AMP	N	152	-	-	0/10/26/26	0/1/3/3

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	152	AMP	C5-C4	4.02	1.49	1.40
2	E	152	AMP	C5-C4	3.85	1.49	1.40
2	J	152	AMP	C5-C4	3.83	1.49	1.40
2	F	152	AMP	C5-C4	3.77	1.49	1.40
2	B	152	AMP	C5-C4	3.70	1.48	1.40

The worst 5 of 120 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	152	AMP	O4'-C1'-N9	13.85	121.32	108.44
2	B	152	AMP	C4'-O4'-C1'	-8.64	100.36	109.75
2	K	152	AMP	N3-C2-N1	-7.58	122.37	128.71
2	E	152	AMP	N3-C2-N1	-7.34	122.58	128.71
2	A	152	AMP	C4'-O4'-C1'	-7.30	101.81	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	150/151 (99%)	-0.02	4 (2%) 52 50	20, 33, 46, 56	10 (6%)
1	B	148/151 (98%)	-0.20	0 100 100	19, 32, 44, 49	9 (6%)
1	C	148/151 (98%)	0.04	2 (1%) 72 73	20, 35, 46, 58	8 (5%)
1	D	150/151 (99%)	-0.08	3 (2%) 62 61	23, 34, 50, 59	8 (5%)
1	E	150/151 (99%)	0.04	2 (1%) 74 75	23, 37, 51, 57	10 (6%)
1	F	150/151 (99%)	-0.03	2 (1%) 74 75	20, 36, 52, 62	10 (6%)
1	G	149/151 (98%)	0.08	5 (3%) 43 41	28, 44, 64, 67	14 (9%)
1	H	147/151 (97%)	0.24	6 (4%) 35 33	29, 47, 68, 75	13 (8%)
1	I	150/151 (99%)	0.14	4 (2%) 52 50	29, 44, 58, 63	14 (9%)
1	J	148/151 (98%)	0.05	1 (0%) 84 87	26, 38, 53, 55	8 (5%)
1	K	149/151 (98%)	-0.16	1 (0%) 84 87	26, 38, 48, 59	7 (4%)
1	L	125/151 (82%)	0.52	10 (8%) 12 10	34, 53, 68, 72	9 (7%)
1	M	149/151 (98%)	-0.22	0 100 100	21, 30, 44, 48	8 (5%)
1	N	150/151 (99%)	0.04	2 (1%) 74 75	19, 32, 50, 57	8 (5%)
All	All	2063/2114 (97%)	0.03	42 (2%) 62 61	19, 38, 58, 75	136 (6%)

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	3	SER	4.1
1	H	59	PHE	4.0
1	L	2	SER	3.8
1	C	151	GLU	3.5
1	A	122	VAL	3.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	AMP	E	152	23/23	0.26	3.57	55,58,60,63	0
2	AMP	A	152	23/23	0.27	3.19	48,52,57,57	0
2	AMP	M	152	23/23	0.26	2.08	49,58,65,65	0
2	AMP	F	152	23/23	0.22	1.57	59,69,74,75	0
2	AMP	H	152	23/23	0.31	1.51	84,86,87,88	0
2	AMP	K	152	23/23	0.24	1.48	59,65,72,72	0
2	AMP	C	152	23/23	0.25	1.42	56,60,61,64	0
2	AMP	D	152	23/23	0.23	1.41	66,67,69,70	0
2	AMP	B	152	23/23	0.23	1.36	63,68,73,74	0
2	AMP	J	152	23/23	0.23	1.21	65,67,70,70	0
2	AMP	I	152	23/23	0.25	1.06	73,75,80,81	0
2	AMP	G	152	23/23	0.24	0.99	78,80,84,84	0
2	AMP	N	152	23/23	0.23	0.79	67,71,79,80	0

6.5 Other polymers ⓘ

There are no such residues in this entry.