



Full wwPDB X-ray Structure Validation Report

Nov 11, 2014 – 06:30 PM EST

PDB ID : 4NQE
Title : Crystal structure of TCR-MR1 ternary complex bound to 5-(2-oxoethylidene amino)-6-D-ribitylaminouracil
Authors : Birkinshaw, R.W.; Rossjohn, J.
Deposited on : 2013-11-25
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable24103
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.1.3
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable24103

1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	271	
1	C	271	
2	B	99	
2	F	99	
3	D	203	
3	G	203	
4	E	245	
4	H	245	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13326 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2152	1380	373	388	11			
1	C	249	Total	C	N	O	S	0	1	0
			2044	1307	358	368	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP Q95460
A	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460
C	0	MET	-	EXPRESSION TAG	UNP Q95460
C	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	96	Total	C	N	O	S	0	0	0
			769	494	131	142	2			
2	F	97	Total	C	N	O	S	0	0	0
			766	492	130	142	2			


- Molecule 3 is a protein called TCR alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	200	Total	C	N	O	S	0	0	0
			1546	978	248	311	9			
3	G	189	Total	C	N	O	S	0	1	0
			1452	927	231	285	9			

- Molecule 4 is a protein called TCR beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	243	Total	C	N	O	S	0	2	0
			1913	1204	333	367	9			
4	H	231	Total	C	N	O	S	0	0	0
			1754	1109	298	338	9			

- Molecule 5 is 1-DEOXY-1-({2,6-DIOXO-5-[(E)-(2-OXOETHYLIDENE)AMINO]-1,2,3,6-TETRAHYDROPYRIMIDIN-4-YL}AMINO)-D-RIBITOL (three-letter code: 2L4) (formula: C₁₁H₁₆N₄O₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			21	11	4	6		
5	C	1	Total	C	N	O	0	0
			21	11	4	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	159	Total	O	0	0
			159	159		
6	B	68	Total	O	0	0
			68	68		
6	C	162	Total	O	0	0
			162	162		
6	D	133	Total	O	0	0
			133	133		
6	E	197	Total	O	0	0
			197	197		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	40	Total	O	0	0
			40	40		
6	G	69	Total	O	0	0
			69	69		
6	H	60	Total	O	0	0
			60	60		




• Molecule 4: TCR beta

Chain E: 



• Molecule 4: TCR beta

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	218.11Å 70.60Å 143.86Å 90.00° 104.63° 90.00°	Depositor
Resolution (Å)	42.00 – 2.10 75.42 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.6 (42.00-2.10) 98.5 (75.42-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.10Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.182 , 0.222 0.196 , 0.239	Depositor DCC
R_{free} test set	6144 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	28.5	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 122098 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13326	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2L4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2218	0.64	0/3017
1	C	0.53	0/2106	0.63	0/2859
2	B	0.47	0/792	0.64	0/1080
2	F	0.48	0/789	0.63	0/1078
3	D	0.54	0/1581	0.67	0/2145
3	G	0.51	0/1487	0.67	0/2018
4	E	0.53	0/1971	0.67	0/2684
4	H	0.45	0/1799	0.66	0/2456
All	All	0.51	0/12743	0.65	0/17337

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2152	0	2028	10	0
1	C	2044	0	1926	8	0
2	B	769	0	715	1	0
2	F	766	0	700	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1546	0	1456	3	0
3	G	1452	0	1360	13	0
4	E	1913	0	1799	8	0
4	H	1754	0	1621	8	0
5	A	21	0	0	0	0
5	C	21	0	0	0	0
6	A	159	0	0	2	0
6	B	68	0	0	0	0
6	C	162	0	0	1	0
6	D	133	0	0	0	0
6	E	197	0	0	1	0
6	F	40	0	0	0	0
6	G	69	0	0	0	0
6	H	60	0	0	0	0
All	All	13326	0	11605	47	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (47) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:D:1:GLY:HA2	3:D:27:SER:H	1.47	0.78
3:G:1:GLY:N	3:G:27:SER:H	1.89	0.70
4:E:210:ARG:HH21	4:E:212:GLN:HE22	1.42	0.68
1:C:111:GLN:HG2	1:C:121:ILE:HG12	1.77	0.67
3:D:78:LEU:HG	3:D:107:ILE:HD12	1.79	0.64
1:C:259:GLU:HG3	1:C:264:HIS:CE1	2.34	0.62
6:C:441:HOH:O	4:E:97:THR:HG21	1.98	0.62
1:A:61:ARG:HG2	3:G:94:ASN:HB3	1.84	0.60
1:A:175:THR:HG23	6:A:491:HOH:O	2.04	0.57
3:G:1:GLY:H2	3:G:27:SER:H	1.52	0.56
4:E:210:ARG:HH21	4:E:212:GLN:NE2	2.02	0.56
1:A:0:MET:HE2	1:A:169:LEU:HD13	1.90	0.54
4:H:155:HIS:HB3	4:H:216:TYR:HB2	1.89	0.54
4:E:128:VAL:HG23	4:E:238:ALA:HB3	1.90	0.53
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.43	0.53
1:A:151:LEU:HD22	3:G:51:LEU:HD12	1.89	0.53
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.89	0.53
1:C:4:SER:HB3	1:C:99:GLU:HG2	1.92	0.51
4:E:200:THR:HG23	6:E:335:HOH:O	2.11	0.50
4:H:30:ASN:O	4:H:51:SER:HA	2.11	0.49
3:G:159:LEU:HB3	4:H:172:CYS:HB2	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:G:121:LEU:HB3	4:H:129:PHE:HB3	1.96	0.48
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.49	0.48
3:G:1:GLY:H1	3:G:27:SER:H	1.60	0.48
1:C:212:MET:HG2	1:C:258:VAL:HG22	1.95	0.48
4:E:210:ARG:HE	4:E:212:GLN:HE21	1.62	0.48
1:A:186:VAL:HG11	1:A:269:VAL:HG22	1.95	0.47
3:D:4:ILE:HD12	3:D:98:ILE:O	2.14	0.47
3:G:28:GLY:HA3	3:G:93[A]:SER:HB3	1.97	0.46
1:A:215:MET:HG3	1:A:257:HIS:CD2	2.51	0.46
1:C:46:ARG:HA	1:C:46:ARG:HD3	1.68	0.45
1:C:259:GLU:HG3	1:C:264:HIS:ND1	2.33	0.44
2:F:24:ASN:HB3	2:F:65:LEU:HD11	2.00	0.44
1:C:233:SER:HB3	2:F:12:ARG:HG3	1.99	0.44
1:A:46:ARG:HD2	6:A:473:HOH:O	2.17	0.43
3:G:151:VAL:HA	3:G:175:SER:HB2	2.01	0.43
1:A:28:VAL:HG23	1:A:33:ILE:HD13	2.00	0.43
4:E:8:PRO:HD2	4:E:21:LEU:HD22	2.01	0.42
4:H:46:ILE:HG22	4:H:47:TYR:HD2	1.84	0.42
4:H:181:GLN:O	4:H:187:SER:HB2	2.20	0.42
3:G:89:ALA:HB1	3:G:97:LEU:HD22	2.02	0.42
3:G:78:LEU:HG	3:G:107:ILE:HD12	2.01	0.41
4:H:70:ASN:HD22	4:H:72:ARG:H	1.69	0.41
3:G:13:THR:O	3:G:16:ALA:HB3	2.20	0.41
4:H:148:ALA:HB2	4:H:213:VAL:HG21	2.03	0.41
3:G:18:VAL:HG12	3:G:78:LEU:HD11	2.03	0.40
4:E:2:ALA:HB3	4:E:26:ASP:OD2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/271 (96%)	252 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	240/271 (89%)	237 (99%)	3 (1%)	0	100	100
2	B	94/99 (95%)	94 (100%)	0	0	100	100
2	F	95/99 (96%)	93 (98%)	2 (2%)	0	100	100
3	D	198/203 (98%)	195 (98%)	3 (2%)	0	100	100
3	G	184/203 (91%)	176 (96%)	7 (4%)	1 (0%)	38	33
4	E	243/245 (99%)	238 (98%)	5 (2%)	0	100	100
4	H	227/245 (93%)	224 (99%)	3 (1%)	0	100	100
All	All	1541/1636 (94%)	1509 (98%)	31 (2%)	1 (0%)	59	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	2	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/241 (93%)	218 (97%)	7 (3%)	52	54
1	C	215/241 (89%)	206 (96%)	9 (4%)	40	38
2	B	83/94 (88%)	81 (98%)	2 (2%)	61	65
2	F	81/94 (86%)	79 (98%)	2 (2%)	60	63
3	D	171/180 (95%)	166 (97%)	5 (3%)	55	57
3	G	156/180 (87%)	149 (96%)	7 (4%)	38	35
4	E	206/211 (98%)	200 (97%)	6 (3%)	55	57
4	H	184/211 (87%)	173 (94%)	11 (6%)	27	22
All	All	1321/1452 (91%)	1272 (96%)	49 (4%)	45	45

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	LEU
1	A	152	TYR
1	A	179	THR
1	A	221	ILE
1	A	222	VAL
1	A	259	GLU
2	B	64	LEU
2	B	70	PHE
1	C	16	ILE
1	C	57	ASP
1	C	73	PHE
1	C	82	ARG
1	C	100	LEU
1	C	152	TYR
1	C	169	LEU
1	C	179	THR
1	C	259	GLU
3	D	78	LEU
3	D	93	SER
3	D	122	ARG
3	D	125	LYS
3	D	179	ASP
4	E	9	LYS
4	E	97	THR
4	E	132	SER
4	E	185	ASN
4	E	194	ARG
4	E	206	ARG
2	F	70	PHE
2	F	88	SER
3	G	27	SER
3	G	75	LEU
3	G	78	LEU
3	G	85	SER
3	G	131	VAL
3	G	157	CYS
3	G	159	LEU
4	H	18	SER
4	H	19	MET
4	H	21	LEU
4	H	70	ASN
4	H	78	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	H	111	ARG
4	H	112	LEU
4	H	155	HIS
4	H	184	LEU
4	H	194	ARG
4	H	226	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	HIS
1	A	153	GLN
1	A	187	ASN
1	A	203	HIS
2	B	31	HIS
4	E	181	GLN
4	E	212	GLN
2	F	8	GLN
4	H	70	ASN
4	H	155	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	2L4	A	301	1	20,21,22	1.03	2 (10%)	23,28,29	1.36	2 (8%)
5	2L4	C	301	1	20,21,22	1.37	3 (15%)	23,28,29	1.29	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2L4	A	301	1	-	0/18/18/19	0/1/1/1
5	2L4	C	301	1	-	0/18/18/19	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	2L4	C4A-C8A	4.06	1.48	1.42
5	C	301	2L4	O2-C2	3.19	1.26	1.21
5	A	301	2L4	C4A-C8A	2.84	1.46	1.42
5	A	301	2L4	O2-C2	2.18	1.24	1.21
5	C	301	2L4	C8A-N8	2.04	1.35	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	301	2L4	C4A-C8A-N1	3.41	119.38	114.86
5	A	301	2L4	C4A-C8A-N8	-3.26	117.90	124.54
5	A	301	2L4	C4A-C8A-N1	3.21	119.11	114.86
5	C	301	2L4	C4A-N5-C6	2.65	125.11	118.60
5	C	301	2L4	C4A-C8A-N8	-2.32	119.83	124.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/271 (97%)	-0.07	2 (0%) 83 87	20, 31, 55, 75	0
1	C	249/271 (91%)	-0.08	3 (1%) 75 80	18, 33, 58, 73	0
2	B	96/99 (96%)	-0.16	0 100 100	19, 37, 60, 64	0
2	F	97/99 (97%)	0.16	1 (1%) 79 83	25, 51, 68, 76	0
3	D	200/203 (98%)	-0.14	0 100 100	16, 27, 52, 67	0
3	G	189/203 (93%)	0.84	39 (20%) 1 1	22, 47, 88, 96	0
4	E	243/245 (99%)	-0.19	1 (0%) 90 92	16, 26, 47, 80	1 (0%)
4	H	231/245 (94%)	0.28	11 (4%) 29 32	26, 47, 78, 89	0
All	All	1569/1636 (95%)	0.07	57 (3%) 41 45	16, 34, 71, 96	1 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	146	SER	5.2
3	G	197	PHE	5.1
3	G	180	PHE	5.1
3	G	150	ASP	4.9
4	H	136	ILE	4.8
3	G	149	SER	4.5
3	G	166	PHE	4.4
3	G	191	ILE	4.2
4	H	142	ALA	4.0
3	G	190	ILE	3.8
4	H	137	SER	3.7
3	G	196	PHE	3.5
4	E	184	LEU	3.5
4	H	197	VAL	3.4
3	G	198	PRO	3.4
3	G	119	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	G	189	SER	3.3
3	G	153	ILE	3.2
3	G	111	ILE	3.1
3	G	132	CYS	3.1
3	G	164	MET	3.1
3	G	163	SER	3.1
3	G	110	ASP	3.0
3	G	192	PRO	3.0
3	G	175	SER	3.0
3	G	186	PHE	2.9
4	H	162	VAL	2.9
4	H	144	LEU	2.9
3	G	187	ASN	2.8
3	G	195	THR	2.8
4	H	202	TRP	2.8
1	A	17	HIS	2.8
3	G	139	SER	2.8
3	G	183	ALA	2.8
3	G	151	VAL	2.7
3	G	182	CYS	2.5
1	C	199	PHE	2.5
3	G	141	THR	2.5
3	G	138	ASP	2.5
1	A	221	ILE	2.4
3	G	179	ASP	2.4
4	H	143	THR	2.4
3	G	188	ASN	2.4
4	H	200	THR	2.4
3	G	147	LYS	2.4
2	F	78	TYR	2.3
3	G	117	ALA	2.3
3	G	143	VAL	2.3
3	G	159	LEU	2.2
3	G	144	SER	2.2
3	G	161	MET	2.1
1	C	265	MET	2.1
4	H	199	ALA	2.1
1	C	225	ILE	2.1
3	G	185	ALA	2.0
4	H	133	GLU	2.0
3	G	140	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	2L4	A	301	21/22	0.13	1.71	19,23,29,30	0
5	2L4	C	301	21/22	0.09	-0.99	17,22,28,30	0

6.5 Other polymers

There are no such residues in this entry.