



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 08:44 AM GMT

PDB ID : 3NSP
Title : Crystal structure of tetrameric RXRalpha-LBD
Authors : Zhang, H.; Hu, T.; Li, L.; Zhou, R.; Chen, L.; Hu, L.; Jiang, H.; Shen, X.
Deposited on : 2010-07-02
Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	240	
1	B	240	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3074 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoid X receptor, alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	0	0	0
			1526	983	264	270	9			
1	B	190	Total	C	N	O	S	0	0	0
			1514	976	263	266	9			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	19	Total	O	0	0
			19	19		
2	B	15	Total	O	0	0
			15	15		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retinoid X receptor, alpha

Chain A: 



- Molecule 1: Retinoid X receptor, alpha

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.83Å 100.27Å 47.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.90 15.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	91.1 (15.00-2.90) 91.1 (15.00-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.80 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.222 , 0.308 0.217 , 0.301	Depositor DCC
R_{free} test set	556 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 14.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 11445 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3074	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/1558	0.81	5/2106 (0.2%)
1	B	0.69	0/1546	0.77	1/2086 (0.0%)
All	All	0.74	0/3104	0.79	6/4192 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	ALA	CB-CA-C	-6.93	99.70	110.10
1	B	331	HIS	N-CA-C	6.56	128.71	111.00
1	A	276	LEU	CB-CA-C	-6.19	98.45	110.20
1	A	277	PHE	N-CA-C	-5.82	95.28	111.00
1	A	326	LEU	CA-CB-CG	5.81	128.66	115.30
1	A	264	PRO	N-CA-C	5.12	125.41	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1526	0	1567	62	0
1	B	1514	0	1551	43	0
2	A	19	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	15	0	0	1	0
All	All	3074	0	3118	105	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 17.

All (105) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:334:ARG:C	1:A:335:ASN:OD1	2.00	1.00
1:A:326:LEU:HD21	1:A:332:VAL:CG2	1.95	0.96
1:A:326:LEU:HD21	1:A:332:VAL:HG21	1.49	0.94
1:B:442:ILE:O	1:B:446:PRO:HD3	1.70	0.92
1:B:442:ILE:O	1:B:446:PRO:CD	2.21	0.89
1:A:316:ARG:HG2	1:A:325:LEU:HD23	1.56	0.85
1:A:369:CYS:CB	1:A:400:LEU:HD13	2.10	0.81
1:B:287:PRO:O	1:B:288:HIS:HB2	1.80	0.81
1:A:275:GLN:O	1:A:276:LEU:HB2	1.81	0.80
1:B:456:GLU:H	1:B:456:GLU:CD	1.88	0.77
1:A:275:GLN:O	1:A:276:LEU:CB	2.31	0.77
1:B:307:GLU:HG2	1:B:425:LEU:HG	1.66	0.77
1:B:446:PRO:O	1:B:450:PHE:HB2	1.86	0.75
1:A:369:CYS:HB2	1:A:400:LEU:HD13	1.67	0.75
1:A:327:ALA:O	2:A:30:HOH:O	2.05	0.74
1:B:288:HIS:HD2	1:B:291:GLU:OE1	1.74	0.70
1:A:390:GLU:O	1:A:394:GLU:HG3	1.91	0.69
1:B:411:GLN:HE22	1:B:414:ARG:HH11	1.41	0.68
1:B:444:ASP:N	1:B:444:ASP:OD1	2.25	0.67
1:B:322:ASP:OD1	1:B:333:HIS:ND1	2.28	0.67
1:A:326:LEU:O	1:A:327:ALA:HB3	1.95	0.67
1:B:460:GLN:O	1:B:461:MET:SD	2.52	0.67
1:A:390:GLU:OE2	2:A:28:HOH:O	2.15	0.64
1:A:335:ASN:OD1	1:A:335:ASN:N	2.29	0.64
1:A:445:THR:HB	1:A:446:PRO:HD3	1.80	0.63
1:A:326:LEU:HD22	1:A:441:LEU:HG	1.80	0.63
1:B:377:ASN:HA	1:B:393:ARG:NH2	2.15	0.62
1:B:443:GLY:C	1:B:444:ASP:OD1	2.38	0.62
1:A:341:GLY:HA3	1:A:435:HIS:CE1	2.36	0.61
1:B:320:VAL:HG21	1:B:331:HIS:HE1	1.65	0.60
1:B:442:ILE:O	1:B:446:PRO:HD2	2.02	0.60
1:B:451:LEU:O	1:B:455:LEU:HD13	2.02	0.59
1:A:313:PHE:HA	1:A:316:ARG:NH1	2.18	0.59
1:A:326:LEU:CD2	1:A:332:VAL:CG2	2.79	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:370:LEU:HD21	1:B:418:LEU:HB3	1.85	0.58
1:B:273:ASP:HA	1:B:276:LEU:HB2	1.85	0.58
1:B:394:GLU:HA	1:B:397:TYR:CE2	2.40	0.57
1:A:454:MET:C	1:A:455:LEU:HG	2.25	0.57
1:B:320:VAL:HG22	1:B:321:LYS:H	1.70	0.56
1:B:331:HIS:C	1:B:331:HIS:CD2	2.80	0.56
1:A:295:ASP:O	1:A:299:ILE:HG13	2.07	0.55
1:A:333:HIS:O	1:A:335:ASN:OD1	2.25	0.54
1:B:394:GLU:HG2	1:B:397:TYR:OH	2.07	0.54
1:A:411:GLN:HE22	1:A:414:ARG:HH11	1.55	0.53
1:A:439:PHE:C	1:A:439:PHE:CD1	2.81	0.53
1:B:377:ASN:HA	1:B:393:ARG:HH21	1.75	0.52
1:A:264:PRO:HG2	1:A:265:VAL:N	2.25	0.52
1:B:421:ARG:HH11	1:B:421:ARG:HG2	1.74	0.51
1:B:407:LYS:NZ	2:B:27:HOH:O	2.39	0.51
1:A:326:LEU:O	1:A:328:THR:N	2.43	0.51
1:A:334:ARG:O	1:A:335:ASN:OD1	2.27	0.51
1:A:326:LEU:O	1:A:328:THR:OG1	2.30	0.50
1:A:332:VAL:HG12	1:A:333:HIS:H	1.76	0.50
1:B:362:MET:HE3	1:B:367:LEU:HB2	1.94	0.49
1:B:373:ILE:HD11	1:B:397:TYR:HB3	1.95	0.49
1:B:443:GLY:O	1:B:447:ILE:HD13	2.12	0.49
1:A:385:ASN:HD21	1:A:388:GLU:HG2	1.79	0.48
1:A:306:ASN:O	1:A:310:ILE:HG13	2.14	0.48
1:A:264:PRO:HG2	1:A:265:VAL:H	1.79	0.48
1:A:272:ALA:O	1:A:275:GLN:O	2.32	0.47
1:B:407:LYS:HD2	1:B:408:TYR:CE1	2.49	0.47
1:B:313:PHE:C	1:B:313:PHE:CD1	2.88	0.47
1:A:333:HIS:C	1:A:335:ASN:OD1	2.53	0.47
1:B:411:GLN:HE22	1:B:414:ARG:NH1	2.10	0.47
1:B:318:ILE:HD11	1:B:357:MET:HB3	1.98	0.46
1:A:345:ILE:O	1:A:349:VAL:HG23	2.16	0.46
1:B:449:THR:O	1:B:453:GLU:HG3	2.15	0.46
1:B:288:HIS:CD2	1:B:291:GLU:OE1	2.63	0.46
1:A:326:LEU:HD21	1:A:332:VAL:HG23	1.90	0.45
1:B:440:LYS:HE3	1:B:440:LYS:HB3	1.76	0.45
1:A:310:ILE:HA	1:A:313:PHE:CE2	2.51	0.45
1:A:326:LEU:CD2	1:A:332:VAL:HG23	2.47	0.45
1:A:445:THR:HB	1:A:446:PRO:CD	2.44	0.45
1:A:288:HIS:HD2	1:A:291:GLU:OE1	2.00	0.45
1:A:423:PRO:O	1:A:426:ARG:HB3	2.15	0.45
1:A:310:ILE:HA	1:A:313:PHE:CZ	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:322:ASP:OD1	1:B:333:HIS:CE1	2.71	0.44
1:A:314:SER:O	1:A:354:VAL:HG22	2.16	0.44
1:A:373:ILE:O	1:A:393:ARG:NH2	2.51	0.44
1:B:411:GLN:NE2	1:B:414:ARG:HH11	2.12	0.44
1:A:421:ARG:HD2	1:A:421:ARG:N	2.33	0.44
1:A:326:LEU:O	1:A:327:ALA:CB	2.61	0.43
1:A:385:ASN:ND2	1:A:388:GLU:HG2	2.34	0.43
1:A:307:GLU:HG2	1:A:425:LEU:HG	2.00	0.43
1:A:373:ILE:HD11	1:A:397:TYR:HB3	2.00	0.43
1:A:451:LEU:O	1:A:454:MET:SD	2.77	0.42
1:A:287:PRO:O	1:A:288:HIS:HB2	2.20	0.42
1:A:310:ILE:CD1	1:A:429:GLY:HA2	2.49	0.42
1:B:320:VAL:HG21	1:B:331:HIS:CE1	2.51	0.42
1:A:326:LEU:C	1:A:328:THR:H	2.22	0.42
1:A:310:ILE:HG12	1:A:313:PHE:CZ	2.55	0.42
1:B:274:LYS:HG3	1:B:275:GLN:N	2.35	0.42
1:A:326:LEU:C	1:A:328:THR:N	2.72	0.41
1:A:451:LEU:HA	1:A:451:LEU:HD12	1.68	0.41
1:A:305:TRP:CZ2	1:A:306:ASN:ND2	2.88	0.41
1:A:326:LEU:HG	1:A:330:LEU:O	2.20	0.41
1:A:310:ILE:O	1:A:314:SER:HB2	2.21	0.41
1:B:306:ASN:ND2	1:B:429:GLY:O	2.51	0.41
1:A:321:LYS:HG3	1:A:321:LYS:O	2.21	0.41
1:A:407:LYS:HG2	1:A:408:TYR:CE1	2.56	0.41
1:A:288:HIS:CD2	1:A:291:GLU:OE1	2.75	0.40
1:B:446:PRO:O	1:B:450:PHE:N	2.52	0.40
1:A:400:LEU:HD23	1:A:415:PHE:HE1	1.86	0.40
1:B:331:HIS:CD2	1:B:331:HIS:O	2.75	0.40
1:B:422:LEU:N	1:B:423:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/240 (80%)	180 (94%)	11 (6%)	0	100	100
1	B	186/240 (78%)	167 (90%)	19 (10%)	0	100	100
All	All	377/480 (78%)	347 (92%)	30 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/207 (80%)	145 (87%)	21 (13%)	6	18
1	B	164/207 (79%)	137 (84%)	27 (16%)	3	9
All	All	330/414 (80%)	282 (86%)	48 (14%)	5	13

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	264	PRO
1	A	266	THR
1	A	276	LEU
1	A	321	LYS
1	A	322	ASP
1	A	326	LEU
1	A	328	THR
1	A	332	VAL
1	A	335	ASN
1	A	346	PHE
1	A	383	LEU
1	A	384	SER
1	A	421	ARG
1	A	427	SER
1	A	436	LEU
1	A	439	PHE
1	A	444	ASP
1	A	451	LEU
1	A	452	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	454	MET
1	A	455	LEU
1	B	273	ASP
1	B	274	LYS
1	B	276	LEU
1	B	290	SER
1	B	307	GLU
1	B	312	SER
1	B	313	PHE
1	B	314	SER
1	B	322	ASP
1	B	325	LEU
1	B	326	LEU
1	B	336	SER
1	B	379	ASP
1	B	381	LYS
1	B	383	LEU
1	B	394	GLU
1	B	395	LYS
1	B	399	SER
1	B	407	LYS
1	B	431	LYS
1	B	436	LEU
1	B	440	LYS
1	B	444	ASP
1	B	450	PHE
1	B	451	LEU
1	B	456	GLU
1	B	462	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	288	HIS
1	A	338	HIS
1	A	385	ASN
1	A	411	GLN
1	A	435	HIS
1	B	288	HIS
1	B	297	GLN
1	B	331	HIS
1	B	411	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	193/240 (80%)	-0.50	3 (1%) 68 78	6, 20, 43, 51	0
1	B	190/240 (79%)	-0.53	1 (0%) 88 93	4, 19, 40, 52	0
All	All	383/480 (79%)	-0.51	4 (1%) 79 86	4, 20, 43, 52	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	455	LEU	3.4
1	B	462	THR	2.6
1	A	327	ALA	2.0
1	A	337	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.