



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 06:49 PM GMT

PDB ID : 1NTK
Title : Crystal Structure of Mitochondrial Cytochrome bc1 in Complex with Antimycin A1
Authors : Gao, X.; Wen, X.; Esser, L.; Quinn, B.; Yu, L.; Yu, C.-A.; Xia, D.
Deposited on : 2003-01-30
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

| | | |
|--------------------------------|---|--------------------------|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.15 2013 |
| Xtriage (Phenix) | : | NOT EXECUTED |
| EDS | : | NOT EXECUTED |
| Percentile statistics | : | 21963 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et. al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | stable22683 |

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.












Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| Clashscore | 79885 | 2154 (2.60-2.60) |
| Ramachandran outliers | 78287 | 2113 (2.60-2.60) |
| Sidechain outliers | 78261 | 2113 (2.60-2.60) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | A | 446 |  |
| 2 | B | 439 |  |
| 3 | C | 379 |  |
| 4 | D | 241 |  |
| 5 | E | 196 |  |
| 6 | F | 110 |  |
| 7 | G | 81 |  |
| 8 | H | 78 |  |
| 9 | I | 57 |  |
| 10 | J | 62 |  |
| 11 | K | 56 |  |

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 17118 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquinol-cytochrome C reductase complex core protein I, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| 1 | A | 446 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 3457 | 2161 | 609 | 667 | 20 | | | |

- Molecule 2 is a protein called Ubiquinol-cytochrome C reductase complex core protein 2, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 2 | B | 423 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 3171 | 1993 | 562 | 609 | 7 | | | |

- Molecule 3 is a protein called Cytochrome b.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| 3 | C | 378 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 3003 | 2013 | 471 | 501 | 18 | | | |

- Molecule 4 is a protein called cytochrome c1.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| 4 | D | 241 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1918 | 1225 | 330 | 348 | 15 | | | |

- Molecule 5 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE IRON-SULFUR SUBUNIT, MITOCHONDRIAL.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 5 | E | 196 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1519 | 957 | 263 | 291 | 8 | | | |

- Molecule 6 is a protein called Ubiquinol-cytochrome C reductase complex 14 kDa protein.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 6 | F | 105 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 910 | 576 | 165 | 167 | 2 | | | |

- Molecule 7 is a protein called Ubiquinol-cytochrome C reductase complex ubiquinone-binding protein QP-C.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 7 | G | 79 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 659 | 430 | 122 | 106 | 1 | | | |

- Molecule 8 is a protein called Ubiquinol-cytochrome C reductase complex 11 kDa protein.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 8 | H | 76 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 626 | 378 | 109 | 134 | 5 | | | |

- Molecule 9 is a protein called Ubiquinol-cytochrome C reductase 8 kDa protein.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|---------|-------|
| 9 | I | 57 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 406 | 253 | 77 | 74 | 2 | | | |

- Molecule 10 is a protein called Ubiquinol-cytochrome C reductase complex 7.2 kDa protein.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|---------|-------|
| 10 | J | 60 | Total | C | N | O | 0 | 0 | 0 |
| | | | 495 | 324 | 86 | 85 | | | |

- Molecule 11 is a protein called Ubiquinol-cytochrome C reductase complex 6.4 kDa protein.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|---------|-------|
| 11 | K | 53 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 441 | 295 | 79 | 66 | 1 | | | |

There is a discrepancy between the modelled and reference sequences:


| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| K | 22 | GLN | SER | SEE REMARK 999 | UNP P07552 |

- Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------------|---------|---------|--------|--------|---------|---------|
| 12 | C | 1 | Total 43 | C 34 | Fe 1 | N 4 | O 4 | 0 | 0 |
| 12 | C | 1 | Total 43 | C 34 | Fe 1 | N 4 | O 4 | 0 | 0 |
| 12 | D | 1 | Total 43 | C 34 | Fe 1 | N 4 | O 4 | 0 | 0 |

- Molecule 13 is ANTIMYCIN A1 (three-letter code: AY1) (formula: $C_{27}H_{44}N_2O_9$).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---|---------|---------|
| 13 | C | 1 | Total | C | N | O | 0 | 0 |
| | | | 38 | 27 | 2 | 9 | | |

- Molecule 14 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---------|---------|
| 14 | E | 1 | Total | Fe | S | 0 | 0 |
| | | | 4 | 2 | 2 | | |

- Molecule 15 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 15 | A | 95 | Total | O | 0 | 0 |
| | | | 95 | 95 | | |
| 15 | B | 83 | Total | O | 0 | 0 |
| | | | 83 | 83 | | |
| 15 | C | 61 | Total | O | 0 | 0 |
| | | | 61 | 61 | | |
| 15 | D | 27 | Total | O | 0 | 0 |
| | | | 27 | 27 | | |
| 15 | E | 3 | Total | O | 0 | 0 |
| | | | 3 | 3 | | |
| 15 | F | 30 | Total | O | 0 | 0 |
| | | | 30 | 30 | | |
| 15 | G | 22 | Total | O | 0 | 0 |
| | | | 22 | 22 | | |
| 15 | I | 6 | Total | O | 0 | 0 |
| | | | 6 | 6 | | |
| 15 | J | 5 | Total | O | 0 | 0 |
| | | | 5 | 5 | | |
| 15 | K | 10 | Total | O | 0 | 0 |
| | | | 10 | 10 | | |


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Ubiquinol-cytochrome C reductase complex core protein I, mitochondrial

Chain A:




- Molecule 2: Ubiquinol-cytochrome C reductase complex core protein 2, mitochondrial

Chain B:




- Molecule 3: Cytochrome b

Chain C:



- Molecule 4: cytochrome c1

Chain D:




- Molecule 5: UBIQUINOL-CYTOCHROME C REDUCTASE IRON-SULFUR SUBUNIT, MITOCHONDRIAL

Chain E:




- Molecule 6: Ubiquinol-cytochrome C reductase complex 14 kDa protein

Chain F:




- Molecule 7: Ubiquinol-cytochrome C reductase complex ubiquinone-binding protein QP-C

Chain G:




- Molecule 8: Ubiquinol-cytochrome C reductase complex 11 kDa protein

Chain H:




- Molecule 9: Ubiquinol-cytochrome C reductase 8 kDa protein

Chain I:




- Molecule 10: Ubiquinol-cytochrome C reductase complex 7.2 kDa protein

Chain J:



- Molecule 11: Ubiquinol-cytochrome C reductase complex 6.4 kDa protein

Chain K:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | I 41 2 2 | Depositor |
| Cell constants a, b, c, α , β , γ | 153.78Å 153.78Å 592.50Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 40.00 – 2.60 | Depositor |
| % Data completeness (in resolution range) | 98.5 (40.00-2.60) | Depositor |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | REFMAC 5 | Depositor |
| R, R_{free} | 0.233 , 0.270 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 17118 | wwPDB-VP |
| Average B, all atoms (Å ²) | 33.0 | wwPDB-VP |

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, AY1, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | A | 0.84 | 2/3530 (0.1%) | 0.85 | 11/4792 (0.2%) |
| 2 | B | 1.11 | 1/3231 (0.0%) | 0.94 | 11/4386 (0.3%) |
| 3 | C | 0.78 | 3/3100 (0.1%) | 0.82 | 9/4242 (0.2%) |
| 4 | D | 0.60 | 0/1977 | 0.87 | 11/2684 (0.4%) |
| 5 | E | 0.53 | 0/1553 | 0.78 | 8/2100 (0.4%) |
| 6 | F | 0.96 | 0/929 | 0.91 | 5/1246 (0.4%) |
| 7 | G | 0.77 | 0/681 | 0.79 | 0/922 |
| 8 | H | 0.56 | 0/632 | 0.83 | 3/847 (0.4%) |
| 9 | I | 1.07 | 1/411 (0.2%) | 1.18 | 1/558 (0.2%) |
| 10 | J | 0.56 | 0/508 | 0.76 | 2/686 (0.3%) |
| 11 | K | 0.66 | 0/457 | 0.77 | 2/625 (0.3%) |
| All | All | 0.83 | 7/17009 (0.0%) | 0.86 | 63/23088 (0.3%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 2 | B | 1 | 0 |
| 3 | C | 2 | 2 |
| 7 | G | 2 | 0 |
| 9 | I | 1 | 0 |
| All | All | 6 | 2 |

The worst 5 of 7 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 9 | I | 25 | ALA | CA-CB | -6.95 | 1.37 | 1.52 |
| 3 | C | 26 | ASN | CB-CG | -6.36 | 1.36 | 1.51 |
| 3 | C | 194 | MET | SD-CE | 5.82 | 2.10 | 1.77 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|-------|-------------|----------|
| 2 | B | 159 | VAL | CB-CG2 | -5.71 | 1.40 | 1.52 |
| 3 | C | 379 | TRP | CB-CG | -5.44 | 1.40 | 1.50 |

The worst 5 of 63 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|------|-------------|----------|
| 6 | F | 42 | ASP | CB-CG-OD2 | 7.55 | 125.10 | 118.30 |
| 3 | C | 345 | HIS | N-CA-C | 6.71 | 129.12 | 111.00 |
| 1 | A | 333 | ASP | CB-CG-OD2 | 6.66 | 124.30 | 118.30 |
| 3 | C | 252 | ASP | CB-CG-OD2 | 6.55 | 124.19 | 118.30 |
| 5 | E | 67 | ASP | CB-CG-OD2 | 6.40 | 124.06 | 118.30 |

5 of 6 chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 2 | B | 304 | HIS | CA |
| 3 | C | 58 | ASP | CA |
| 3 | C | 221 | HIS | CA |
| 7 | G | 11 | ARG | CA |
| 7 | G | 73 | ASN | CA |

All (2) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-------------------|
| 3 | C | 221 | HIS | Mainchain,Peptide |

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 3457 | 0 | 3356 | 229 | 0 |
| 2 | B | 3171 | 0 | 3152 | 199 | 0 |
| 3 | C | 3003 | 0 | 3065 | 153 | 0 |
| 4 | D | 1918 | 0 | 1870 | 113 | 0 |
| 5 | E | 1519 | 0 | 1504 | 34 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 6 | F | 910 | 0 | 904 | 43 | 0 |
| 7 | G | 659 | 0 | 658 | 27 | 0 |
| 8 | H | 626 | 0 | 595 | 25 | 0 |
| 9 | I | 406 | 0 | 435 | 260 | 0 |
| 10 | J | 495 | 0 | 493 | 27 | 0 |
| 11 | K | 441 | 0 | 450 | 18 | 0 |
| 12 | C | 86 | 0 | 60 | 16 | 0 |
| 12 | D | 43 | 0 | 30 | 6 | 0 |
| 13 | C | 38 | 0 | 37 | 12 | 0 |
| 14 | E | 4 | 0 | 0 | 1 | 0 |
| 15 | A | 95 | 0 | 0 | 10 | 0 |
| 15 | B | 83 | 0 | 0 | 21 | 0 |
| 15 | C | 61 | 0 | 0 | 7 | 0 |
| 15 | D | 27 | 0 | 0 | 2 | 0 |
| 15 | E | 3 | 0 | 0 | 2 | 0 |
| 15 | F | 30 | 0 | 0 | 4 | 0 |
| 15 | G | 22 | 0 | 0 | 3 | 0 |
| 15 | I | 6 | 0 | 0 | 3 | 0 |
| 15 | J | 5 | 0 | 0 | 0 | 0 |
| 15 | K | 10 | 0 | 0 | 2 | 0 |
| All | All | 17118 | 0 | 16609 | 883 | 0 |

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 883 close contacts within the same asymmetric unit are listed below.

| Atom-1 | Atom-2 | Distance(Å) | Clash(Å) |
|-----------------|----------------|-------------|----------|
| 1:A:284:TYR:HE1 | 9:I:20:ARG:CG | 1.09 | 1.65 |
| 1:A:284:TYR:CE1 | 9:I:20:ARG:HG3 | 1.24 | 1.61 |
| 1:A:279:HIS:CE1 | 9:I:52:ARG:HB2 | 1.56 | 1.40 |
| 2:B:283:PRO:HG3 | 9:I:31:GLN:CG | 1.52 | 1.39 |
| 3:C:194:MET:SD | 3:C:194:MET:CE | 2.10 | 1.38 |

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|----------|-------------|-----|
| 1 | A | 444/446 (100%) | 396 (89%) | 42 (10%) | 6 (1%) | 16 | 32 |
| 2 | B | 421/439 (96%) | 389 (92%) | 24 (6%) | 8 (2%) | 12 | 23 |
| 3 | C | 376/379 (99%) | 317 (84%) | 37 (10%) | 22 (6%) | 2 | 2 |
| 4 | D | 239/241 (99%) | 172 (72%) | 47 (20%) | 20 (8%) | 1 | 1 |
| 5 | E | 194/196 (99%) | 181 (93%) | 9 (5%) | 4 (2%) | 11 | 19 |
| 6 | F | 103/110 (94%) | 95 (92%) | 8 (8%) | 0 | 100 | 100 |
| 7 | G | 77/81 (95%) | 65 (84%) | 8 (10%) | 4 (5%) | 3 | 3 |
| 8 | H | 74/78 (95%) | 58 (78%) | 11 (15%) | 5 (7%) | 2 | 2 |
| 9 | I | 55/57 (96%) | 25 (46%) | 17 (31%) | 13 (24%) | 0 | 0 |
| 10 | J | 58/62 (94%) | 46 (79%) | 8 (14%) | 4 (7%) | 2 | 2 |
| 11 | K | 51/56 (91%) | 41 (80%) | 7 (14%) | 3 (6%) | 2 | 2 |
| All | All | 2092/2145 (98%) | 1785 (85%) | 218 (10%) | 89 (4%) | 4 | 5 |

5 of 89 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 74 | ALA |
| 1 | A | 225 | GLU |
| 2 | B | 234 | GLY |
| 2 | B | 304 | HIS |
| 2 | B | 305 | GLN |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------------|-----------|----------|-------------|----|
| 1 | A | 370/370 (100%) | 324 (88%) | 46 (12%) | 7 | 13 |
| 2 | B | 332/343 (97%) | 291 (88%) | 41 (12%) | 7 | 13 |
| 3 | C | 326/327 (100%) | 286 (88%) | 40 (12%) | 7 | 13 |
| 4 | D | 206/206 (100%) | 172 (84%) | 34 (16%) | 3 | 5 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 5 | E | 168/168 (100%) | 158 (94%) | 10 (6%) | 27 | 51 |
| 6 | F | 96/98 (98%) | 81 (84%) | 15 (16%) | 4 | 6 |
| 7 | G | 68/71 (96%) | 53 (78%) | 15 (22%) | 1 | 2 |
| 8 | H | 73/74 (99%) | 62 (85%) | 11 (15%) | 4 | 7 |
| 9 | I | 44/44 (100%) | 33 (75%) | 11 (25%) | 1 | 1 |
| 10 | J | 50/52 (96%) | 45 (90%) | 5 (10%) | 11 | 20 |
| 11 | K | 43/46 (94%) | 31 (72%) | 12 (28%) | 0 | 1 |
| All | All | 1776/1799 (99%) | 1536 (86%) | 240 (14%) | 6 | 10 |

5 of 240 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | C | 263 | ASN |
| 4 | D | 58 | GLU |
| 9 | I | 47 | ARG |
| 3 | C | 268 | ILE |
| 3 | C | 349 | THR |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 162 | ASN |
| 2 | B | 329 | GLN |
| 8 | H | 49 | GLN |
| 2 | B | 174 | ASN |
| 2 | B | 248 | ASN |

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|-------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | # $ Z > 2$ | Counts | RMSZ | # $ Z > 2$ |
| 12 | HEM | C | 381 | 3 | 49,50,50 | 11.88 | 28 (57%) | 46,82,82 | 3.59 | 23 (50%) |
| 12 | HEM | C | 382 | 3 | 49,50,50 | 11.38 | 29 (59%) | 46,82,82 | 3.34 | 22 (47%) |
| 13 | AY1 | C | 383 | - | 39,39,39 | 4.53 | 13 (33%) | 50,53,53 | 3.44 | 23 (46%) |
| 12 | HEM | D | 242 | 4 | 49,50,50 | 12.83 | 29 (59%) | 46,82,82 | 3.51 | 20 (43%) |
| 14 | FES | E | 197 | 5 | 0,4,4 | 0.00 | - | 0,4,4 | 0.00 | - |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|-----------|--------------|---------|
| 12 | HEM | C | 381 | 3 | - | 0/14/114/114 | 0/0/8/8 |
| 12 | HEM | C | 382 | 3 | - | 0/14/114/114 | 0/0/8/8 |
| 13 | AY1 | C | 383 | - | 3/3/10/13 | 0/40/53/53 | 0/1/2/2 |
| 12 | HEM | D | 242 | 4 | - | 0/14/114/114 | 0/0/8/8 |
| 14 | FES | E | 197 | 5 | - | 0/0/4/4 | 0/0/1/1 |

The worst 5 of 99 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|--------|-------------|----------|
| 12 | C | 381 | HEM | C3D-C4D | -51.53 | 1.31 | 1.44 |
| 12 | D | 242 | HEM | C2D-C1D | -50.18 | 1.32 | 1.44 |
| 12 | D | 242 | HEM | C3D-C4D | -49.09 | 1.32 | 1.44 |
| 12 | D | 242 | HEM | C2B-C1B | -49.01 | 1.32 | 1.44 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|--------|-------------|----------|
| 12 | C | 381 | HEM | C2D-C1D | -43.87 | 1.33 | 1.44 |

The worst 5 of 88 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 13 | C | 383 | AY1 | O5-C12-C13 | 10.01 | 128.52 | 109.33 |
| 12 | C | 382 | HEM | C4A-CHB-C1B | -9.96 | 114.36 | 127.47 |
| 12 | C | 381 | HEM | C4A-CHB-C1B | -9.82 | 114.54 | 127.47 |
| 12 | D | 242 | HEM | C1A-CHA-C4D | -9.74 | 114.65 | 127.47 |
| 12 | C | 381 | HEM | C1A-CHA-C4D | -9.66 | 114.76 | 127.47 |

All (3) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 13 | C | 383 | AY1 | C8 |
| 13 | C | 383 | AY1 | C17 |
| 13 | C | 383 | AY1 | C12 |

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.