



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 02:02 PM GMT

PDB ID : 1NYT
Title : SHIKIMATE DEHYDROGENASE AroE COMPLEXED WITH NADP+
Authors : Roszak, A.W.; Lapthorn, A.J.
Deposited on : 2003-02-13
Resolution : 1.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1768 (1.50-1.50)
Ramachandran outliers	78287	1720 (1.50-1.50)
Sidechain outliers	78261	1718 (1.50-1.50)
RSRZ outliers	66119	1514 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	271	
1	B	271	
1	C	271	
1	D	271	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	A	1402	-	X
2	SO4	A	1404	-	X
2	SO4	B	1414	-	X
2	SO4	B	1434	-	X
2	SO4	C	1423	-	X
2	SO4	C	1426	-	X
2	SO4	D	1433	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
4	DTV	A	1405	-	X

2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 9929 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Shikimate 5-dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	6	0
			2088	1331	358	391	8			
1	B	271	Total	C	N	O	S	0	8	0
			2090	1329	359	393	9			
1	C	269	Total	C	N	O	S	0	6	0
			2066	1314	355	388	9			
1	D	270	Total	C	N	O	S	0	10	0
			2082	1323	355	395	9			

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is (2S,3S)-1,4-DIMERCAPTOBUTANE-2,3-DIOL (three-letter code: DTV) (formula: C₄H₁₀O₂S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	365	Total	O	0	0
			365	365		
5	B	391	Total	O	0	0
			391	391		
5	C	302	Total	O	0	0
			302	302		
5	D	275	Total	O	0	0
			275	275		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Shikimate 5-dehydrogenase

Chain A: 




- Molecule 1: Shikimate 5-dehydrogenase

Chain B: 




- Molecule 1: Shikimate 5-dehydrogenase

Chain C: 



- Molecule 1: Shikimate 5-dehydrogenase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.52Å 140.02Å 102.71Å 90.00° 122.07° 90.00°	Depositor
Resolution (Å)	38.92 – 1.50 38.92 – 1.50	Depositor EDS
% Data completeness (in resolution range)	93.1 (38.92-1.50) 93.1 (38.92-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.1.27	Depositor
R, R_{free}	0.132 , 0.169 0.150 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.5	EDS
Estimated twinning fraction	0.087 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 196162 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9929	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, SO4, DTV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/2162	0.85	1/2934 (0.0%)
1	B	0.83	1/2174 (0.0%)	0.84	0/2949
1	C	0.79	0/2139	0.90	7/2901 (0.2%)
1	D	0.76	0/2175	0.88	2/2954 (0.1%)
All	All	0.80	1/8650 (0.0%)	0.87	10/11738 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	175	GLU	CD-OE1	6.85	1.33	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	112	ARG	NE-CZ-NH2	-9.12	115.74	120.30
1	D	112	ARG	NE-CZ-NH2	-8.73	115.93	120.30
1	D	112	ARG	NE-CZ-NH1	7.92	124.26	120.30
1	C	112	ARG	NE-CZ-NH1	7.29	123.94	120.30
1	C	233	ARG	NE-CZ-NH2	7.04	123.82	120.30
1	C	258[A]	ASP	CB-CG-OD2	6.08	123.77	118.30
1	C	258[B]	ASP	CB-CG-OD2	6.08	123.77	118.30
1	A	142	ASP	CB-CG-OD2	5.71	123.44	118.30
1	C	195	ASP	CB-CG-OD2	5.23	123.01	118.30
1	C	102	ASP	CB-CG-OD1	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2088	0	0	7	0
1	B	2090	0	0	9	0
1	C	2066	0	0	5	0
1	D	2082	0	0	6	0
2	A	20	0	0	3	0
2	B	20	0	0	0	0
2	C	20	0	0	1	0
2	D	10	0	0	1	0
3	A	48	0	0	2	0
3	B	48	0	0	0	0
3	C	48	0	0	0	0
3	D	48	0	0	0	0
4	A	8	0	0	3	0
5	A	365	0	0	5	0
5	B	391	0	0	5	0
5	C	302	0	0	2	0
5	D	275	0	0	5	0
All	All	9929	0	0	31	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (31) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:1671:HOH:O	1:C:89[B]:MET:CE	1.68	1.29
2:A:1404:SO4:O3	5:A:1681:HOH:O	1.59	1.20
1:D:140[A]:SER:OG	5:D:1601:HOH:O	1.63	1.15
1:D:30[A]:GLU:OE2	5:D:1626:HOH:O	1.76	1.02
1:A:215[B]:TYR:CD2	1:A:238[B]:LEU:CD2	2.44	1.01
1:A:215[B]:TYR:CE2	1:A:238[B]:LEU:CD2	2.46	0.98
1:B:89[B]:MET:CE	5:D:1499:HOH:O	2.12	0.96
1:A:61:THR:OG1	4:A:1405:DTV:S4	2.25	0.94
1:A:215[A]:TYR:OH	5:A:1724:HOH:O	2.00	0.79
2:A:1404:SO4:S	5:A:1681:HOH:O	2.31	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:108[A]:SER:OG	5:A:1527:HOH:O	2.02	0.76
1:C:152[B]:VAL:CG1	5:C:1708:HOH:O	2.34	0.75
1:B:59[B]:ASN:ND2	5:B:1775:HOH:O	2.31	0.64
1:D:65:LYS:NZ	5:D:1597:HOH:O	2.32	0.62
1:C:89[A]:MET:CE	5:C:1626:HOH:O	2.47	0.62
1:A:2[A]:GLU:OE1	1:A:32:PRO:CB	2.49	0.61
1:B:267[B]:GLN:NE2	5:B:1686:HOH:O	2.33	0.61
1:C:61:THR:OG1	2:C:1423:SO4:O2	2.24	0.54
1:B:28:ASN:ND2	5:B:1539:HOH:O	2.45	0.49
1:D:196[A]:ILE:CG2	5:D:1625:HOH:O	2.60	0.49
3:A:1401:NAP:C5N	4:A:1405:DTV:C4	2.92	0.48
1:B:74:GLU:OE2	1:B:95:ARG:NH2	2.49	0.46
1:B:141:LEU:O	1:B:142:ASP:CB	2.64	0.45
1:B:59[B]:ASN:CG	5:B:1775:HOH:O	2.55	0.44
3:A:1401:NAP:C4N	4:A:1405:DTV:C4	2.94	0.44
1:D:61[A]:THR:OG1	2:D:1433:SO4:O4	2.35	0.44
1:C:177:GLU:OE2	1:C:202:SER:OG	2.36	0.43
1:A:16:SER:OG	2:A:1404:SO4:O4	2.38	0.42
1:D:215:TYR:CB	1:D:238:LEU:CD2	2.97	0.42
1:B:75:LEU:N	5:B:1550:HOH:O	2.54	0.41
1:B:196:ILE:CD1	1:B:225:TRP:CD1	3.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/271 (102%)	270 (98%)	5 (2%)	0	100	100
1	B	277/271 (102%)	269 (97%)	8 (3%)	0	100	100
1	C	273/271 (101%)	266 (97%)	7 (3%)	0	100	100
1	D	278/271 (103%)	274 (99%)	4 (1%)	0	100	100
All	All	1103/1084 (102%)	1079 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/217 (103%)	223 (100%)	0	100	100
1	B	225/217 (104%)	219 (97%)	6 (3%)	57	20
1	C	220/217 (101%)	218 (99%)	2 (1%)	87	69
1	D	224/217 (103%)	217 (97%)	7 (3%)	52	15
All	All	892/868 (103%)	877 (98%)	15 (2%)	78	41

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1	MET
1	B	2	GLU
1	B	33	TYR
1	B	141	LEU
1	B	255[A]	VAL
1	B	255[B]	VAL
1	C	1	MET
1	C	269	GLU
1	D	135	LEU
1	D	136	LEU
1	D	196[A]	ILE
1	D	196[B]	ILE
1	D	217	LYS
1	D	269[A]	GLU
1	D	269[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	A	1401	-	52,52,52	1.02	3 (5%)	80,80,80	1.64	7 (8%)
2	SO4	A	1402	-	4,4,4	0.43	0	6,6,6	0.45	0
2	SO4	A	1403	-	4,4,4	0.53	0	6,6,6	0.84	0
2	SO4	A	1404	-	4,4,4	0.81	0	6,6,6	0.66	0
4	DTV	A	1405	-	7,7,7	0.93	1 (14%)	8,8,8	3.39	5 (62%)
2	SO4	A	1424	-	4,4,4	0.34	0	6,6,6	0.52	0
3	NAP	B	1411	-	52,52,52	1.19	4 (7%)	80,80,80	1.81	7 (8%)
2	SO4	B	1412	-	4,4,4	0.12	0	6,6,6	0.25	0
2	SO4	B	1413	-	4,4,4	0.86	0	6,6,6	0.59	0
2	SO4	B	1414	-	4,4,4	1.05	0	6,6,6	0.85	0
2	SO4	B	1434	-	4,4,4	0.33	0	6,6,6	0.50	0
3	NAP	C	1421	-	52,52,52	1.34	5 (9%)	80,80,80	2.03	11 (13%)
2	SO4	C	1422	-	4,4,4	0.23	0	6,6,6	0.12	0
2	SO4	C	1423	-	4,4,4	0.52	0	6,6,6	0.66	0
2	SO4	C	1425	-	4,4,4	0.73	0	6,6,6	0.50	0
2	SO4	C	1426	-	4,4,4	0.54	0	6,6,6	0.60	0
3	NAP	D	1431	-	52,52,52	1.24	4 (7%)	80,80,80	1.75	9 (11%)
2	SO4	D	1432	-	4,4,4	0.17	0	6,6,6	0.15	0
2	SO4	D	1433	-	4,4,4	0.30	0	6,6,6	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	1401	-	-	0/35/67/67	0/3/5/5
2	SO4	A	1402	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1403	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1404	-	-	0/0/0/0	0/0/0/0
4	DTV	A	1405	-	-	0/8/8/8	0/0/0/0
2	SO4	A	1424	-	-	0/0/0/0	0/0/0/0
3	NAP	B	1411	-	-	0/35/67/67	0/3/5/5
2	SO4	B	1412	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1413	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1414	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1434	-	-	0/0/0/0	0/0/0/0
3	NAP	C	1421	-	-	0/35/67/67	0/3/5/5
2	SO4	C	1422	-	-	0/0/0/0	0/0/0/0
2	SO4	C	1423	-	-	0/0/0/0	0/0/0/0
2	SO4	C	1425	-	-	0/0/0/0	0/0/0/0
2	SO4	C	1426	-	-	0/0/0/0	0/0/0/0
3	NAP	D	1431	-	-	0/35/67/67	0/3/5/5
2	SO4	D	1432	-	-	0/0/0/0	0/0/0/0
2	SO4	D	1433	-	-	0/0/0/0	0/0/0/0

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1421	NAP	O7N-C7N	5.77	1.37	1.24
3	D	1431	NAP	O7N-C7N	5.11	1.36	1.24
3	A	1401	NAP	O7N-C7N	4.02	1.33	1.24
3	B	1411	NAP	O7N-C7N	4.02	1.33	1.24
3	B	1411	NAP	C2A-N3A	3.88	1.39	1.32
3	C	1421	NAP	C2A-N3A	3.75	1.39	1.32
3	D	1431	NAP	C2A-N1A	3.05	1.39	1.33
3	D	1431	NAP	C2A-N3A	2.87	1.37	1.32
3	B	1411	NAP	C8A-N9A	2.80	1.40	1.36
3	D	1431	NAP	PA-O3	2.74	1.64	1.59
3	C	1421	NAP	C2A-N1A	2.66	1.39	1.33
3	A	1401	NAP	C2A-N3A	2.62	1.37	1.32
3	B	1411	NAP	C2A-N1A	2.56	1.39	1.33
3	C	1421	NAP	PN-O3	2.32	1.65	1.60
3	C	1421	NAP	PA-O3	2.26	1.64	1.59
3	A	1401	NAP	C8A-N9A	2.25	1.40	1.36
4	A	1405	DTV	C1-C2	2.11	1.55	1.50

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1411	NAP	N3A-C2A-N1A	-11.48	119.11	128.71
3	C	1421	NAP	N3A-C2A-N1A	-11.44	119.15	128.71
3	D	1431	NAP	N3A-C2A-N1A	-9.97	120.37	128.71
3	A	1401	NAP	N3A-C2A-N1A	-8.34	121.73	128.71
3	C	1421	NAP	C3N-C7N-N7N	7.91	126.77	117.77
4	A	1405	DTV	O2-C2-C1	6.19	121.51	109.87
3	D	1431	NAP	C3N-C7N-N7N	5.87	124.45	117.77
3	A	1401	NAP	O7N-C7N-C3N	-5.87	112.96	119.58
3	B	1411	NAP	C3N-C7N-N7N	5.53	124.06	117.77
3	C	1421	NAP	O7N-C7N-C3N	-5.37	113.52	119.58
3	A	1401	NAP	C3N-C7N-N7N	4.80	123.24	117.77
3	B	1411	NAP	O7N-C7N-C3N	-4.62	114.37	119.58
4	A	1405	DTV	C3-C4-S4	4.22	121.31	114.66
3	D	1431	NAP	O7N-C7N-C3N	-4.17	114.87	119.58
3	C	1421	NAP	C8A-N9A-C4A	4.03	109.97	106.90
4	A	1405	DTV	O3-C3-C2	3.90	117.45	109.83
3	C	1421	NAP	N3A-C4A-N9A	3.66	132.05	125.43
4	A	1405	DTV	C4-C3-C2	-3.59	104.86	113.07
3	D	1431	NAP	C2N-C3N-C4N	3.42	122.19	118.31
3	B	1411	NAP	C2N-C3N-C4N	3.23	121.97	118.31
3	D	1431	NAP	N3A-C4A-N9A	3.16	131.13	125.43
3	A	1401	NAP	O4D-C1D-C2D	-2.92	102.29	106.77
3	C	1421	NAP	N7A-C8A-N9A	-2.79	106.47	114.36
3	A	1401	NAP	N3A-C4A-N9A	2.71	130.33	125.43
4	A	1405	DTV	O3-C3-C4	-2.42	105.31	109.87
3	C	1421	NAP	O4D-C1D-C2D	-2.33	103.20	106.77
3	D	1431	NAP	O4B-C1B-C2B	-2.33	104.77	106.95
3	D	1431	NAP	O3X-P2B-O2X	2.28	116.50	107.61
3	C	1421	NAP	C3B-C2B-C1B	-2.26	98.33	102.73
3	B	1411	NAP	N3A-C4A-N9A	2.26	129.51	125.43
3	A	1401	NAP	O4B-C1B-C2B	-2.24	104.85	106.95
3	C	1421	NAP	C1B-N9A-C4A	-2.20	122.83	126.64
3	B	1411	NAP	N7A-C8A-N9A	-2.15	108.27	114.36
3	A	1401	NAP	O3-PN-O1N	2.12	113.88	108.83
3	C	1421	NAP	C2A-N3A-C4A	2.12	120.03	114.01
3	B	1411	NAP	O2X-P2B-O2B	-2.08	101.11	107.09
3	D	1431	NAP	N7A-C8A-N9A	-2.05	108.57	114.36
3	D	1431	NAP	C5N-C4N-C3N	-2.04	117.67	120.32
3	C	1421	NAP	O7N-C7N-N7N	-2.01	119.70	122.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/271 (100%)	0.20	14 (5%) 26 29	6, 11, 23, 42	0
1	B	271/271 (100%)	0.21	11 (4%) 35 39	6, 11, 24, 43	0
1	C	269/271 (99%)	0.53	30 (11%) 6 6	7, 14, 28, 53	0
1	D	270/271 (99%)	0.56	32 (11%) 5 5	7, 16, 29, 54	0
All	All	1081/1084 (99%)	0.37	87 (8%) 12 14	6, 13, 27, 54	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	270	LEU	10.6
1	B	1	MET	9.5
1	A	1	MET	6.3
1	C	268	GLU	6.1
1	A	270	LEU	5.7
1	D	268	GLU	5.0
1	C	192	ILE	4.7
1	C	196	ILE	4.7
1	D	192	ILE	4.6
1	C	41	ASN	4.5
1	A	192	ILE	4.4
1	C	217	LYS	4.1
1	C	40	ILE	4.0
1	C	13	HIS	3.9
1	D	40	ILE	3.9
1	D	13	HIS	3.8
1	D	225	TRP	3.7
1	C	218	GLY	3.7
1	D	12	ALA	3.7
1	C	193	SER	3.7
1	B	271	SER	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	233	ARG	3.5
1	D	264	LYS	3.4
1	D	194	GLY	3.4
1	C	224	ALA	3.4
1	D	41[A]	ASN	3.4
1	C	195	ASP	3.3
1	A	174	ASP	3.3
1	A	271	SER	3.3
1	C	225	TRP	3.2
1	C	194	GLY	3.1
1	D	217	LYS	3.1
1	D	206	PRO	3.1
1	C	228	GLN	3.0
1	D	204	ILE	2.9
1	D	228	GLN	2.8
1	D	250	LEU	2.8
1	C	191	GLY	2.7
1	B	134	VAL	2.7
1	D	174	ASP	2.7
1	D	196[A]	ILE	2.7
1	B	103	GLY	2.7
1	C	267	GLN	2.6
1	C	190	SER	2.6
1	A	178	GLY	2.6
1	D	269[A]	GLU	2.6
1	A	106	LEU	2.6
1	C	214	PHE	2.6
1	B	106	LEU	2.5
1	C	264	LYS	2.5
1	C	25	GLN	2.5
1	C	12	ALA	2.5
1	A	104	VAL	2.4
1	D	136	LEU	2.4
1	B	104	VAL	2.4
1	D	38	ALA	2.4
1	C	250	LEU	2.4
1	A	134	VAL	2.4
1	A	217	LYS	2.4
1	D	231	SER	2.4
1	B	250	LEU	2.4
1	C	63	PRO	2.4
1	B	251	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	255[A]	VAL	2.3
1	C	39	PRO	2.3
1	C	232	LYS	2.3
1	D	218	GLY	2.2
1	C	265	GLN	2.2
1	B	142	ASP	2.2
1	A	101	THR	2.2
1	A	177	GLU	2.2
1	A	268	GLU	2.2
1	D	198	ALA	2.2
1	D	265	GLN	2.2
1	B	174	ASP	2.1
1	D	193	SER	2.1
1	D	224	ALA	2.1
1	D	39	PRO	2.1
1	A	142	ASP	2.1
1	C	230	GLY	2.1
1	C	38	ALA	2.1
1	B	270	LEU	2.1
1	C	269	GLU	2.1
1	D	227	GLU	2.0
1	C	174	ASP	2.0
1	D	62	VAL	2.0
1	D	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	DTV	A	1405	8/8	0.47	13.55	10,22,23,23	8
2	SO4	B	1434	5/5	0.22	7.23	26,26,28,29	5
2	SO4	C	1423	5/5	0.33	6.60	20,25,27,30	5
2	SO4	A	1404	5/5	0.21	5.73	22,23,26,30	5
2	SO4	C	1426	5/5	0.21	4.30	23,27,28,34	5
2	SO4	B	1414	5/5	0.24	4.07	16,17,21,25	5
2	SO4	A	1402	5/5	0.20	3.71	21,24,26,27	5
2	SO4	D	1433	5/5	0.20	3.40	32,33,35,35	5
2	SO4	D	1432	5/5	0.12	1.01	27,28,31,31	5
2	SO4	A	1424	5/5	0.11	0.98	23,24,26,26	5
2	SO4	C	1425	5/5	0.17	0.92	23,27,28,29	5
2	SO4	A	1403	5/5	0.13	0.91	22,24,28,30	0
2	SO4	B	1413	5/5	0.10	0.51	20,21,26,28	0
3	NAP	C	1421	48/48	0.13	0.03	14,24,31,33	0
2	SO4	B	1412	5/5	0.09	-0.13	24,25,25,27	5
2	SO4	C	1422	5/5	0.08	-0.37	21,21,25,25	5
3	NAP	D	1431	48/48	0.08	-0.57	13,17,21,23	0
3	NAP	B	1411	48/48	0.07	-0.79	7,10,13,15	0
3	NAP	A	1401	48/48	0.07	-0.79	8,11,14,17	0

6.5 Other polymers ⓘ

There are no such residues in this entry.