



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 10, 2014 – 11:08 AM EDT

PDB ID : 4O10  
Title : Structural and Biochemical Analyses of the Catalysis and Potency Impact of Inhibitor Phosphoribosylation by Human Nicotinamide Phosphoribosyltransferase  
Authors : Oh, A.; Wang, W.  
Deposited on : 2013-12-14  
Resolution : 1.55 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23161  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23161

# 1 Overall quality at a glance

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1117 (1.58-1.54)
Clashscore	79885	1249 (1.58-1.54)
Ramachandran outliers	78287	1212 (1.58-1.54)
Sidechain outliers	78261	1210 (1.58-1.54)
RSRZ outliers	66119	1117 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	501	
1	B	501	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	PO4	B	602	-	X
4	EDO	A	605	-	X
4	EDO	A	607	-	X
4	EDO	A	608	-	X
4	EDO	B	608	-	X
4	EDO	B	610	-	X

## 2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 8745 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nicotinamide phosphoribosyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	471	Total	C	N	O	S	0	0	0
			3765	2422	622	714	7			
1	B	469	Total	C	N	O	S	0	1	0
			3766	2422	624	713	7			

- Molecule 2 is N-{4-[(3,5-DIFLUOROPHENYL)SULFONYL]BENZYL}INDOLIZINE-7-CARBOXAMIDE (three-letter code: 2QF) (formula: C<sub>22</sub>H<sub>16</sub>F<sub>2</sub>N<sub>2</sub>O<sub>3</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0
			30	22	2	2	3	1	
2	B	1	Total	C	F	N	O	S	0
			30	22	2	2	3	1	

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	540	Total O 540 540	0	0
5	B	542	Total O 542 542	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.47Å 106.36Å 82.79Å 90.00° 96.35° 90.00°	Depositor
Resolution (Å)	46.22 – 1.55 46.21 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.1 (46.22-1.55) 98.1 (46.21-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.00 (at 1.55Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.151 , 0.179 0.150 , 0.179	Depositor DCC
$R_{free}$ test set	7393 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.5	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 41.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 147577 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8745	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2QF, PO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/3853	0.74	2/5220 (0.0%)
1	B	0.64	0/3854	0.71	2/5220 (0.0%)
All	All	0.65	0/7707	0.72	4/10440 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	311	ARG	NE-CZ-NH1	10.01	125.31	120.30
1	A	311	ARG	NE-CZ-NH2	-7.48	116.56	120.30
1	B	311	ARG	NE-CZ-NH1	7.39	124.00	120.30
1	B	311	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3765	0	3739	12	0
1	B	3766	0	3747	8	0
2	A	30	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	30	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	1	0
4	A	24	0	36	1	0
4	B	28	0	42	1	0
5	A	540	0	0	12	1
5	B	542	0	0	6	1
All	All	8745	0	7564	23	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (23) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:263:GLU:OE2	5:B:1122:HOH:O	2.10	0.69
1:A:235:ASP:OD1	5:A:1044:HOH:O	2.11	0.69
1:A:23:TYR:O	5:A:983:HOH:O	2.13	0.67
1:A:140:GLU:OE2	5:A:1237:HOH:O	2.13	0.66
1:A:177:LEU:HD22	1:A:484:ILE:HD11	1.79	0.63
1:A:363:GLU:OE2	5:A:1094:HOH:O	2.16	0.62
1:A:88:LYS:NZ	5:A:1168:HOH:O	2.27	0.62
1:B:187:GLU:H	1:B:187:GLU:CD	2.04	0.61
1:B:53:LYS:NZ	5:B:1152:HOH:O	2.31	0.61
1:B:255:LYS:O	5:B:832:HOH:O	2.16	0.61
1:A:439:ASN:HB3	5:A:1049:HOH:O	2.07	0.53
5:A:1029:HOH:O	4:B:606:EDO:H22	2.08	0.53
4:A:609:EDO:H22	5:B:1043:HOH:O	2.10	0.52
1:A:89:GLU:OE1	5:A:873:HOH:O	2.19	0.51
1:B:187:GLU:OE2	5:B:1209:HOH:O	2.20	0.49
1:A:42:LYS:NZ	5:A:985:HOH:O	2.41	0.46
1:A:184:ASP:O	5:A:1122:HOH:O	2.19	0.46
3:B:603:PO4:O3	5:B:1129:HOH:O	2.21	0.45
1:A:326:GLU:HG3	5:A:986:HOH:O	2.18	0.43
1:B:331:LYS:HD2	1:B:331:LYS:HA	1.78	0.43
1:B:409:LEU:HD13	1:B:410:GLY:O	2.20	0.41
1:A:257:HIS:HA	5:A:1031:HOH:O	2.20	0.41
1:B:53:LYS:HE2	1:B:53:LYS:HB3	1.39	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:1237:HOH:O	5:B:1138:HOH:O[2_646]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/501 (93%)	458 (98%)	9 (2%)	0	100	100
1	B	466/501 (93%)	457 (98%)	9 (2%)	0	100	100
All	All	933/1002 (93%)	915 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/440 (94%)	410 (99%)	3 (1%)	91	77
1	B	414/440 (94%)	408 (99%)	6 (1%)	78	52
All	All	827/880 (94%)	818 (99%)	9 (1%)	85	63

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	TYR
1	A	195	TYR
1	A	370	GLN
1	B	18	TYR
1	B	94	ASP
1	B	195	TYR
1	B	409	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	434[A]	ARG
1	B	434[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	2QF	A	601	-	33,33,33	1.22	3 (9%)	48,48,48	3.28	14 (29%)
3	PO4	A	602	-	4,4,4	0.42	0	6,6,6	0.31	0
3	PO4	A	603	-	4,4,4	0.80	0	6,6,6	0.29	0
4	EDO	A	604	-	3,3,3	0.58	0	2,2,2	0.34	0
4	EDO	A	605	-	3,3,3	0.54	0	2,2,2	0.44	0
4	EDO	A	606	-	3,3,3	0.58	0	2,2,2	0.42	0
4	EDO	A	607	-	3,3,3	0.66	0	2,2,2	0.15	0
4	EDO	A	608	-	3,3,3	0.48	0	2,2,2	0.73	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	609	-	3,3,3	0.48	0	2,2,2	0.22	0
2	2QF	B	601	-	33,33,33	1.14	2 (6%)	48,48,48	2.82	10 (20%)
3	PO4	B	602	-	4,4,4	1.46	1 (25%)	6,6,6	0.34	0
3	PO4	B	603	-	4,4,4	0.46	0	6,6,6	0.30	0
4	EDO	B	604	-	3,3,3	0.65	0	2,2,2	0.61	0
4	EDO	B	605	-	3,3,3	0.71	0	2,2,2	0.48	0
4	EDO	B	606	-	3,3,3	0.61	0	2,2,2	0.27	0
4	EDO	B	607	-	3,3,3	0.47	0	2,2,2	0.65	0
4	EDO	B	608	-	3,3,3	0.60	0	2,2,2	0.09	0
4	EDO	B	609	-	3,3,3	0.59	0	2,2,2	0.32	0
4	EDO	B	610	-	3,3,3	0.62	0	2,2,2	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2QF	A	601	-	-	0/21/21/21	0/4/4/4
3	PO4	A	602	-	-	0/0/0/0	0/0/0/0
3	PO4	A	603	-	-	0/0/0/0	0/0/0/0
4	EDO	A	604	-	-	0/1/1/1	0/0/0/0
4	EDO	A	605	-	-	0/1/1/1	0/0/0/0
4	EDO	A	606	-	-	0/1/1/1	0/0/0/0
4	EDO	A	607	-	-	0/1/1/1	0/0/0/0
4	EDO	A	608	-	-	0/1/1/1	0/0/0/0
4	EDO	A	609	-	-	0/1/1/1	0/0/0/0
2	2QF	B	601	-	-	0/21/21/21	0/4/4/4
3	PO4	B	602	-	-	0/0/0/0	0/0/0/0
3	PO4	B	603	-	-	0/0/0/0	0/0/0/0
4	EDO	B	604	-	-	0/1/1/1	0/0/0/0
4	EDO	B	605	-	-	0/1/1/1	0/0/0/0
4	EDO	B	606	-	-	0/1/1/1	0/0/0/0
4	EDO	B	607	-	-	0/1/1/1	0/0/0/0
4	EDO	B	608	-	-	0/1/1/1	0/0/0/0
4	EDO	B	609	-	-	0/1/1/1	0/0/0/0
4	EDO	B	610	-	-	0/1/1/1	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	2QF	C6-C5	4.17	1.46	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	2QF	C6-C5	3.61	1.45	1.39
2	A	601	2QF	C4-C3	3.23	1.42	1.40
3	B	602	PO4	P-O4	-2.70	1.42	1.52
2	B	601	2QF	C3-N8	-2.36	1.37	1.40
2	A	601	2QF	C7-N8	-2.18	1.36	1.39

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	2QF	C27-C28-C23	17.77	122.59	116.83
2	B	601	2QF	C27-C28-C23	14.44	121.51	116.83
2	B	601	2QF	C25-C24-C23	7.94	119.40	116.83
2	A	601	2QF	C25-C24-C23	7.74	119.34	116.83
2	A	601	2QF	C17-S20-C23	-4.99	95.84	104.58
2	B	601	2QF	C17-S20-C23	-4.82	96.13	104.58
2	B	601	2QF	O22-S20-C23	3.79	111.70	107.94
2	A	601	2QF	C2-C3-N8	-3.61	105.17	107.40
2	B	601	2QF	C28-C27-C26	-3.45	118.77	123.51
2	A	601	2QF	C18-C17-S20	3.18	123.82	119.58
2	A	601	2QF	C28-C27-C26	-3.01	119.37	123.51
2	A	601	2QF	O22-S20-C17	2.71	110.63	107.94
2	A	601	2QF	C13-N12-C10	-2.67	114.70	121.79
2	A	601	2QF	O21-S20-C23	-2.60	105.37	107.94
2	A	601	2QF	C4-C3-N8	2.58	120.74	118.21
2	B	601	2QF	O21-S20-C23	-2.50	105.47	107.94
2	B	601	2QF	C24-C23-S20	2.49	121.59	119.24
2	A	601	2QF	C28-C23-S20	2.48	121.58	119.24
2	A	601	2QF	O21-S20-C17	2.41	110.34	107.94
2	A	601	2QF	C9-N8-C3	2.37	111.36	108.12
2	B	601	2QF	O21-S20-O22	2.36	122.49	119.11
2	A	601	2QF	C5-C4-C3	-2.19	117.74	121.58
2	B	601	2QF	C13-N12-C10	-2.17	116.02	121.79
2	B	601	2QF	F29-C27-C28	2.06	121.23	118.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	471/501 (94%)	-0.43	3 (0%) 86 90	6, 13, 27, 60	0
1	B	469/501 (93%)	-0.38	2 (0%) 90 92	6, 12, 27, 58	0
All	All	940/1002 (93%)	-0.41	5 (0%) 88 91	6, 12, 27, 60	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	409	LEU	2.8
1	A	43	LYS	2.1
1	A	455	GLN	2.1
1	A	486	LEU	2.1
1	B	94	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	A	607	4/4	0.12	6.72	22,27,28,31	0
4	EDO	A	608	4/4	0.20	5.48	32,34,34,35	0
4	EDO	B	610	4/4	0.13	3.97	18,24,33,34	0
3	PO4	B	602	5/5	0.10	2.70	10,14,16,17	0
4	EDO	A	605	4/4	0.09	2.68	12,12,13,18	0
4	EDO	B	608	4/4	0.06	2.43	25,27,27,33	0
4	EDO	B	605	4/4	0.09	1.88	15,17,18,19	0
3	PO4	A	602	5/5	0.09	1.84	17,19,19,20	0
4	EDO	B	609	4/4	0.09	1.81	17,25,26,28	0
4	EDO	B	604	4/4	0.07	1.37	10,14,16,22	0
4	EDO	B	607	4/4	0.10	0.73	30,31,34,40	0
4	EDO	B	606	4/4	0.12	0.72	22,26,27,33	0
2	2QF	B	601	30/30	0.07	0.72	9,13,21,23	0
4	EDO	A	606	4/4	0.07	0.57	14,19,26,32	0
4	EDO	A	604	4/4	0.08	0.54	16,18,18,21	0
2	2QF	A	601	30/30	0.07	0.45	10,13,20,24	0
4	EDO	A	609	4/4	0.10	0.18	19,21,24,32	0
3	PO4	A	603	5/5	0.06	0.02	10,13,13,15	0
3	PO4	B	603	5/5	0.07	-0.02	13,14,17,17	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.