



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 03:05 PM GMT

PDB ID : 2O2S
Title : The structure of *T. gondii* enoyl acyl carrier protein reductase in complex with NAD and triclosan
Authors : Muench, S.P.; Prigge, S.T.; McLeod, R.; Rafferty, J.B.; Kirisits, M.J.; Roberts, C.W.; Mui, E.J.; Rice, D.W.
Deposited on : 2006-11-30
Resolution : 2.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4672 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-acyl carrier reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2255	1426	391	434	4			
1	B	302	Total	C	N	O	S	0	0	0
			2259	1431	392	432	4			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is TRICLOSAN (three-letter code: TCL) (formula: $C_{12}H_7Cl_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	O	0	0
			17	12	3	2		
3	B	1	Total	C	Cl	O	0	0
			17	12	3	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	16	Total	O	0	0
			16	16		
4	B	20	Total	O	0	0
			20	20		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enoyl-acyl carrier reductase

Chain A: 



- Molecule 1: Enoyl-acyl carrier reductase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.08Å 78.08Å 187.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.60 19.80 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.60) 99.8 (19.80-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.59Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.205 , 0.267 0.214 , 0.275	Depositor DCC
R_{free} test set	1077 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.0	EDS
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 21070 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4672	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TCL, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	1/2303 (0.0%)	0.76	3/3126 (0.1%)
1	B	0.65	0/2307	0.72	1/3130 (0.0%)
All	All	0.66	1/4610 (0.0%)	0.74	4/6256 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	SER	C-N	-8.69	1.14	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	304	MET	O-C-N	-7.53	106.79	121.10
1	A	259	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	B	184	ARG	NE-CZ-NH1	5.51	123.05	120.30
1	A	304	MET	CA-C-N	5.38	132.17	117.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	237	LYS	Peptide
1	A	4	PRO	Peptide
1	B	241	LYS	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2255	0	2228	35	0
1	B	2259	0	2240	32	1
2	A	44	0	26	3	0
2	B	44	0	26	0	0
3	A	17	0	6	1	0
3	B	17	0	6	0	0
4	A	16	0	0	1	0
4	B	20	0	0	1	0
All	All	4672	0	4532	65	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (65) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:304:MET:HB2	1:B:305:PRO:CD	1.52	1.37
1:B:304:MET:CB	1:B:305:PRO:HD3	1.65	1.12
1:B:304:MET:HB2	1:B:305:PRO:HD3	1.14	1.01
1:B:304:MET:CB	1:B:305:PRO:CD	2.31	1.00
1:B:304:MET:HB2	1:B:305:PRO:HD2	1.57	0.83
1:A:254:ASN:HD22	1:A:299:VAL:H	1.30	0.79
1:B:236:GLY:O	1:B:237:LYS:C	2.30	0.69
1:B:51:GLN:HE21	1:B:51:GLN:HA	1.63	0.63
1:B:124:LEU:HD22	1:B:168:MET:CE	2.29	0.63
1:A:5:ILE:CG2	1:A:277:LEU:HD13	2.30	0.62
1:B:183:GLU:OE2	1:B:287:TYR:OH	2.16	0.61
1:A:254:ASN:ND2	1:A:299:VAL:H	1.96	0.61
1:A:5:ILE:HG23	1:A:277:LEU:HD13	1.84	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:254:ASN:HD22	1:B:299:VAL:H	1.49	0.60
1:A:65:LEU:HD21	1:A:71:ILE:HB	1.83	0.58
1:A:259:ARG:NH2	1:B:279:ARG:HD3	2.18	0.58
1:A:259:ARG:HH21	1:B:279:ARG:HD3	1.68	0.58
1:A:161:LEU:HD23	1:A:207:LEU:HD22	1.86	0.57
1:B:29:LYS:HG2	1:B:71:ILE:HD11	1.88	0.55
1:B:164:PHE:O	1:B:168:MET:HG3	2.07	0.55
1:A:178:SER:HA	1:A:197:LYS:HD2	1.89	0.55
1:A:181:ALA:HA	1:A:184:ARG:O	2.08	0.53
1:A:44:PRO:HB3	1:A:102:VAL:HG21	1.90	0.53
1:A:254:ASN:ND2	1:A:299:VAL:N	2.56	0.53
1:A:10:GLN:HA	1:A:122:ASP:OD2	2.09	0.53
1:A:254:ASN:HD21	1:A:300:ASP:H	1.57	0.52
1:A:197:LYS:NZ	2:A:500:NAD:O2D	2.29	0.52
1:A:262:HIS:HB2	4:A:707:HOH:O	2.10	0.52
1:A:81:ALA:HB1	1:A:156:SER:HB2	1.93	0.50
1:B:306:PRO:HA	4:B:716:HOH:O	2.11	0.50
2:A:500:NAD:H2N	2:A:500:NAD:O1N	2.12	0.50
1:B:97:LYS:O	1:B:100:ALA:HB3	2.12	0.49
1:A:63:ARG:O	1:A:71:ILE:HG22	2.14	0.48
1:B:178:SER:HA	1:B:197:LYS:HD2	1.94	0.48
1:B:17:VAL:HG21	1:B:40:LEU:HD22	1.95	0.47
1:A:32:ALA:CB	1:A:71:ILE:HD13	2.45	0.47
1:B:305:PRO:HA	1:B:306:PRO:HD3	1.27	0.47
1:A:262:HIS:HD2	1:A:264:ASP:H	1.63	0.47
1:B:96:ASN:HD21	1:B:98:ARG:HD3	1.80	0.47
1:A:262:HIS:CD2	1:A:264:ASP:H	2.33	0.47
1:A:32:ALA:HB2	1:A:71:ILE:HD13	1.97	0.45
1:B:95:ASP:O	1:B:96:ASN:C	2.55	0.45
1:A:173:SER:HB3	1:A:274:LEU:HD23	1.99	0.45
1:A:103:ASP:OD1	1:A:104:GLY:N	2.50	0.45
1:A:220:ASN:OD1	1:A:284:VAL:HG22	2.17	0.44
1:B:177:LEU:HD23	1:B:222:ILE:HB	2.00	0.44
1:A:209:TRP:CH2	1:B:299:VAL:HG11	2.53	0.44
1:B:223:SER:HB3	1:B:287:TYR:CD2	2.53	0.44
1:B:85:LYS:HG2	1:B:88:ASP:OD2	2.18	0.43
1:A:108:LYS:O	1:A:112:VAL:HG23	2.17	0.43
2:A:500:NAD:H3D	3:A:705:TCL:CL16	2.55	0.43
1:A:206:THR:O	1:A:209:TRP:HB3	2.19	0.43
1:B:222:ILE:HG23	1:B:288:VAL:HG23	2.01	0.42
1:A:124:LEU:O	1:A:174:ALA:HA	2.18	0.42
1:A:87:GLU:OE1	1:A:87:GLU:N	2.43	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:124:LEU:O	1:B:174:ALA:HA	2.20	0.42
1:B:165:GLY:N	1:B:166:PRO:CD	2.83	0.42
1:B:304:MET:HE2	1:B:304:MET:HB3	1.70	0.41
1:B:241:LYS:HE3	1:B:245:ASP:HB3	2.02	0.41
1:A:266:VAL:O	1:A:267:GLY:C	2.59	0.41
1:A:65:LEU:HB3	1:A:66:PRO:CD	2.51	0.41
1:A:90:PRO:HG2	1:A:93:ILE:HD13	2.03	0.41
1:A:122:ASP:OD1	1:A:169:ASN:ND2	2.46	0.40
1:A:209:TRP:CZ2	1:B:299:VAL:HG11	2.57	0.40
1:B:123:ILE:HG23	1:B:274:LEU:HD22	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:300:ASP:OD2	1:B:303:SER:OG[5.675]	1.65	0.55

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/315 (96%)	282 (94%)	19 (6%)	0	100	100
1	B	298/315 (95%)	277 (93%)	17 (6%)	4 (1%)	18	35
All	All	599/630 (95%)	559 (93%)	36 (6%)	4 (1%)	30	58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	304	MET
1	B	305	PRO
1	B	96	ASN
1	B	156	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/242 (95%)	222 (96%)	8 (4%)	48	77
1	B	232/242 (96%)	224 (97%)	8 (3%)	49	78
All	All	462/484 (96%)	446 (96%)	16 (4%)	48	77

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	71	ILE
1	A	79	LEU
1	A	92	ASP
1	A	195	SER
1	A	251	SER
1	A	258	ARG
1	A	259	ARG
1	B	51	GLN
1	B	153	SER
1	B	161	LEU
1	B	240	GLU
1	B	271	LEU
1	B	293	HIS
1	B	303	SER
1	B	304	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	254	ASN
1	A	262	HIS
1	B	51	GLN
1	B	96	ASN
1	B	254	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	500	-	48,48,48	1.29	3 (6%)	73,73,73	2.23	13 (17%)
3	TCL	A	705	-	18,18,18	1.80	4 (22%)	25,25,25	1.31	5 (20%)
2	NAD	B	501	-	48,48,48	1.46	4 (8%)	73,73,73	2.35	18 (24%)
3	TCL	B	706	-	18,18,18	1.67	2 (11%)	25,25,25	1.30	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	500	-	-	0/30/62/62	0/3/5/5
3	TCL	A	705	-	-	0/4/4/4	0/2/2/2
2	NAD	B	501	-	-	0/30/62/62	0/3/5/5
3	TCL	B	706	-	-	0/4/4/4	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAD	C4A-N3A	7.11	1.46	1.35
2	A	500	NAD	C4A-N3A	6.33	1.45	1.35
3	B	706	TCL	C6-C5	4.65	1.48	1.40
3	A	705	TCL	C8-C9	4.32	1.47	1.39
3	A	705	TCL	C6-C5	4.12	1.47	1.40
3	B	706	TCL	C8-C9	3.71	1.46	1.39
2	B	501	NAD	C4A-N9A	-2.76	1.33	1.37
2	B	501	NAD	C2N-N1N	-2.45	1.32	1.35
2	B	501	NAD	C2D-C1D	-2.35	1.50	1.53
3	A	705	TCL	C9-CL16	2.33	1.79	1.73
3	A	705	TCL	C11-CL15	2.26	1.79	1.74
2	A	500	NAD	C4A-N9A	-2.19	1.34	1.37
2	A	500	NAD	C2D-C1D	-2.18	1.50	1.53

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAD	N3A-C2A-N1A	-11.01	119.51	128.71
2	A	500	NAD	N3A-C2A-N1A	-9.94	120.39	128.71
2	A	500	NAD	O4B-C1B-N9A	8.05	115.93	108.44
2	B	501	NAD	O4D-C1D-N1N	7.66	115.79	107.95
2	B	501	NAD	O4B-C1B-N9A	6.83	114.79	108.44
2	A	500	NAD	O4D-C1D-N1N	6.65	114.75	107.95
2	A	500	NAD	C4A-C5A-N7A	-4.71	105.48	109.52
2	B	501	NAD	C4A-C5A-N7A	-3.51	106.51	109.52
2	B	501	NAD	C2B-C1B-N9A	3.28	121.67	113.27
2	B	501	NAD	C2D-C3D-C4D	3.18	108.99	102.65
2	B	501	NAD	O7N-C7N-C3N	-3.18	116.00	119.58
2	B	501	NAD	O5B-C5B-C4B	-3.06	97.71	108.94
2	A	500	NAD	C2B-C3B-C4B	3.00	108.63	102.65
2	A	500	NAD	C5A-C4A-N9A	2.93	111.40	107.16
3	B	706	TCL	C3-C2-CL14	-2.90	114.56	119.34
2	A	500	NAD	C2D-C3D-C4D	2.83	108.30	102.65
2	B	501	NAD	C2B-C3B-C4B	2.73	108.09	102.65
2	A	500	NAD	C3N-C7N-N7N	2.70	120.84	117.77
2	B	501	NAD	O3D-C3D-C2D	-2.66	103.17	111.83
2	B	501	NAD	C1B-N9A-C4A	-2.65	122.06	126.64
3	B	706	TCL	C1-C6-C5	-2.53	116.96	119.82
2	A	500	NAD	C5A-C4A-N3A	-2.51	120.23	125.70
2	B	501	NAD	C3N-C7N-N7N	2.46	120.57	117.77
2	A	500	NAD	C4D-O4D-C1D	2.38	112.33	109.75
2	B	501	NAD	C8A-N9A-C1B	2.33	130.97	126.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAD	C2D-C1D-N1N	2.29	117.74	113.86
3	B	706	TCL	C1-C2-CL14	2.28	121.97	119.14
2	B	501	NAD	C5A-C4A-N9A	2.25	110.42	107.16
3	B	706	TCL	O7-C5-C6	2.23	120.88	116.33
2	B	501	NAD	O3-PN-O5D	-2.23	92.44	101.36
3	A	705	TCL	C1-C6-C5	-2.22	117.31	119.82
3	A	705	TCL	C3-C2-CL14	-2.21	115.69	119.34
2	A	500	NAD	C2D-C1D-N1N	2.20	117.58	113.86
3	A	705	TCL	C8-C9-CL16	2.15	122.26	119.45
3	A	705	TCL	O17-C6-C1	2.14	125.07	119.42
2	B	501	NAD	O4D-C1D-C2D	2.11	110.01	106.77
3	A	705	TCL	C13-C12-C11	2.05	121.50	119.22
2	A	500	NAD	O3D-C3D-C4D	-2.04	105.07	111.08
2	A	500	NAD	C2A-N3A-C4A	2.02	119.77	114.01
2	B	501	NAD	N7A-C8A-N9A	-2.02	108.64	114.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/315 (96%)	-0.20	7 (2%) 57 54	42, 54, 75, 80	0
1	B	302/315 (95%)	-0.17	9 (2%) 48 45	40, 54, 75, 83	0
All	All	605/630 (96%)	-0.18	16 (2%) 53 50	40, 54, 75, 83	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	4.7
1	B	3	PHE	4.1
1	B	66	PRO	3.5
1	B	240	GLU	3.4
1	A	238	SER	2.9
1	B	68	GLY	2.9
1	B	237	LYS	2.7
1	A	67	ASP	2.7
1	A	237	LYS	2.5
1	B	97	LYS	2.4
1	B	67	ASP	2.4
1	A	70	LEU	2.3
1	B	230	ARG	2.2
1	A	97	LYS	2.1
1	A	240	GLU	2.1
1	B	306	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	TCL	A	705	17/17	0.15	0.60	52,52,53,54	0
3	TCL	B	706	17/17	0.14	0.21	51,54,58,59	0
2	NAD	A	500	44/44	0.14	0.01	41,55,70,71	0
2	NAD	B	501	44/44	0.11	-0.67	44,51,59,63	0

6.5 Other polymers

There are no such residues in this entry.