



# wwPDB X-ray Structure Validation Summary Report

Aug 26, 2014 – 01:50 AM EDT

PDB ID : 4O41  
Title : Amide linked RNA  
Authors : Pallan, P.S.; Egli, M.  
Deposited on : 2013-12-18  
Resolution : 1.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23489  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23489

# 1 Overall quality at a glance

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1038 (1.26-1.14)
Clashscore	79885	1158 (1.26-1.14)
RSRZ outliers	66119	1038 (1.26-1.14)
RNA backbone	1838	1002 (2.92-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	12	
1	B	12	
1	C	12	
1	D	12	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1365 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called AMIDE LINKED RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			
1	B	12	Total	C	N	O	P	0	3	0
			320	145	55	106	14			
1	C	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			
1	D	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			

- Molecule 2 is STRONTIUM ION (three-letter code: SR) (formula: Sr).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Sr	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total	O	0	0
			71	71		
3	B	76	Total	O	0	0
			76	76		
3	C	71	Total	O	0	0
			71	71		
3	D	66	Total	O	0	0
			66	66		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AMIDE LINKED RNA

Chain A: 



- Molecule 1: AMIDE LINKED RNA

Chain B: 




- Molecule 1: AMIDE LINKED RNA

Chain C: 



- Molecule 1: AMIDE LINKED RNA

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	22.36Å 34.33Å 43.30Å 110.03° 94.24° 95.83°	Depositor
Resolution (Å)	28.00 – 1.20 20.69 – 1.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.00-1.20) 93.9 (20.69-1.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.81 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.147 , 0.212 0.146 , 0.207	Depositor DCC
$R_{free}$ test set	1753 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.5	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 73.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 35926 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	1365	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: URU, SR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.14	0/237	1.67	4/365 (1.1%)
1	B	1.64	2/303 (0.7%)	2.14	12/466 (2.6%)
1	C	1.07	1/237 (0.4%)	1.78	5/365 (1.4%)
1	D	1.21	1/237 (0.4%)	1.98	13/365 (3.6%)
All	All	1.31	4/1014 (0.4%)	1.92	34/1561 (2.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	204[A]	G	C2'-O2'	-15.67	1.21	1.41
1	B	204[B]	G	C2'-O2'	-15.67	1.21	1.41
1	D	404	G	C2'-O2'	-5.16	1.34	1.41
1	C	311	C	C2'-O2'	5.02	1.48	1.41

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	201	U	C5-C4-O4	-9.37	120.28	125.90
1	B	204[A]	G	C3'-C2'-C1'	-9.20	94.14	101.50
1	B	204[B]	G	C3'-C2'-C1'	-9.20	94.14	101.50
1	D	409	G	C2-N3-C4	8.39	116.09	111.90
1	B	209	G	C2-N3-C4	7.55	115.68	111.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	253	0	108	0	0
1	B	320	0	143	2	0
1	C	253	0	110	0	0
1	D	253	0	110	0	0
2	A	2	0	0	0	0
3	A	71	0	0	0	0
3	B	76	0	0	0	0
3	C	71	0	0	0	0
3	D	66	0	0	0	0
All	All	1365	0	471	2	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (2) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:204[B]:G:N9	1:B:204[B]:G:C1'	2.00	1.25
1:B:204[B]:G:N9	1:B:204[B]:G:O4'	2.46	0.48

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

There are no protein chains in this entry.

### 5.3.2 Protein sidechains ⓘ

There are no protein chains in this entry.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	8/12 (66%)	0	0
1	B	7/12 (58%)	0	0
1	C	8/12 (66%)	0	0
1	D	8/12 (66%)	0	0
All	All	31/48 (64%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	URU	A	106	1	40,42,43	1.49	3 (7%)	50,61,64	1.30	5 (10%)
1	URU	B	206[A]	1	0,5,43	0.00	-	0,3,64	0.00	-
1	URU	B	206[B]	1	0,5,43	0.00	-	0,3,64	0.00	-
1	URU	C	306	1	40,42,43	1.61	7 (17%)	50,61,64	2.00	12 (24%)
1	URU	D	406	1	40,42,43	1.68	6 (15%)	50,61,64	2.05	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	URU	A	106	1	-	0/17/54/55	0/4/4/4
1	URU	B	206[A]	1	-	0/0/1/55	0/0/0/4
1	URU	B	206[B]	1	-	0/0/1/55	0/0/0/4
1	URU	C	306	1	-	0/17/54/55	0/4/4/4
1	URU	D	406	1	-	0/17/54/55	0/4/4/4

The worst 5 of 16 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	106	URU	C5-C4	5.71	1.43	1.37
1	D	406	URU	OW'-CW'	5.46	1.56	1.43
1	C	306	URU	C5-C4	4.75	1.42	1.37
1	D	406	URU	C15-C14	4.72	1.42	1.37
1	D	406	URU	C5-C4	4.67	1.42	1.37

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	306	URU	C2-N1-C1'	-7.57	113.46	118.21
1	D	406	URU	O3-C-C6'	7.04	131.56	121.25
1	C	306	URU	C12-N11-CV'	-5.71	114.63	118.21
1	D	406	URU	C12-N11-CV'	-5.14	114.98	118.21
1	D	406	URU	O3-C-NZ'	-4.00	115.06	122.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	10/12 (83%)	-0.47	0 100 100	12, 15, 16, 18	0
1	B	11/12 (91%)	0.13	1 (9%) 9 10	12, 15, 18, 19	1 (9%)
1	C	10/12 (83%)	-0.26	0 100 100	12, 15, 19, 19	0
1	D	10/12 (83%)	-0.27	0 100 100	12, 17, 22, 23	0
All	All	41/48 (85%)	-0.21	1 (2%) 56 60	12, 15, 21, 23	1 (2%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	201	U	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	URU	D	406	39/40	0.08	2.79	11,14,18,22	0
1	URU	B	206[B]	7/40	0.07	-0.25	14,14,17,27	7
1	URU	B	206[A]	7/40	0.07	-0.55	12,15,18,20	7
1	URU	A	106	39/40	0.06	-1.08	9,11,15,17	0
1	URU	C	306	39/40	0.06	-1.47	11,13,18,19	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SR	A	701	1/1	0.03	-4.78	13,13,13,13	1
2	SR	A	702	1/1	0.02	-6.96	13,13,13,13	1

### 6.5 Other polymers

There are no such residues in this entry.