



wwPDB X-ray Structure Validation Summary Report

Mar 12, 2014 – 07:07 PM GMT

PDB ID : 4O4H
Title : Tubulin-Laulimalide complex
Authors : Prota, A.E.; Bargsten, K.; Northcote, P.T.; Marsh, M.; Altmann, K.H.; Miller, J.H.; Diaz, J.F.; Steinmetz, M.O.
Deposited on : 2013-12-18
Resolution : 2.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1323
EDS : trunk22714
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk22714

1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	451	
1	C	451	
2	B	445	
2	D	445	
3	E	143	
4	F	384	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
10	GOL	B	504	-	X
10	GOL	D	503	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
6	MG	A	502	-	X
6	MG	B	502	-	X
6	MG	C	502	-	X
6	MG	F	402	-	X
7	CA	A	503	-	X
9	LLM	B	503	-	X

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 18548 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	12	0
			3481	2214	584	658	25			
1	C	440	Total	C	N	O	S	0	19	0
			3519	2235	586	672	26			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	426	Total	C	N	O	S	0	10	0
			3397	2137	573	660	27			
2	D	427	Total	C	N	O	S	0	6	0
			3370	2117	571	653	29			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	123	Total	C	N	O	S	0	5	0
			1039	642	187	205	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	ILE	CLONING ARTIFACT	UNP P63043
E	4	ALA	SER	CLONING ARTIFACT	UNP P63043


- Molecule 4 is a protein called Tubulin-tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	348	Total	C	N	O	S	0	6	0
			2878	1852	487	525	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	380	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	381	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	382	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	383	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	384	HIS	-	EXPRESSION TAG	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	A	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total	Ca	0	0
			2	2		
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
8	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 9 is LAULIMALIDE (three-letter code: LLM) (formula: C₃₀H₄₂O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			37	30	7		


- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			6	3	3		
10	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is PHOSPHOMETHYLPHOSPHONICACID ADENYLATE ESTER (three-

letter code: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	110	Total	O	0	0
			110	110		
12	B	142	Total	O	0	0
			142	142		
12	C	231	Total	O	0	0
			231	231		
12	D	102	Total	O	0	0
			102	102		
12	E	31	Total	O	0	0
			31	31		
12	F	40	Total	O	0	0
			40	40		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain



- Molecule 2: Tubulin beta-2B chain




- Molecule 2: Tubulin beta-2B chain




- Molecule 3: Stathmin-4

Chain E:



- Molecule 4: Tubulin-tyrosine ligase

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.69Å 156.87Å 180.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.90 – 2.10 78.43 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (71.90-2.10) 99.8 (78.43-2.10)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.192 , 0.224 0.193 , 0.224	Depositor DCC
R_{free} test set	8656 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 172824 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18548	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GOL, MG, CA, GTP, ACP, LLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	1/3596 (0.0%)	0.51	1/4883 (0.0%)
1	C	0.33	0/3655	0.52	0/4965
2	B	0.31	0/3498	0.50	0/4738
2	D	0.31	0/3461	0.52	1/4689 (0.0%)
3	E	0.30	0/1063	0.42	0/1412
4	F	0.38	1/2961 (0.0%)	0.49	0/4002
All	All	0.33	2/18234 (0.0%)	0.50	2/24689 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	90	SER	CB-OG	-12.87	1.25	1.42
1	A	89	PRO	N-CD	5.28	1.55	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	338	LYS	CD-CE-NZ	-12.12	83.83	111.70
1	A	88	HIS	C-N-CD	5.62	140.19	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3481	0	0	16	2
1	C	3519	0	0	11	0
2	B	3397	0	11	22	0
2	D	3370	0	0	21	1
3	E	1039	0	0	1	0
4	F	2878	0	0	29	3
5	A	32	0	12	0	0
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
8	B	28	0	12	0	0
8	D	28	0	12	0	0
9	B	37	0	0	1	0
10	B	6	0	8	1	0
10	D	6	0	8	0	0
11	F	31	0	14	5	0
12	A	110	0	0	8	0
12	B	142	0	0	11	0
12	C	231	0	0	8	0
12	D	102	0	0	9	0
12	E	31	0	0	0	0
12	F	40	0	0	3	0
All	All	18548	0	89	103	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:163:LYS:NZ	12:C:770:HOH:O	2.09	0.85
2:B:320:ARG:NH1	10:B:504:GOL:O3	2.09	0.84
4:F:146:VAL:O	4:F:184:LYS:NZ	2.10	0.84
2:D:164:ARG:NH1	12:D:616:HOH:O	2.11	0.82
1:C:293[A]:ASN:OD1	1:C:339:ARG:NH1	2.13	0.82

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:88:HIS:CE1	4:F:142:ARG:NH2[2.564]	1.14	1.06
1:A:88:HIS:ND1	4:F:142:ARG:NH2[2.564]	1.95	0.25
2:D:338:LYS:NZ	4:F:90:SER:OG[3.545]	2.05	0.15

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/451 (100%)	440 (98%)	9 (2%)	0	100	100
1	C	457/451 (101%)	447 (98%)	10 (2%)	0	100	100
2	B	432/445 (97%)	417 (96%)	14 (3%)	1 (0%)	56	57
2	D	429/445 (96%)	410 (96%)	18 (4%)	1 (0%)	56	57
3	E	124/143 (87%)	123 (99%)	1 (1%)	0	100	100
4	F	346/384 (90%)	327 (94%)	19 (6%)	0	100	100
All	All	2237/2319 (96%)	2164 (97%)	71 (3%)	2 (0%)	59	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	283	TYR
2	D	181	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/379 (101%)	372 (97%)	10 (3%)	59 62
1	C	390/379 (103%)	380 (97%)	10 (3%)	59 62
2	B	378/383 (99%)	365 (97%)	13 (3%)	49 49
2	D	374/383 (98%)	359 (96%)	15 (4%)	42 41
3	E	115/127 (91%)	113 (98%)	2 (2%)	73 78
4	F	320/342 (94%)	286 (89%)	34 (11%)	10 5
All	All	1959/1993 (98%)	1875 (96%)	84 (4%)	40 37

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	139	HIS
2	D	415	GLU
4	F	260	ASN
2	D	247	GLN
2	D	280	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GTP	A	501	6	34,34,34	0.95	1 (2%)	52,54,54	1.60	7 (13%)
8	GDP	B	501	6	30,30,30	1.28	5 (16%)	45,47,47	3.55	8 (17%)
9	LLM	B	503	-	40,40,40	3.30	14 (35%)	55,55,55	1.92	14 (25%)
10	GOL	B	504	-	5,5,5	0.39	0	5,5,5	0.41	0
5	GTP	C	501	6	34,34,34	1.07	1 (2%)	52,54,54	1.78	7 (13%)
8	GDP	D	501	6	30,30,30	1.26	4 (13%)	45,47,47	3.17	8 (17%)
10	GOL	D	503	-	5,5,5	0.40	0	5,5,5	0.38	0
11	ACP	F	401	-	33,33,33	2.38	12 (36%)	52,52,52	4.60	25 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	0/22/38/38	0/3/3/3
8	GDP	B	501	6	-	0/16/32/32	0/3/3/3
9	LLM	B	503	-	-	0/39/64/64	0/1/4/4
10	GOL	B	504	-	-	0/4/4/4	0/0/0/0
5	GTP	C	501	6	-	0/22/38/38	0/3/3/3
8	GDP	D	501	6	-	0/16/32/32	0/3/3/3
10	GOL	D	503	-	-	0/4/4/4	0/0/0/0
11	ACP	F	401	-	-	0/20/38/38	0/3/3/3

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	B	503	LLM	C14-C13	-11.53	1.39	1.51
9	B	503	LLM	C12-C13	-10.67	1.40	1.51
9	B	503	LLM	C21-C22	6.21	1.52	1.32
11	F	401	ACP	PG-O1G	5.13	1.61	1.50
9	B	503	LLM	C30-C25	-5.02	1.38	1.50

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
8	B	501	GDP	C6-C5-N7	20.55	136.91	134.14
8	D	501	GDP	C6-C5-N7	17.24	136.46	134.14
11	F	401	ACP	O4'-C1'-C2'	-12.72	88.17	106.69
11	F	401	ACP	N3-C2-N1	-12.45	117.93	128.89
11	F	401	ACP	O4'-C4'-C5'	11.57	150.47	109.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/451 (97%)	0.17	7 (1%) 68 72	26, 44, 80, 131	0
1	C	440/451 (97%)	0.11	3 (0%) 84 88	22, 34, 58, 99	0
2	B	426/445 (95%)	0.20	5 (1%) 75 80	25, 41, 72, 126	2 (0%)
2	D	427/445 (95%)	0.33	17 (3%) 36 41	28, 47, 78, 123	6 (1%)
3	E	123/143 (86%)	0.49	2 (1%) 68 72	34, 56, 93, 139	0
4	F	348/384 (90%)	1.50	105 (30%) 1 1	34, 69, 131, 174	0
All	All	2203/2319 (94%)	0.42	139 (6%) 19 22	22, 45, 96, 174	8 (0%)

The worst 5 of 139 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	105	LEU	10.5
4	F	249	TYR	8.1
4	F	104	ASN	7.0
4	F	173	ILE	6.4
4	F	182	ILE	6.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	MG	F	402	1/1	0.26	19.81	30,30,30,30	0
10	GOL	D	503	6/6	0.42	11.64	63,74,92,100	0
6	MG	C	502	1/1	0.22	10.82	34,34,34,34	0
6	MG	B	502	1/1	0.20	7.35	22,22,22,22	0
10	GOL	B	504	6/6	0.23	6.55	55,62,66,75	0
6	MG	A	502	1/1	0.23	6.35	38,38,38,38	0
9	LLM	B	503	37/37	0.23	2.70	91,93,102,104	0
7	CA	A	503	1/1	0.16	2.53	80,80,80,80	0
5	GTP	C	501	32/32	0.13	0.59	21,26,32,36	0
7	CA	B	505	1/1	0.17	0.52	109,109,109,109	0
5	GTP	A	501	32/32	0.14	0.33	25,30,34,43	0
8	GDP	D	501	28/28	0.13	-0.24	34,44,53,59	0
8	GDP	B	501	28/28	0.13	-0.29	21,27,33,37	0
11	ACP	F	401	31/31	0.16	-1.28	49,73,130,140	0
6	MG	D	502	1/1	0.07	-1.66	44,44,44,44	0
7	CA	B	506	1/1	0.09	-5.10	93,93,93,93	0

6.5 Other polymers ⓘ

There are no such residues in this entry.