



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 09:12 PM GMT

PDB ID : 3O52
Title : Structure of the E.coli GDP-mannose hydrolase (yffh) in complex with tartrate
Authors : Amzel, L.M.; Gabelli, S.B.; Boto, A.N.
Deposited on : 2010-07-27
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	191	
1	B	191	
1	C	191	
1	D	191	
1	E	191	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	TAR	A	7112	-	X
2	TAR	C	7110	-	X
2	TAR	D	7110	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
3	NA	A	7113	-	X
3	NA	A	7114	-	X
3	NA	B	7111	-	X
4	CL	D	7113	-	X

2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 7421 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDP-mannose pyrophosphatase nudK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	1	0
			1420	904	237	274	5			
1	B	176	Total	C	N	O	S	0	0	0
			1420	902	236	277	5			
1	C	177	Total	C	N	O	S	0	0	0
			1430	908	239	278	5			
1	D	180	Total	C	N	O	S	0	0	0
			1455	922	245	283	5			
1	E	177	Total	C	N	O	S	0	0	0
			1429	908	238	278	5			

- Molecule 2 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	4	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	4	6		
2	B	1	Total	C	O	0	0
			10	4	6		
2	C	1	Total	C	O	0	0
			10	4	6		
2	D	1	Total	C	O	0	0
			10	4	6		
2	E	1	Total	C	O	0	0
			10	4	6		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	2	Total	Na	0	0
			2	2		
3	D	2	Total	Na	0	0
			2	2		
3	E	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	2	Total	Cl	0	0
			2	2		
4	C	1	Total	Cl	0	0
			1	1		
4	E	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	40	Total	O	0	0
			40	40		
5	B	53	Total	O	0	0
			53	53		
5	C	38	Total	O	0	0
			38	38		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	33	Total 33	O 33	0	0
5	E	33	Total 33	O 33	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GDP-mannose pyrophosphatase nudK

Chain A: 



- Molecule 1: GDP-mannose pyrophosphatase nudK

Chain B: 




- Molecule 1: GDP-mannose pyrophosphatase nudK

Chain C: 




- Molecule 1: GDP-mannose pyrophosphatase nudK

Chain D: 



- Molecule 1: GDP-mannose pyrophosphatase nudK

Chain E: 





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	75.34Å 103.22Å 254.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.73 – 2.50 29.73 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.2 (29.73-2.50) 96.3 (29.73-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.211 , 0.288 0.203 , 0.281	Depositor DCC
R_{free} test set	1684 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 33382 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7421	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TAR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1444	0.70	1/1948 (0.1%)
1	B	0.69	0/1439	0.75	1/1940 (0.1%)
1	C	0.63	0/1451	0.73	1/1958 (0.1%)
1	D	0.81	6/1476 (0.4%)	0.76	3/1991 (0.2%)
1	E	1.34	20/1450 (1.4%)	1.14	10/1957 (0.5%)
All	All	0.87	26/7260 (0.4%)	0.83	16/9794 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	138	ASP	CG-OD2	19.61	1.70	1.25
1	E	31	ASP	CG-OD2	14.81	1.59	1.25
1	E	138	ASP	C-O	14.38	1.50	1.23
1	E	190	MET	CG-SD	12.80	2.14	1.81
1	D	30	LYS	CD-CE	11.61	1.80	1.51
1	D	121	SER	C-N	10.28	1.53	1.34
1	E	136	TYR	CE2-CZ	9.59	1.51	1.38
1	E	136	TYR	CG-CD1	8.14	1.49	1.39
1	D	30	LYS	CE-NZ	7.98	1.69	1.49
1	E	138	ASP	CB-CG	7.80	1.68	1.51
1	E	58	LYS	CE-NZ	7.72	1.68	1.49
1	E	56	LYS	C-O	6.99	1.36	1.23
1	E	54	ASN	CG-ND2	6.90	1.50	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	54	ASN	CB-CG	6.81	1.66	1.51
1	E	59	THR	CB-CG2	6.55	1.74	1.52
1	E	55	THR	C-O	6.14	1.35	1.23
1	E	108	GLU	CD-OE1	5.64	1.31	1.25
1	E	135	GLU	CD-OE1	5.64	1.31	1.25
1	E	56	LYS	CB-CG	5.50	1.67	1.52
1	D	30	LYS	CG-CD	5.47	1.71	1.52
1	E	136	TYR	CG-CD2	5.36	1.46	1.39
1	E	32	GLY	C-O	5.26	1.32	1.23
1	E	57	LYS	CD-CE	5.24	1.64	1.51
1	E	157	GLU	CB-CG	5.19	1.62	1.52
1	D	84	CYS	CB-SG	-5.08	1.73	1.81
1	D	151	GLU	CD-OE1	5.03	1.31	1.25

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	138	ASP	CB-CG-OD2	-23.15	97.47	118.30
1	E	31	ASP	CB-CG-OD1	17.11	133.70	118.30
1	E	138	ASP	CA-C-O	-12.08	94.72	120.10
1	C	27	LEU	CA-CB-CG	9.20	136.46	115.30
1	E	190	MET	CG-SD-CE	-7.53	88.15	100.20
1	E	138	ASP	OD1-CG-OD2	6.93	136.46	123.30
1	D	121	SER	O-C-N	-6.88	108.02	121.10
1	D	30	LYS	CD-CE-NZ	-6.82	96.01	111.70
1	E	31	ASP	OD1-CG-OD2	-6.61	110.75	123.30
1	E	138	ASP	CB-CG-OD1	6.29	123.96	118.30
1	E	32	GLY	O-C-N	6.24	132.69	122.70
1	E	56	LYS	O-C-N	5.94	132.20	122.70
1	D	120	MET	CA-C-N	-5.76	104.53	117.20
1	A	128	LEU	CA-CB-CG	5.67	128.34	115.30
1	E	190	MET	CB-CG-SD	-5.21	96.77	112.40
1	B	158	LEU	CA-CB-CG	5.10	127.04	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	32	GLY	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1420	0	0	9	0
1	B	1420	0	7	12	0
1	C	1430	0	7	7	0
1	D	1455	0	2	16	0
1	E	1429	0	14	14	0
2	A	20	0	8	2	0
2	B	10	0	4	4	0
2	C	10	0	4	0	0
2	D	10	0	3	0	0
2	E	10	0	4	1	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	D	2	0	0	0	0
3	E	1	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
4	E	1	0	0	0	0
5	A	40	0	0	2	0
5	B	53	0	0	4	0
5	C	38	0	0	1	0
5	D	33	0	0	2	0
5	E	33	0	0	1	0
All	All	7421	0	53	56	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (56) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:30:LYS:CD	1:D:30:LYS:CE	1.80	1.60
1:D:30:LYS:NZ	1:D:30:LYS:CE	1.69	1.54
1:E:58:LYS:NZ	1:E:58:LYS:CE	1.68	1.53
1:E:190:MET:CG	1:E:190:MET:SD	2.14	1.36
1:E:138:ASP:CG	1:E:138:ASP:OD2	1.70	1.29

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:44:ARG:NH1	2:B:7110:TAR:O41	1.67	1.26
1:D:31:ASP:OD1	1:D:32:GLY:N	2.13	0.82
1:D:25:TYR:OH	1:D:39:ARG:NH1	2.14	0.80
1:D:30:LYS:CE	1:D:30:LYS:CG	2.63	0.77
1:B:21:HIS:CE1	5:B:208:HOH:O	2.41	0.72
1:E:58:LYS:NZ	1:E:58:LYS:CD	2.54	0.71
1:E:190:MET:CB	1:E:190:MET:SD	2.80	0.69
1:B:31:ASP:O	1:B:33:GLU:N	2.27	0.67
1:B:44:ARG:NH1	2:B:7110:TAR:C4	2.56	0.67
1:D:4:GLN:N	5:D:300:HOH:O	2.27	0.67
2:A:7110:TAR:HO3	1:B:123:GLY:N	1.92	0.67
1:C:120:MET:CE	1:D:175:GLY:CA	2.73	0.67
1:D:30:LYS:CD	1:D:30:LYS:NZ	2.59	0.66
1:D:151:GLU:CD	1:D:151:GLU:N	2.51	0.64
1:A:44:ARG:NH2	5:A:195:HOH:O	2.31	0.63
1:C:117:GLU:CG	5:C:201:HOH:O	2.46	0.63
1:A:37:HIS:CD2	1:E:91:ASP:OD2	2.52	0.62
1:E:59:THR:OG1	1:E:136:TYR:OH	2.19	0.60
1:B:90:ASN:ND2	5:B:207:HOH:O	2.34	0.59
1:A:59:THR:OG1	1:A:136:TYR:OH	2.21	0.57
1:E:190:MET:CG	1:E:190:MET:CE	2.83	0.57
1:E:138:ASP:CB	1:E:138:ASP:OD2	2.54	0.54
1:B:10:ASP:OD1	1:B:21:HIS:CD2	2.60	0.54
1:A:10:ASP:OD1	1:A:21:HIS:CD2	2.61	0.53
1:C:113:ARG:NH1	1:C:113:ARG:CG	2.71	0.53
1:B:138:ASP:CA	1:B:139:ASN:C	2.78	0.52
1:E:99:LYS:NZ	5:E:206:HOH:O	2.44	0.50
1:E:54:ASN:OD1	1:E:57:LYS:N	2.45	0.50
1:E:136:TYR:CD1	1:E:136:TYR:C	2.87	0.47
1:C:152:ASP:N	1:C:152:ASP:OD2	2.48	0.47
1:A:123:GLY:N	2:B:7110:TAR:HO2	2.12	0.46
1:D:29:ARG:O	1:D:31:ASP:O	2.33	0.46
1:A:136:TYR:OH	1:A:157:GLU:OE2	2.34	0.46
1:B:173:ARG:N	5:B:198:HOH:O	2.47	0.46
1:C:114:LYS:NZ	1:C:130:HIS:CD2	2.83	0.46
1:A:182:ASN:ND2	5:B:201:HOH:O	2.49	0.45
1:A:44:ARG:NH1	2:A:7110:TAR:O4	2.50	0.45
1:D:67:ARG:NH2	1:D:82:GLU:OE2	2.49	0.45
1:D:176:LYS:N	5:D:194:HOH:O	2.48	0.44
1:B:39:ARG:NH2	1:B:123:GLY:O	2.50	0.44
1:C:31:ASP:O	1:C:33:GLU:N	2.51	0.43
1:C:85:ALA:O	1:C:100:GLU:OE1	2.37	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:E:127:GLU:OE2	2:E:7110:TAR:H3	2.18	0.42
1:E:187:SER:OG	1:E:187:SER:O	2.37	0.42
1:D:75:ASN:CG	1:D:173:ARG:NH1	2.73	0.42
1:D:49:THR:OG1	1:D:84:CYS:O	2.37	0.42
1:B:127:GLU:CD	2:B:7110:TAR:H3	2.40	0.42
1:D:139:ASN:N	1:D:139:ASN:OD1	2.52	0.42
1:A:21:HIS:CE1	5:A:215:HOH:O	2.72	0.41
1:B:152:ASP:OD2	1:B:152:ASP:C	2.59	0.41
1:D:114:LYS:NZ	1:D:130:HIS:CD2	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/191 (90%)	158 (92%)	11 (6%)	2 (1%)	19	32
1	B	170/191 (89%)	163 (96%)	4 (2%)	3 (2%)	13	20
1	C	173/191 (91%)	163 (94%)	8 (5%)	2 (1%)	19	32
1	D	176/191 (92%)	166 (94%)	9 (5%)	1 (1%)	33	55
1	E	173/191 (91%)	154 (89%)	16 (9%)	3 (2%)	14	22
All	All	863/955 (90%)	804 (93%)	48 (6%)	11 (1%)	18	29

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	30	LYS
1	A	30	LYS
1	A	32	GLY
1	B	32	GLY
1	E	110	GLY
1	C	32	GLY
1	C	138	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	127	GLU
1	E	188	HIS
1	B	124	GLY
1	E	124	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/168 (94%)	145 (92%)	13 (8%)	17	29
1	B	158/168 (94%)	152 (96%)	6 (4%)	44	71
1	C	159/168 (95%)	148 (93%)	11 (7%)	22	39
1	D	161/168 (96%)	151 (94%)	10 (6%)	26	45
1	E	159/168 (95%)	145 (91%)	14 (9%)	14	26
All	All	795/840 (95%)	741 (93%)	54 (7%)	22	39

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	14	SER
1	A	39	ARG
1	A	44	ARG
1	A	54	ASN
1	A	58	LYS
1	A	63	ILE
1	A	108	GLU
1	A	116	PHE
1	A	135	GLU
1	A	158	LEU
1	A	169	THR
1	A	191	ASP
1	B	33	GLU
1	B	44	ARG
1	B	68	VAL
1	B	109	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	116	PHE
1	B	158	LEU
1	C	4	GLN
1	C	27	LEU
1	C	31	ASP
1	C	33	GLU
1	C	68	VAL
1	C	84	CYS
1	C	113	ARG
1	C	116	PHE
1	C	138	ASP
1	C	152	ASP
1	C	158	LEU
1	D	11	LYS
1	D	33	GLU
1	D	56	LYS
1	D	68	VAL
1	D	116	PHE
1	D	141	ARG
1	D	151	GLU
1	D	165	GLU
1	D	182	ASN
1	D	191	ASP
1	E	6	THR
1	E	22	ASN
1	E	31	ASP
1	E	39	ARG
1	E	44	ARG
1	E	76	GLU
1	E	90	ASN
1	E	98	ARG
1	E	99	LYS
1	E	116	PHE
1	E	152	ASP
1	E	164	LEU
1	E	165	GLU
1	E	173	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 10 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TAR	A	7110	3	9,9,9	1.20	0	12,12,12	1.07	0
2	TAR	A	7112	-	9,9,9	0.99	0	12,12,12	0.86	0
2	TAR	B	7110	-	9,9,9	1.04	0	12,12,12	1.36	2 (16%)
2	TAR	C	7110	-	9,9,9	1.01	0	12,12,12	1.31	2 (16%)
2	TAR	D	7110	3	9,9,9	1.18	1 (11%)	12,12,12	1.12	1 (8%)
2	TAR	E	7110	3	9,9,9	1.01	0	12,12,12	1.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TAR	A	7110	3	-	0/12/12/12	0/0/0/0
2	TAR	A	7112	-	-	0/12/12/12	0/0/0/0
2	TAR	B	7110	-	-	0/12/12/12	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TAR	C	7110	-	-	0/12/12/12	0/0/0/0
2	TAR	D	7110	3	-	0/12/12/12	0/0/0/0
2	TAR	E	7110	3	-	0/12/12/12	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	7110	TAR	C2-C1	-2.04	1.50	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7110	TAR	O41-C4-C3	2.36	120.78	113.89
2	B	7110	TAR	O11-C1-C2	2.29	120.57	113.89
2	D	7110	TAR	O11-C1-C2	2.27	120.50	113.89
2	B	7110	TAR	O1-C1-C2	-2.03	115.21	121.15
2	C	7110	TAR	O11-C1-C2	2.00	119.73	113.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/191 (91%)	0.17	8 (4%) 31 32	32, 48, 75, 87	0
1	B	176/191 (92%)	-0.15	4 (2%) 57 60	30, 44, 71, 83	0
1	C	177/191 (92%)	0.18	8 (4%) 32 33	34, 50, 74, 87	0
1	D	180/191 (94%)	0.28	11 (6%) 21 20	37, 54, 83, 109	0
1	E	177/191 (92%)	0.27	9 (5%) 27 27	29, 58, 105, 121	0
All	All	885/955 (92%)	0.15	40 (4%) 28 33	29, 50, 84, 121	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	142	ALA	8.7
1	A	31	ASP	5.8
1	E	30	LYS	5.8
1	A	32	GLY	4.8
1	E	31	ASP	4.7
1	E	2	THR	4.6
1	B	139	ASN	4.2
1	E	108	GLU	3.9
1	E	32	GLY	3.8
1	C	139	ASN	3.7
1	D	191	ASP	3.7
1	A	30	LYS	3.4
1	D	141	ARG	3.4
1	A	29	ARG	3.1
1	D	29	ARG	3.0
1	C	110	GLY	3.0
1	D	140	GLN	2.9
1	C	179	LEU	2.9
1	D	30	LYS	2.8
1	E	29	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	139	ASN	2.7
1	C	76	GLU	2.7
1	D	179	LEU	2.7
1	C	30	LYS	2.6
1	A	76	GLU	2.6
1	C	108	GLU	2.6
1	E	111	GLU	2.5
1	E	191	ASP	2.5
1	D	108	GLU	2.4
1	A	33	GLU	2.4
1	D	138	ASP	2.3
1	E	58	LYS	2.3
1	B	138	ASP	2.3
1	A	56	LYS	2.2
1	C	177	THR	2.2
1	A	57	LYS	2.1
1	B	108	GLU	2.1
1	B	31	ASP	2.0
1	C	107	TYR	2.0
1	D	56	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	A	7114	1/1	0.52	20.20	58,58,58,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	TAR	A	7112	10/10	0.36	13.22	90,92,92,92	0
2	TAR	C	7110	10/10	0.55	10.18	92,93,93,93	0
4	CL	D	7113	1/1	0.34	7.93	71,71,71,71	1
3	NA	A	7113	1/1	0.29	4.94	61,61,61,61	0
2	TAR	D	7110	10/10	0.25	3.63	69,73,74,74	0
3	NA	B	7111	1/1	0.34	2.58	48,48,48,48	0
4	CL	E	7112	1/1	0.18	2.00	77,77,77,77	0
2	TAR	A	7110	10/10	0.22	1.69	49,53,55,57	0
2	TAR	B	7110	10/10	0.17	0.80	57,59,62,62	0
3	NA	D	7112	1/1	0.16	0.42	59,59,59,59	0
2	TAR	E	7110	10/10	0.15	-0.40	53,55,56,57	0
3	NA	E	7111	1/1	0.13	-1.01	43,43,43,43	0
3	NA	D	7111	1/1	0.10	-1.10	63,63,63,63	0
4	CL	D	7114	1/1	0.23	-	78,78,78,78	0
4	CL	C	192	1/1	0.42	-	69,69,69,69	0

6.5 Other polymers ⓘ

There are no such residues in this entry.