



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 08:56 PM GMT

PDB ID : 3O6B
Title : A Dual E3 Mechanism for Rub1 Ligation to Cdc53: Dcn1(P)-Cdc53(WHB)
low resolution
Authors : Scott, D.C.; Monda, J.K; Grace, C.R.R.; Duda, D.M.; Kriwacki, R.W.; Kurz,
T.; Schulman, B.A.
Deposited on : 2010-07-28
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1007 (3.18-3.02)
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)
RSRZ outliers	66119	1008 (3.18-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	202	
1	C	202	
1	E	202	
1	G	202	
1	I	202	
2	B	76	
2	D	76	
2	F	76	
2	H	76	
2	J	76	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10662 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Defective in cullin neddylation protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1664	1090	256	310	8			
1	C	198	Total	C	N	O	S	0	0	0
			1689	1107	259	315	8			
1	E	195	Total	C	N	O	S	0	0	0
			1667	1094	255	310	8			
1	G	192	Total	C	N	O	S	0	0	0
			1639	1074	251	306	8			
1	I	157	Total	C	N	O	S	0	0	0
			1348	888	208	246	6			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	68	GLY	-	EXPRESSION TAG	UNP Q12395
A	69	SER	-	EXPRESSION TAG	UNP Q12395
C	68	GLY	-	EXPRESSION TAG	UNP Q12395
C	69	SER	-	EXPRESSION TAG	UNP Q12395
E	68	GLY	-	EXPRESSION TAG	UNP Q12395
E	69	SER	-	EXPRESSION TAG	UNP Q12395
G	68	GLY	-	EXPRESSION TAG	UNP Q12395
G	69	SER	-	EXPRESSION TAG	UNP Q12395
I	68	GLY	-	EXPRESSION TAG	UNP Q12395
I	69	SER	-	EXPRESSION TAG	UNP Q12395

- Molecule 2 is a protein called Cell division control protein 53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	70	Total	C	N	O	S	0	0	0
			559	350	105	100	4			
2	D	68	Total	C	N	O	S	0	0	0
			543	341	99	99	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	68	Total	C	N	O	S	0	0	0
			540	339	97	100	4			
2	H	68	Total	C	N	O	S	0	0	0
			539	339	99	97	4			
2	J	61	Total	C	N	O	S	0	0	0
			474	298	84	88	4			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	740	GLY	-	EXPRESSION TAG	UNP Q12018
B	741	SER	-	EXPRESSION TAG	UNP Q12018
D	740	GLY	-	EXPRESSION TAG	UNP Q12018
D	741	SER	-	EXPRESSION TAG	UNP Q12018
F	740	GLY	-	EXPRESSION TAG	UNP Q12018
F	741	SER	-	EXPRESSION TAG	UNP Q12018
H	740	GLY	-	EXPRESSION TAG	UNP Q12018
H	741	SER	-	EXPRESSION TAG	UNP Q12018
J	740	GLY	-	EXPRESSION TAG	UNP Q12018
J	741	SER	-	EXPRESSION TAG	UNP Q12018

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Defective in cullin neddylation protein 1

Chain A:




- Molecule 1: Defective in cullin neddylation protein 1

Chain C:



- Molecule 1: Defective in cullin neddylation protein 1

Chain E:



- Molecule 1: Defective in cullin neddylation protein 1

Chain G:



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	123.91Å 123.91Å 192.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 3.10 39.69 – 3.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.10) 99.1 (39.69-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 3.12Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.258 , 0.308 0.264 , 0.308	Depositor DCC
R_{free} test set	1529 reflections (5.09%)	DCC
Wilson B-factor (Å ²)	95.1	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 67.3	EDS
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 30166 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10662	wwPDB-VP
Average B, all atoms (Å ²)	114.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.61	0/1710	0.80	2/2317 (0.1%)
1	C	0.56	0/1736	0.72	0/2354
1	E	0.73	5/1714 (0.3%)	0.81	2/2324 (0.1%)
1	G	0.51	0/1684	0.72	0/2283
1	I	0.46	0/1381	0.71	1/1860 (0.1%)
2	B	0.62	0/566	0.82	0/757
2	D	0.55	0/549	0.70	0/733
2	F	0.50	0/545	0.77	1/728 (0.1%)
2	H	0.56	0/546	0.84	0/731
2	J	0.45	0/478	0.73	1/640 (0.2%)
All	All	0.58	5/10909 (0.0%)	0.76	7/14727 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	170	ALA	C-O	11.32	1.44	1.23
1	E	170	ALA	CA-CB	-6.26	1.39	1.52
1	E	173	LEU	C-O	5.98	1.34	1.23
1	E	172	ASN	C-O	5.98	1.34	1.23
1	E	168	ASN	CG-ND2	5.03	1.45	1.32

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	ARG	NE-CZ-NH2	8.92	124.76	120.30
1	A	237	ARG	NH1-CZ-NH2	-6.14	112.64	119.40
2	F	800	GLY	N-CA-C	5.76	127.50	113.10
1	E	247	ASP	N-CA-C	5.71	126.42	111.00
1	E	253	ALA	N-CA-C	5.61	126.15	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1664	0	1613	88	3
1	C	1689	0	1635	60	0
1	E	1667	0	1613	150	3
1	G	1639	0	1584	57	0
1	I	1348	0	1311	89	0
2	B	559	0	572	59	0
2	D	543	0	554	57	0
2	F	540	0	554	80	0
2	H	539	0	551	39	0
2	J	474	0	473	29	0
All	All	10662	0	10460	680	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 32.

The worst 5 of 680 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:758:ILE:HD11	2:B:777:GLN:OE1	1.25	1.24
1:E:71:TYR:HB3	1:E:72:PRO:HD2	1.21	1.17
2:B:779:HIS:CG	2:B:780:GLN:H	1.60	1.15
2:D:762:LYS:HD2	2:D:765:LEU:HD11	1.24	1.15
1:E:261:PHE:O	1:E:265:LEU:HD11	1.47	1.15

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:237:ARG:NH2	1:E:71:TYR:N[2_655]	2.06	0.14
1:A:247:ASP:OD2	1:E:73:LYS:NZ[2_655]	2.07	0.13
1:A:237:ARG:NH2	1:E:71:TYR:CA[2_655]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/202 (96%)	171 (89%)	21 (11%)	1 (0%)	38	81
1	C	196/202 (97%)	177 (90%)	18 (9%)	1 (0%)	38	81
1	E	193/202 (96%)	167 (86%)	22 (11%)	4 (2%)	11	48
1	G	190/202 (94%)	172 (90%)	18 (10%)	0	100	100
1	I	145/202 (72%)	128 (88%)	15 (10%)	2 (1%)	16	58
2	B	68/76 (90%)	58 (85%)	6 (9%)	4 (6%)	2	17
2	D	64/76 (84%)	58 (91%)	5 (8%)	1 (2%)	14	55
2	F	64/76 (84%)	57 (89%)	6 (9%)	1 (2%)	14	55
2	H	66/76 (87%)	60 (91%)	6 (9%)	0	100	100
2	J	55/76 (72%)	48 (87%)	5 (9%)	2 (4%)	5	31
All	All	1234/1390 (89%)	1096 (89%)	122 (10%)	16 (1%)	18	60

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	764	ASN
2	B	779	HIS
2	B	782	PHE
2	D	764	ASN
1	E	72	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/190 (97%)	174 (95%)	10 (5%)	31	72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	187/190 (98%)	176 (94%)	11 (6%)	28	68
1	E	184/190 (97%)	167 (91%)	17 (9%)	13	45
1	G	181/190 (95%)	177 (98%)	4 (2%)	64	92
1	I	148/190 (78%)	131 (88%)	17 (12%)	8	31
2	B	59/65 (91%)	53 (90%)	6 (10%)	11	37
2	D	58/65 (89%)	51 (88%)	7 (12%)	7	27
2	F	58/65 (89%)	48 (83%)	10 (17%)	3	11
2	H	57/65 (88%)	54 (95%)	3 (5%)	32	72
2	J	50/65 (77%)	45 (90%)	5 (10%)	11	38
All	All	1166/1275 (92%)	1076 (92%)	90 (8%)	18	57

5 of 90 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	175	LEU
1	E	265	LEU
1	I	257	ILE
1	E	206	ASP
1	E	250	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	168	ASN
2	F	767	HIS
1	I	196	GLN
1	E	242	GLN
2	F	764	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/202 (96%)	-0.02	0 100 100	67, 93, 113, 129	0
1	C	198/202 (98%)	-0.00	0 100 100	66, 88, 118, 133	0
1	E	195/202 (96%)	-0.01	0 100 100	80, 118, 139, 144	0
1	G	192/202 (95%)	0.05	2 (1%) 79 23	79, 113, 137, 145	0
1	I	157/202 (77%)	0.43	8 (5%) 27 4	135, 153, 179, 191	0
2	B	70/76 (92%)	0.06	0 100 100	80, 103, 141, 149	0
2	D	68/76 (89%)	0.04	1 (1%) 70 16	80, 110, 135, 151	0
2	F	68/76 (89%)	0.24	1 (1%) 70 16	106, 129, 157, 163	0
2	H	68/76 (89%)	-0.00	0 100 100	89, 111, 129, 136	0
2	J	61/76 (80%)	0.30	3 (4%) 28 4	116, 135, 154, 156	0
All	All	1272/1390 (91%)	0.09	15 (1%) 75 20	66, 111, 159, 191	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	265	LEU	5.6
1	I	130	PHE	2.6
1	I	239	PRO	2.6
2	J	765	LEU	2.6
1	G	127	ARG	2.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.