



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 11:12 AM GMT

PDB ID : 1O7G
Title : NAPHTHALENE 1,2-DIOXYGENASE WITH NAPHTHALENE BOUND IN THE ACTIVE SITE.
Authors : Karlsson, A.; Parales, J.V.; Parales, R.E.; Gibson, D.T.; Eklund, H.; Ramaswamy, S.
Deposited on : 2002-11-05
Resolution : 1.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	449	
2	B	194	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	EDO	B	1697	-	X
4	NPY	A	1450	-	X
5	SO4	B	1699	-	X

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 5662 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAPHTHALENE 1,2-DIOXYGENASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	0	1
			3481	2202	598	665	16			

- Molecule 2 is a protein called NAPHTHALENE 1,2-DIOXYGENASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	193	Total	C	N	O	S	0	0	0
			1608	1007	302	293	6			

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is NAPHTHALENE (three-letter code: NPY) (formula: C₁₀H₈).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C	0	0
			10	10		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	Fe 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	368	Total 368	O 368	0	0
8	B	151	Total 151	O 151	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NAPHTHALENE 1,2-DIOXYGENASE ALPHA SUBUNIT



• Molecule 2: NAPHTHALENE 1,2-DIOXYGENASE BETA SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	139.67Å 139.67Å 208.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 1.70 19.80 – 1.70	Depositor EDS
% Data completeness (in resolution range)	95.9 (25.00-1.70) 95.9 (19.80-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.0.36	Depositor
R, R_{free}	0.163 , 0.187 0.175 , 0.198	Depositor DCC
R_{free} test set	4117 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 82025 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5662	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NPY, FE, FES, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	0/3573	0.98	14/4841 (0.3%)
2	B	0.80	0/1638	0.98	7/2209 (0.3%)
All	All	0.86	0/5211	0.98	21/7050 (0.3%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	671	ARG	NE-CZ-NH2	-8.57	116.01	120.30
1	A	161	ASP	CB-CG-OD2	7.99	125.49	118.30
1	A	205	ASP	CB-CG-OD2	7.60	125.14	118.30
1	A	293	ARG	NE-CZ-NH1	-7.41	116.59	120.30
1	A	3	TYR	CB-CG-CD2	7.19	125.32	121.00
2	B	671	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	A	45	ASP	CB-CG-OD2	6.51	124.16	118.30
1	A	286	ASP	CB-CG-OD2	6.44	124.09	118.30
1	A	293	ARG	NE-CZ-NH2	6.41	123.50	120.30
1	A	320	ASP	CB-CG-OD2	6.40	124.06	118.30
1	A	47	LEU	CB-CG-CD1	6.04	121.26	111.00
1	A	122	ASP	CB-CG-OD2	5.65	123.39	118.30
2	B	527	ASP	CB-CG-OD2	5.61	123.35	118.30
2	B	572	ARG	NE-CZ-NH1	5.57	123.08	120.30
1	A	218	ARG	NE-CZ-NH1	5.37	122.98	120.30
2	B	572	ARG	NE-CZ-NH2	-5.33	117.63	120.30
2	B	508	ASP	CB-CG-OD2	5.26	123.03	118.30
1	A	339	ASP	CB-CG-OD2	5.25	123.03	118.30
2	B	638	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	A	255	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	62	ASP	CB-CG-OD2	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3481	0	3316	28	0
2	B	1608	0	1583	23	0
3	A	8	0	12	0	0
3	B	16	0	24	6	0
4	A	10	0	8	0	0
5	A	5	0	0	1	0
5	B	10	0	0	0	0
6	A	4	0	0	1	0
7	A	1	0	0	0	0
8	A	368	0	0	1	0
8	B	151	0	0	4	0
All	All	5662	0	4943	51	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (51) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:577:ARG:H	2:B:577:ARG:HD2	1.06	1.15
2:B:533:GLU:OE2	2:B:671:ARG:HD2	1.45	1.14
2:B:660:GLU:HG3	3:B:1697:EDO:H12	1.29	1.12
2:B:578:ARG:HH11	2:B:578:ARG:HB2	1.15	1.08
2:B:577:ARG:N	2:B:577:ARG:HD2	1.70	0.95
1:A:198:PRO:HA	1:A:366:MET:HE1	1.49	0.94
1:A:198:PRO:HA	1:A:366:MET:CE	2.02	0.88
1:A:240:LEU:N	1:A:240:LEU:HD13	1.92	0.85
2:B:658:ALA:O	3:B:1697:EDO:H22	1.80	0.81
2:B:678:VAL:HA	3:B:1697:EDO:H21	1.62	0.80
2:B:578:ARG:CB	2:B:578:ARG:HH11	1.95	0.77
2:B:660:GLU:HG3	3:B:1697:EDO:C1	2.12	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:131:LYS:HG2	5:A:1451:SO4:O2	1.85	0.75
2:B:577:ARG:CD	2:B:577:ARG:N	2.47	0.74
2:B:555:GLU:HG3	8:B:2047:HOH:O	1.86	0.74
2:B:660:GLU:CG	3:B:1697:EDO:H12	2.15	0.73
2:B:578:ARG:NH1	2:B:578:ARG:HB2	2.00	0.67
1:A:240:LEU:N	1:A:240:LEU:CD1	2.56	0.67
2:B:523:PHE:HE1	8:B:2104:HOH:O	1.77	0.66
1:A:181:GLU:HG3	1:A:334:LYS:HG2	1.78	0.65
1:A:201:ASN:CB	1:A:366:MET:HE1	2.27	0.64
1:A:201:ASN:HB3	1:A:366:MET:HE1	1.82	0.61
2:B:561:GLU:H	2:B:561:GLU:CD	2.02	0.61
1:A:240:LEU:HD13	1:A:240:LEU:H	1.66	0.58
1:A:266:VAL:HB	1:A:267:PRO:HD3	1.88	0.55
2:B:575:SER:O	2:B:577:ARG:NE	2.37	0.55
2:B:556:HIS:CE1	8:B:2037:HOH:O	2.60	0.54
1:A:240:LEU:HD13	1:A:252:VAL:HB	1.88	0.53
2:B:537:LEU:HD13	8:B:2033:HOH:O	2.08	0.53
1:A:405:LYS:NZ	8:A:2329:HOH:O	2.42	0.53
1:A:224:PHE:CE1	1:A:253:LEU:HD23	2.45	0.52
1:A:201:ASN:CB	1:A:366:MET:CE	2.90	0.48
1:A:198:PRO:HA	1:A:366:MET:HE3	1.88	0.47
1:A:201:ASN:HB2	1:A:366:MET:HE1	1.96	0.47
1:A:129:ASN:OD1	1:A:132:CYS:SG	2.65	0.47
1:A:47:LEU:HD22	1:A:182:LEU:HD23	1.99	0.45
1:A:240:LEU:CD1	1:A:252:VAL:HB	2.47	0.44
2:B:504:ASN:OD1	2:B:506:GLN:HB2	2.17	0.44
1:A:104:HIS:HB2	6:A:1452:FES:S1	2.59	0.43
1:A:239:GLY:HA3	1:A:252:VAL:O	2.20	0.42
1:A:308:THR:O	1:A:309:CYS:HB2	2.19	0.42
1:A:220:GLY:O	1:A:221:GLU:HB2	2.19	0.42
2:B:531:GLN:NE2	2:B:532:GLN:OE1	2.53	0.41
2:B:578:ARG:HH11	2:B:578:ARG:CG	2.33	0.41
1:A:198:PRO:HB2	1:A:316:TRP:CE2	2.56	0.41
1:A:261:HIS:NE2	1:A:344:LEU:HD11	2.36	0.41
1:A:203:VAL:HG23	1:A:299:THR:HB	2.03	0.41
1:A:311:GLY:HA2	1:A:331:ILE:HG13	2.02	0.41
2:B:561:GLU:CD	2:B:561:GLU:N	2.70	0.40
1:A:14:LEU:HD23	1:A:14:LEU:HA	1.93	0.40
2:B:658:ALA:O	3:B:1697:EDO:C2	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/449 (99%)	433 (97%)	13 (3%)	0	100	100
2	B	191/194 (98%)	185 (97%)	5 (3%)	1 (0%)	38	17
All	All	637/643 (99%)	618 (97%)	18 (3%)	1 (0%)	56	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	524	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/369 (100%)	359 (98%)	8 (2%)	64	43
2	B	172/173 (99%)	158 (92%)	14 (8%)	17	4
All	All	539/542 (99%)	517 (96%)	22 (4%)	41	17

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LEU
1	A	103	TYR
1	A	131	LYS
1	A	221	GLU
1	A	240	LEU
1	A	278	GLU
1	A	293	ARG
1	A	331	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	508	ASP
2	B	519	ILE
2	B	531	GLN
2	B	532	GLN
2	B	537	LEU
2	B	543	HIS
2	B	571	LEU
2	B	577	ARG
2	B	578	ARG
2	B	593	GLN
2	B	603	LEU
2	B	667	GLU
2	B	671	ARG
2	B	690	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	115	GLN
2	B	514	HIS
2	B	531	GLN
2	B	556	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1448	-	3,3,3	0.40	0	2,2,2	0.08	0
3	EDO	A	1449	-	3,3,3	0.38	0	2,2,2	0.71	0
4	NPY	A	1450	-	11,11,11	1.31	1 (9%)	14,14,14	0.73	0
5	SO4	A	1451	-	4,4,4	0.42	0	6,6,6	1.35	2 (33%)
6	FES	A	1452	1	0,4,4	0.00	-	0,4,4	0.00	-
3	EDO	B	1695	-	3,3,3	0.54	0	2,2,2	0.06	0
3	EDO	B	1696	-	3,3,3	0.60	0	2,2,2	0.79	0
3	EDO	B	1697	-	3,3,3	0.57	0	2,2,2	0.22	0
3	EDO	B	1698	-	3,3,3	0.34	0	2,2,2	0.65	0
5	SO4	B	1699	-	4,4,4	0.15	0	6,6,6	0.20	0
5	SO4	B	1700	-	4,4,4	0.46	0	6,6,6	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1448	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1449	-	-	0/1/1/1	0/0/0/0
4	NPY	A	1450	-	-	0/0/0/0	0/0/2/2
5	SO4	A	1451	-	-	0/0/0/0	0/0/0/0
6	FES	A	1452	1	-	0/0/4/4	0/0/1/1
3	EDO	B	1695	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1696	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1697	-	-	0/1/1/1	0/0/0/0
3	EDO	B	1698	-	-	0/1/1/1	0/0/0/0
5	SO4	B	1699	-	-	0/0/0/0	0/0/0/0
5	SO4	B	1700	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1450	NPY	C8A-C4A	2.58	1.48	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1451	SO4	O4-S-O3	2.49	119.59	109.08
5	A	1451	SO4	O2-S-O1	-2.04	102.78	109.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	448/449 (99%)	-0.01	17 (3%)	38 43	13, 20, 36, 57	0
2	B	193/194 (99%)	0.15	12 (6%)	20 24	14, 21, 42, 54	0
All	All	641/643 (99%)	0.04	29 (4%)	31 36	13, 20, 39, 57	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	447	THR	12.0
1	A	448	ASP	8.3
2	B	502	MET	7.8
2	B	577	ARG	6.8
2	B	525	CYS	6.1
1	A	236	GLU	5.4
2	B	667	GLU	4.5
1	A	446	THR	4.2
1	A	235	PRO	4.1
2	B	523	PHE	4.0
1	A	223	ILE	3.9
2	B	629	VAL	3.9
2	B	630	ASN	3.6
1	A	221	GLU	3.6
1	A	283	GLU	3.5
1	A	232	ALA	3.5
1	A	224	PHE	3.3
1	A	282	LYS	3.2
2	B	632	LYS	2.7
1	A	234	PRO	2.5
2	B	580	LYS	2.4
1	A	226	SER	2.4
1	A	225	SER	2.3
1	A	129	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	532	GLN	2.1
2	B	524	ASN	2.1
1	A	264	ASP	2.0
2	B	520	LEU	2.0
1	A	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	B	1697	4/4	0.36	13.08	54,59,60,62	0
5	SO4	B	1699	5/5	0.33	2.79	51,51,54,55	0
4	NPY	A	1450	10/10	0.19	2.66	37,37,38,39	0
5	SO4	A	1451	5/5	0.27	1.52	41,44,44,44	0
3	EDO	B	1698	4/4	0.19	0.67	53,54,54,55	0
3	EDO	A	1448	4/4	0.07	0.52	17,18,20,20	0
3	EDO	A	1449	4/4	0.08	0.22	18,24,25,26	0
3	EDO	B	1695	4/4	0.08	-0.05	25,27,27,29	0
3	EDO	B	1696	4/4	0.10	-0.18	40,41,43,44	0
5	SO4	B	1700	5/5	0.14	-0.19	44,44,46,46	0
6	FES	A	1452	4/4	0.06	-2.53	14,15,17,17	0
7	FE	A	1453	1/1	0.02	-8.37	22,22,22,22	0

6.5 Other polymers ⓘ

There are no such residues in this entry.