



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 05:54 AM GMT

PDB ID : 2O8W
Title : Crystal Structure and Binding Epitopes of Urokinase-type Plasminogen Activator (C122A/N145Q/S195A) in complex with Inhibitors
Authors : Zhao, G.; Yuan, C.; Jiang, L.; Huang, Z.; Huang, M.
Deposited on : 2006-12-12
Resolution : 1.86 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance


The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1269 (1.86-1.86)
Clashscore	79885	1470 (1.86-1.86)
Ramachandran outliers	78287	1451 (1.86-1.86)
Sidechain outliers	78261	1451 (1.86-1.86)
RSRZ outliers	66119	1269 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	253	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 2252 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1932	1220	338	359	15	0	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	122	ALA	CYS	ENGINEERED	UNP Q5PY49
A	145	GLN	ASN	ENGINEERED	UNP Q5PY49
A	195	ALA	SER	ENGINEERED	UNP Q5PY49

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is 1-PHENYLGUANIDINE (three-letter code: PL0) (formula: $C_7H_9N_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			10	7	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	292	Total 292	O 292	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	120.88Å 120.88Å 42.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	60.24 – 1.86 60.44 – 1.86	Depositor EDS
% Data completeness (in resolution range)	79.5 (60.24-1.86) 79.6 (60.44-1.86)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.86Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.255 0.212 , 0.254	Depositor DCC
R_{free} test set	797 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.4	EDS
Estimated twinning fraction	0.021 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 15503 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2252	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, SO4, PL0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1981	0.63	1/2685 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	199	LEU	N-CA-C	-5.49	96.19	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1932	0	1882	69	0
2	A	5	0	0	0	0
3	A	13	0	18	2	0
4	A	10	0	8	0	0
5	A	292	0	0	18	0
All	All	2252	0	1908	69	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 18.

All (69) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:136:CYS:SG	1:A:162:LEU:HD11	2.06	0.96
1:A:24:ILE:HG12	1:A:69:GLY:HA2	1.65	0.78
1:A:131:GLN:HG2	5:A:5154:HOH:O	1.84	0.77
1:A:128:ASN:HD21	1:A:203:LEU:HD21	1.53	0.74
1:A:91:HIS:HD2	1:A:93:ASP:H	1.35	0.73
1:A:81:MET:HE1	1:A:113:GLN:H	1.54	0.71
1:A:209:LEU:CD1	1:A:231:VAL:HG21	2.23	0.69
1:A:235:LEU:O	1:A:239:ARG:HG3	1.93	0.68
1:A:39:THR:HG23	5:A:5404:HOH:O	1.93	0.67
1:A:115:SER:HB3	5:A:5360:HOH:O	1.96	0.65
1:A:84:GLU:HG3	5:A:5396:HOH:O	1.96	0.65
1:A:81:MET:CE	1:A:113:GLN:H	2.13	0.62
1:A:239:ARG:HD2	5:A:5187:HOH:O	2.00	0.61
1:A:156:LYS:NZ	5:A:5205:HOH:O	2.32	0.61
1:A:37(A):ARG:HD2	5:A:5353:HOH:O	2.00	0.61
1:A:170:GLN:HG3	5:A:5146:HOH:O	2.00	0.60
1:A:235:LEU:HD23	5:A:5201:HOH:O	2.02	0.59
1:A:53:ILE:HD11	1:A:103:ILE:HD11	1.83	0.58
1:A:162:LEU:HD12	1:A:162:LEU:N	2.18	0.58
1:A:91:HIS:CD2	1:A:93:ASP:H	2.21	0.57
1:A:170:GLN:HB3	1:A:170(A):PRO:HD2	1.85	0.57
1:A:16:ILE:O	5:A:5205:HOH:O	2.17	0.57
1:A:45:SER:OG	1:A:198:PRO:HB3	2.05	0.56
1:A:97(A):THR:O	1:A:97(B):LEU:HD23	2.05	0.56
1:A:110(D):ARG:HG2	1:A:110(D):ARG:HH11	1.70	0.56
1:A:185(B):GLN:HB2	1:A:187:LYS:HD3	1.87	0.56
1:A:209:LEU:HD12	1:A:231:VAL:HG21	1.88	0.56
1:A:166:ARG:O	1:A:170:GLN:HG2	2.05	0.56
1:A:97(A):THR:HG22	1:A:97(B):LEU:HG	1.89	0.54
1:A:95:SER:OG	1:A:100:HIS:HB2	2.08	0.54
1:A:202:SER:HB2	3:A:5115:PG4:H51	1.90	0.54
1:A:187:LYS:NZ	1:A:187:LYS:HB2	2.22	0.54
1:A:72:ARG:NE	5:A:5158:HOH:O	2.41	0.53
1:A:77:THR:O	1:A:80:GLU:HG2	2.09	0.51
1:A:161:LYS:HD3	5:A:5393:HOH:O	2.11	0.51
1:A:109:ARG:HG3	1:A:109:ARG:O	2.11	0.51
1:A:202:SER:CB	3:A:5115:PG4:H51	2.42	0.50
1:A:81:MET:CE	1:A:118:ILE:HD13	2.42	0.49
1:A:27:GLN:HG2	1:A:157:MET:HE1	1.94	0.49
1:A:233:HIS:HE1	5:A:5327:HOH:O	1.97	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:59:PHE:HD1	1:A:64:TYR:CE2	2.32	0.48
1:A:81:MET:HE1	1:A:112:ALA:HA	1.96	0.47
1:A:22:THR:HG21	1:A:157:MET:HE3	1.97	0.47
1:A:86:GLU:O	1:A:87:ASN:ND2	2.48	0.46
1:A:170(A):PRO:HD3	5:A:5140:HOH:O	2.14	0.46
1:A:25:GLU:H	1:A:25:GLU:CD	2.18	0.46
1:A:170(A):PRO:HG2	1:A:170(B):HIS:H	1.81	0.46
1:A:27:GLN:HG3	1:A:139:THR:OG1	2.17	0.45
1:A:98:ALA:HA	1:A:215:TRP:CE2	2.50	0.45
1:A:139:THR:HG22	1:A:155:LEU:HD11	1.98	0.45
1:A:238:ILE:HD12	5:A:5201:HOH:O	2.16	0.44
1:A:197:GLY:HA3	5:A:5116:HOH:O	2.17	0.44
1:A:217:ARG:HG2	1:A:217:ARG:HH11	1.83	0.44
1:A:176:VAL:HA	1:A:180:MET:SD	2.57	0.43
1:A:145:GLN:HB2	1:A:148:ASP:CG	2.38	0.43
1:A:27:GLN:NE2	1:A:137:GLU:OE2	2.52	0.43
1:A:70:ARG:HD2	1:A:80:GLU:OE1	2.19	0.43
1:A:81:MET:HE3	1:A:118:ILE:HD13	2.00	0.42
1:A:86:GLU:HB2	1:A:109:ARG:HG2	2.01	0.42
1:A:188:THR:O	1:A:189:ASP:HB2	2.19	0.42
1:A:223(A):ASP:O	1:A:224:LYS:HD3	2.19	0.42
1:A:26:ASN:O	1:A:27:GLN:HG2	2.20	0.42
1:A:131:GLN:HE21	1:A:131:GLN:HA	1.85	0.42
1:A:181:LEU:HG	1:A:230:ARG:HG3	2.01	0.41
1:A:203:LEU:HD13	5:A:5283:HOH:O	2.19	0.41
1:A:206:ARG:HG3	1:A:206:ARG:HH11	1.85	0.41
1:A:61:LYS:HE2	1:A:62(A):GLU:HB2	2.02	0.41
1:A:206:ARG:NH1	1:A:206:ARG:HG3	2.36	0.40
1:A:170(A):PRO:HB3	5:A:5140:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/253 (96%)	228 (94%)	15 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/218 (97%)	209 (99%)	3 (1%)	78	68

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	LEU
1	A	97	ASP
1	A	168	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	37	HIS
1	A	91	HIS
1	A	128	ASN
1	A	131	GLN
1	A	165	HIS
1	A	204	GLN
1	A	233	HIS
1	A	241	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	404	-	4,4,4	0.26	0	6,6,6	0.11	0
3	PG4	A	5115	-	12,12,12	0.55	0	11,11,11	3.59	1 (9%)
4	PL0	A	800	-	10,10,10	2.41	5 (50%)	12,12,12	4.10	5 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	404	-	-	0/0/0/0	0/0/0/0
3	PG4	A	5115	-	-	0/10/10/10	0/0/0/0
4	PL0	A	800	-	-	0/4/4/4	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800	PL0	C5-N1	5.13	1.51	1.41
4	A	800	PL0	C6-C5	2.51	1.43	1.39
4	A	800	PL0	C4-C5	2.37	1.43	1.39
4	A	800	PL0	C3-C4	2.37	1.44	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800	PL0	C3-C2	2.17	1.44	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	5115	PG4	O2-C2-C1	11.52	165.99	110.61
4	A	800	PL0	N1-C7-N3	-8.54	99.56	122.86
4	A	800	PL0	N2-C7-N1	7.71	146.13	115.42
4	A	800	PL0	C5-N1-C7	-6.49	113.30	128.35
4	A	800	PL0	C4-C5-N1	-3.55	108.46	120.39
4	A	800	PL0	C6-C5-N1	3.18	131.08	120.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	245/253 (96%)	0.08	12 (4%)	28 27	13, 20, 35, 46	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37(A)	ARG	4.0
1	A	97(A)	THR	3.9
1	A	97(B)	LEU	3.6
1	A	37(C)	GLY	3.5
1	A	132	PHE	2.7
1	A	170(A)	PRO	2.7
1	A	170(B)	HIS	2.4
1	A	203	LEU	2.3
1	A	109	ARG	2.3
1	A	37(D)	SER	2.1
1	A	204	GLN	2.0
1	A	97	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	PL0	A	800	10/10	0.12	-	18,19,20,20	0
2	SO4	A	404	5/5	0.09	-	37,37,38,38	0
3	PG4	A	5115	13/13	0.21	-	57,57,58,58	0

6.5 Other polymers ⓘ

There are no such residues in this entry.