



wwPDB X-ray Structure Validation Summary Report

Mar 31, 2014 – 01:45 PM BST

PDB ID : 4OCL
Title : Crystal Structure of the Rpn8-Rpn11 MPN domain heterodimer, crystal form Ia
Authors : Pathare, G.R.; Bracher, A.
Deposited on : 2014-01-09
Resolution : 2.40 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable23004
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable23004

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	187	
1	D	187	
2	B	220	
2	E	220	
3	C	133	
3	F	133	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7359 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 26S proteasome regulatory subunit RPN8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	163	Total	C	N	O	S	0	0	0
			1252	800	211	236	5			
1	D	164	Total	C	N	O	S	0	0	0
			1265	808	213	239	5			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP Q08723
A	0	HIS	-	CLONING ARTIFACT	UNP Q08723
A	177	GLY	-	SEE REMARK 999	UNP Q08723
A	178	SER	-	SEE REMARK 999	UNP Q08723
A	179	GLY	-	SEE REMARK 999	UNP Q08723
A	180	GLY	-	SEE REMARK 999	UNP Q08723
A	181	SER	-	SEE REMARK 999	UNP Q08723
A	182	GLY	-	SEE REMARK 999	UNP Q08723
A	183	GLY	-	SEE REMARK 999	UNP Q08723
A	184	SER	-	SEE REMARK 999	UNP Q08723
A	185	GLY	-	SEE REMARK 999	UNP Q08723
D	-1	GLY	-	CLONING ARTIFACT	UNP Q08723
D	0	HIS	-	CLONING ARTIFACT	UNP Q08723
D	177	GLY	-	SEE REMARK 999	UNP Q08723
D	178	SER	-	SEE REMARK 999	UNP Q08723
D	179	GLY	-	SEE REMARK 999	UNP Q08723
D	180	GLY	-	SEE REMARK 999	UNP Q08723
D	181	SER	-	SEE REMARK 999	UNP Q08723
D	182	GLY	-	SEE REMARK 999	UNP Q08723
D	183	GLY	-	SEE REMARK 999	UNP Q08723
D	184	SER	-	SEE REMARK 999	UNP Q08723
D	185	GLY	-	SEE REMARK 999	UNP Q08723

- Molecule 2 is a protein called 26S proteasome regulatory subunit RPN11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	0	0
			1396	890	240	254	12			
2	E	184	Total	C	N	O	S	0	0	0
			1402	893	240	257	12			

- Molecule 3 is a protein called Nb1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	125	Total	C	N	O	S	0	0	0
			973	611	173	185	4			
3	F	125	Total	C	N	O	S	0	0	0
			975	612	176	183	4			


- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	E	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total	O	0	0
			11	11		
5	B	21	Total	O	0	0
			21	21		
5	C	14	Total	O	0	0
			14	14		
5	D	17	Total	O	0	0
			17	17		
5	E	11	Total	O	0	0
			11	11		
5	F	20	Total	O	0	0
			20	20		

● Molecule 3: Nb1



● Molecule 3: Nb1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	45.26Å 63.43Å 100.11Å 100.08° 92.75° 90.62°	Depositor
Resolution (Å)	30.00 – 2.40 28.75 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.2 (30.00-2.40) 97.2 (28.75-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.195 , 0.252 0.191 , 0.245	Depositor DCC
R_{free} test set	2088 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	45.6	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.5	EDS
Estimated twinning fraction	0.067 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 41608 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7359	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/1275	0.79	0/1734
1	D	0.88	1/1288 (0.1%)	0.86	2/1750 (0.1%)
2	B	0.84	1/1421 (0.1%)	0.85	2/1919 (0.1%)
2	E	0.86	1/1427 (0.1%)	0.83	1/1933 (0.1%)
3	C	0.80	0/999	0.86	1/1354 (0.1%)
3	F	0.83	0/1001	0.88	1/1356 (0.1%)
All	All	0.83	3/7411 (0.0%)	0.84	7/10046 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	F	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	178	GLY	C-O	7.70	1.35	1.23
1	D	108	GLU	CG-CD	6.62	1.61	1.51
2	E	209	GLU	CG-CD	5.33	1.59	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	20	ARG	NE-CZ-NH1	-7.81	116.39	120.30
1	D	89	LEU	CA-CB-CG	7.81	133.26	115.30
2	B	153	ILE	CG1-CB-CG2	-5.80	98.63	111.40
2	B	100	ARG	NE-CZ-NH2	-5.39	117.60	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	10	ILE	CG1-CB-CG2	-5.38	99.57	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	F	104	GLY	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1252	0	0	2	0
1	D	1265	0	0	9	0
2	B	1396	0	0	13	0
2	E	1402	0	0	7	0
3	C	973	0	0	2	0
3	F	975	0	0	5	0
4	B	1	0	0	0	0
4	E	1	0	0	0	0
5	A	11	0	0	0	0
5	B	21	0	0	2	0
5	C	14	0	0	0	0
5	D	17	0	0	0	0
5	E	11	0	0	0	0
5	F	20	0	0	0	0
All	All	7359	0	0	36	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:59:ASP:O	2:B:135:ARG:NH1	2.08	0.86
2:B:151:VAL:CG1	2:B:153:ILE:CD1	2.59	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:74:SER:O	2:E:76:THR:N	2.19	0.75
2:B:111:HIS:CD2	2:B:115:GLY:O	2.44	0.70
3:F:4:GLN:NE2	3:F:112:PHE:CE1	2.62	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/187 (85%)	154 (97%)	3 (2%)	2 (1%)	18	24
1	D	160/187 (86%)	153 (96%)	7 (4%)	0	100	100
2	B	177/220 (80%)	170 (96%)	6 (3%)	1 (1%)	33	47
2	E	180/220 (82%)	172 (96%)	7 (4%)	1 (1%)	33	47
3	C	123/133 (92%)	120 (98%)	3 (2%)	0	100	100
3	F	123/133 (92%)	118 (96%)	4 (3%)	1 (1%)	27	39
All	All	922/1080 (85%)	887 (96%)	30 (3%)	5 (0%)	38	53

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	59	ASP
2	E	75	GLY
1	A	141	GLU
1	A	30	ASN
3	F	27	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/162 (82%)	122 (92%)	10 (8%)	19	28
1	D	134/162 (83%)	128 (96%)	6 (4%)	38	57
2	B	152/191 (80%)	142 (93%)	10 (7%)	24	35
2	E	150/191 (78%)	143 (95%)	7 (5%)	36	54
3	C	101/109 (93%)	94 (93%)	7 (7%)	22	33
3	F	101/109 (93%)	90 (89%)	11 (11%)	9	13
All	All	770/924 (83%)	719 (93%)	51 (7%)	24	35

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	28	ARG
1	D	9	THR
3	F	75	ASN
3	C	68	ARG
1	D	43	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	163/187 (87%)	0.19	8 (4%) 28 26	27, 52, 77, 92	0
1	D	164/187 (87%)	0.05	2 (1%) 75 75	24, 46, 66, 82	0
2	B	181/220 (82%)	0.10	5 (2%) 50 48	24, 44, 75, 93	0
2	E	184/220 (83%)	0.28	15 (8%) 12 10	22, 42, 92, 144	0
3	C	125/133 (93%)	0.25	6 (4%) 29 27	24, 45, 74, 98	0
3	F	125/133 (93%)	-0.02	5 (4%) 36 34	21, 41, 71, 88	0
All	All	942/1080 (87%)	0.15	41 (4%) 34 30	21, 45, 76, 144	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	180	LEU	8.4
2	E	185	ILE	7.6
2	B	163	ALA	6.9
3	C	30	PHE	6.2
2	E	181	ASN	6.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ZN	E	401	1/1	0.03	-2.97	64,64,64,64	0
4	ZN	B	401	1/1	0.05	-3.86	74,74,74,74	0

6.5 Other polymers

There are no such residues in this entry.