



# wwPDB X-ray Structure Validation Summary Report

Nov 3, 2014 – 11:48 PM EST

PDB ID : 4OG9  
Title : Cytokine complex crystal form 2  
Authors : Desmet, J.; Verstraete, K.; Bloch, Y.; Lorent, E.; Wen, Y.; Devreese, B.; Vandenbroucke, K.; Loverix, S.; Hettmann, T.; Deroo, S.; Somers, K.; Hendrikx, P.; Lasters, I.; Savvides, S.N.  
Deposited on : 2014-01-15  
Resolution : 3.39 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable24103  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.1.3  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable24103

# 1 Overall quality at a glance

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1017 (3.52-3.28)
Clashscore	79885	1214 (3.50-3.30)
Ramachandran outliers	78287	1177 (3.50-3.30)
Sidechain outliers	78261	1177 (3.50-3.30)
RSRZ outliers	66119	1017 (3.52-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	328	
2	B	198	
3	C	118	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8023 atoms, of which 3902 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	289	Total	C	H	N	O	S	0	0	0
			4233	1404	2023	351	443	12			

- Molecule 2 is a protein called Interleukin-23 subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	162	Total	C	H	N	O	S	0	0	0
			2402	779	1173	220	224	6			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	GLY	-	EXPRESSION TAG	UNP Q9NPF7
B	191	THR	-	EXPRESSION TAG	UNP Q9NPF7
B	192	LYS	-	EXPRESSION TAG	UNP Q9NPF7
B	193	HIS	-	EXPRESSION TAG	UNP Q9NPF7
B	194	HIS	-	EXPRESSION TAG	UNP Q9NPF7
B	195	HIS	-	EXPRESSION TAG	UNP Q9NPF7
B	196	HIS	-	EXPRESSION TAG	UNP Q9NPF7
B	197	HIS	-	EXPRESSION TAG	UNP Q9NPF7
B	198	HIS	-	EXPRESSION TAG	UNP Q9NPF7

- Molecule 3 is a protein called alphabody MB23.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	81	Total	C	H	N	O	S	0	0	0
			1275	402	654	100	113	6			

- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	5	113	34	52	2	25	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.85Å 57.85Å 366.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.14 – 3.39 57.14 – 3.39	Depositor EDS
% Data completeness (in resolution range)	99.8 (57.14-3.39) 99.9 (57.14-3.39)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.98 (at 3.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1750)	Depositor
R, $R_{free}$	0.262 , 0.293 0.261 , 0.290	Depositor DCC
$R_{free}$ test set	942 reflections (9.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	109.9	Xtriage
Anisotropy	0.453	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 76.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 9460 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8023	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	127.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/2266	0.51	0/3097
2	B	0.30	0/1262	0.58	0/1722
3	C	0.29	0/622	0.60	0/832
All	All	0.28	0/4150	0.55	0/5651

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2210	2023	1	8	0
2	B	1229	1173	0	9	0
3	C	621	654	0	5	0
4	A	61	52	0	3	0
All	All	4121	3902	1	24	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:61:GLN:NE2	3:C:98:GLN:OE1	2.20	0.74
2:B:58:GLU:N	2:B:62:GLU:OE2	2.26	0.68
2:B:60:GLY:O	2:B:62:GLU:N	2.28	0.66
3:C:91:SER:N	3:C:94:GLU:OE1	2.32	0.63
2:B:135:LEU:O	2:B:137:PRO:HD3	2.03	0.58

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/328 (86%)	271 (96%)	8 (3%)	2 (1%)	30	83
2	B	160/198 (81%)	152 (95%)	5 (3%)	3 (2%)	12	64
3	C	75/118 (64%)	75 (100%)	0	0	100	100
All	All	516/644 (80%)	498 (96%)	13 (2%)	5 (1%)	22	78

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	305	SER
2	B	137	PRO
1	A	304	ALA
2	B	61	ASP
2	B	136	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/297 (80%)	239 (100%)	0	100	100
2	B	129/167 (77%)	128 (99%)	1 (1%)	89	97
3	C	62/81 (76%)	62 (100%)	0	100	100
All	All	430/545 (79%)	429 (100%)	1 (0%)	96	99

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	143	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	222	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	500	1,4	12,14,15	0.28	0	15,19,21	0.82	1 (6%)
4	NAG	A	501	4	12,14,15	0.40	0	15,19,21	0.27	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BMA	A	502	4	10,11,12	0.85	0	11,15,17	1.12	1 (9%)
4	MAN	A	503	4	10,11,12	0.60	0	11,15,17	1.00	1 (9%)
4	MAN	A	504	4	10,11,12	0.44	0	11,15,17	0.86	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	500	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	501	4	-	0/6/23/26	0/1/1/1
4	BMA	A	502	4	-	0/2/19/22	0/1/1/1
4	MAN	A	503	4	-	0/2/19/22	0/1/1/1
4	MAN	A	504	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	BMA	O3-C3-C2	2.97	115.14	109.74
4	A	500	NAG	C2-N2-C7	2.87	126.87	123.39
4	A	504	MAN	O2-C2-C3	-2.25	105.33	110.10
4	A	503	MAN	O2-C2-C3	-2.13	105.58	110.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	289/328 (88%)	0.59	21 (7%) 15 6	78, 124, 166, 173	0
2	B	162/198 (81%)	0.27	0 100 100	73, 104, 132, 152	0
3	C	81/118 (68%)	0.16	0 100 100	93, 131, 145, 156	0
All	All	532/644 (82%)	0.43	21 (3%) 37 14	73, 113, 159, 173	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	46	VAL	4.3
1	A	23	ILE	4.2
1	A	45	MET	4.2
1	A	104	LEU	4.0
1	A	44	GLU	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	500	14/15	0.29	-1.29	99,105,125,125	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	501	14/15	0.27	-1.71	101,104,125,127	0
4	BMA	A	502	11/12	0.24	-2.38	103,106,127,127	0
4	MAN	A	503	11/12	0.32	-	102,107,128,130	0
4	MAN	A	504	11/12	0.44	-	103,107,127,128	0

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.