



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 11:18 PM GMT

PDB ID : 2OIN  
Title : crystal structure of HCV NS3-4A R155K muntant  
Authors : Wei, Y.  
Deposited on : 2007-01-11  
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	200	
1	B	200	
2	C	21	
2	D	21	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	ZN	A	901	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2891 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1110	696	197	209	8			
1	B	181	Total	C	N	O	S	0	0	0
			1323	826	239	249	9			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	MET	-	CLONING ARTIFACT	UNP Q9ELS8
A	17	ALA	-	CLONING ARTIFACT	UNP Q9ELS8
A	18	SER	-	CLONING ARTIFACT	UNP Q9ELS8
A	19	MET	-	CLONING ARTIFACT	UNP Q9ELS8
A	20	THR	-	CLONING ARTIFACT	UNP Q9ELS8
A	21	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
A	22	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
A	23	GLN	-	CLONING ARTIFACT	UNP Q9ELS8
A	24	GLN	-	CLONING ARTIFACT	UNP Q9ELS8
A	25	MET	-	CLONING ARTIFACT	UNP Q9ELS8
A	26	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
A	145	ARG	GLN	CONFLICT	UNP Q9ELS8
A	181	LYS	ARG	ENGINEERED	UNP Q9ELS8
A	208	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
A	209	SER	-	CLONING ARTIFACT	UNP Q9ELS8
A	210	HIS	-	EXPRESSION TAG	UNP Q9ELS8
A	211	HIS	-	EXPRESSION TAG	UNP Q9ELS8
A	212	HIS	-	EXPRESSION TAG	UNP Q9ELS8
A	213	HIS	-	EXPRESSION TAG	UNP Q9ELS8
A	214	HIS	-	EXPRESSION TAG	UNP Q9ELS8
A	215	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1016	MET	-	CLONING ARTIFACT	UNP Q9ELS8
B	1017	ALA	-	CLONING ARTIFACT	UNP Q9ELS8
B	1018	SER	-	CLONING ARTIFACT	UNP Q9ELS8
B	1019	MET	-	CLONING ARTIFACT	UNP Q9ELS8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	1020	THR	-	CLONING ARTIFACT	UNP Q9ELS8
B	1021	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
B	1022	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
B	1023	GLN	-	CLONING ARTIFACT	UNP Q9ELS8
B	1024	GLN	-	CLONING ARTIFACT	UNP Q9ELS8
B	1025	MET	-	CLONING ARTIFACT	UNP Q9ELS8
B	1026	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
B	1145	ARG	GLN	CONFLICT	UNP Q9ELS8
B	1181	LYS	ARG	ENGINEERED	UNP Q9ELS8
B	1208	GLY	-	CLONING ARTIFACT	UNP Q9ELS8
B	1209	SER	-	CLONING ARTIFACT	UNP Q9ELS8
B	1210	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1211	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1212	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1213	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1214	HIS	-	EXPRESSION TAG	UNP Q9ELS8
B	1215	HIS	-	EXPRESSION TAG	UNP Q9ELS8

- Molecule 2 is a protein called NS4A peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	16	Total	C	N	O	0	0	0
			108	70	20	18			
2	D	21	Total	C	N	O	0	0	0
			142	93	25	24			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total	O	0	0
			105	105		
4	B	83	Total	O	0	0
			83	83		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	11	Total 11	O 11	0	0
4	D	7	Total 7	O 7	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyprotein

Chain A: 




- Molecule 1: Polyprotein

Chain B: 



- Molecule 2: NS4A peptide

Chain C: 



- Molecule 2: NS4A peptide

Chain D: 

There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	225.31Å 225.31Å 75.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.93 – 2.50 24.58 – 2.50	Depositor EDS
% Data completeness (in resolution range)	86.2 (19.93-2.50) 87.1 (24.58-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 2.50Å)	Xtriage
Refinement program	CNX 2005	Depositor
R, $R_{free}$	0.200 , 0.246 0.196 , 0.240	Depositor DCC
$R_{free}$ test set	1132 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 30.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 22091 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2891	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/1134	0.51	0/1550
1	B	0.29	0/1349	0.51	0/1843
2	C	0.43	0/108	0.54	0/145
2	D	0.40	0/143	0.57	0/193
All	All	0.31	0/2734	0.51	0/3731

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1110	0	1108	28	0
1	B	1323	0	1322	26	0
2	C	108	0	125	5	0
2	D	142	0	161	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	105	0	0	7	0
4	B	83	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	11	0	0	0	0
4	D	7	0	0	0	0
All	All	2891	0	2716	55	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (55) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:1156:ARG:HH11	1:B:1156:ARG:HG3	1.48	0.78
1:B:1158:ILE:HD12	1:B:1190:ALA:HB2	1.66	0.78
1:A:169:LEU:HG	4:A:939:HOH:O	1.83	0.77
1:A:64:THR:HG22	1:A:65:ALA:N	2.04	0.72
1:A:158:ILE:O	1:A:158:ILE:HD13	1.91	0.71
1:A:193:VAL:HG21	4:A:946:HOH:O	1.92	0.70
1:B:1186:THR:O	1:B:1187:ARG:HB2	1.92	0.67
1:A:169:LEU:HD21	4:A:946:HOH:O	1.94	0.66
1:A:64:THR:CG2	1:A:65:ALA:N	2.60	0.65
1:B:1122:PRO:HB3	1:B:1198:VAL:HG11	1.80	0.63
1:B:1156:ARG:HH11	1:B:1156:ARG:CG	2.13	0.61
1:B:1108:LEU:HD11	1:B:1201:LEU:HD12	1.82	0.60
1:A:64:THR:HG22	1:A:66:THR:H	1.66	0.60
1:A:88:ARG:HG2	2:C:21:GLY:O	2.02	0.59
2:C:30:VAL:O	2:C:30:VAL:HG13	2.02	0.59
1:B:1177:VAL:O	1:B:1198:VAL:HG23	2.04	0.58
1:A:132:LEU:HD23	1:A:133:VAL:N	2.18	0.58
1:B:1103:ASN:ND2	1:B:1106:GLN:HG2	2.20	0.57
1:B:1176:ALA:O	1:B:1198:VAL:HG21	2.05	0.57
1:B:1132:LEU:HD23	1:B:1140:ILE:HD11	1.87	0.57
1:A:64:THR:CG2	1:A:65:ALA:H	2.19	0.55
1:A:115:GLN:HG2	4:A:944:HOH:O	2.09	0.52
1:A:120:LEU:HD13	1:A:176:ALA:HB2	1.91	0.52
1:A:199:GLU:HG2	4:A:993:HOH:O	2.09	0.52
1:B:1177:VAL:C	1:B:1198:VAL:HG23	2.29	0.52
1:B:1143:ARG:HB2	1:B:1153:LEU:HD21	1.93	0.51
1:B:1158:ILE:CD1	1:B:1190:ALA:HB2	2.40	0.51
1:A:61:ILE:O	2:C:26:VAL:HG22	2.11	0.51
1:A:89:THR:HG22	1:A:98:ILE:HA	1.93	0.50
1:B:1156:ARG:HG2	1:B:1160:TYR:CD1	2.47	0.49
1:A:136:HIS:O	1:A:137:ALA:HB3	2.12	0.49
1:A:132:LEU:HD12	4:A:946:HOH:O	2.11	0.49
1:B:1198:VAL:O	1:B:1201:LEU:HB2	2.12	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:60:GLN:HG2	1:A:71:ALA:HB3	1.96	0.48
1:A:128:SER:HB2	1:A:143:ARG:NH2	2.31	0.46
1:B:1156:ARG:NH1	1:B:1156:ARG:CG	2.74	0.46
1:A:158:ILE:HG12	1:A:185:CYS:SG	2.56	0.46
1:A:60:GLN:NE2	1:A:73:CYS:SG	2.87	0.45
1:B:1100:MET:HG3	1:B:1112:PRO:HD3	1.99	0.45
1:B:1156:ARG:NH1	1:B:1160:TYR:CE1	2.85	0.45
1:B:1142:VAL:HG22	1:B:1152:LEU:HD23	2.00	0.44
1:B:1070:LEU:HD21	1:B:1135:ARG:HG3	2.00	0.44
1:A:75:ASN:HD22	1:A:205:MET:CE	2.31	0.44
1:B:1069:PHE:N	1:B:1069:PHE:CD2	2.86	0.43
1:B:1156:ARG:NH1	1:B:1160:TYR:HE1	2.16	0.43
1:B:1158:ILE:HD13	1:B:1185:CYS:HB2	1.99	0.43
1:A:55:VAL:HG21	2:C:36:ALA:HA	2.02	0.42
1:B:1099:GLN:HG3	4:B:43:HOH:O	2.18	0.42
1:A:55:VAL:CG2	2:C:36:ALA:HA	2.50	0.41
1:A:116:GLY:HA2	4:A:974:HOH:O	2.19	0.41
1:B:1058:GLU:HA	1:B:1118:ARG:O	2.19	0.41
1:B:1186:THR:O	1:B:1187:ARG:CB	2.64	0.41
1:A:101:TYR:CE2	1:A:205:MET:HA	2.56	0.41
1:A:73:CYS:HA	1:A:77:VAL:O	2.21	0.41
1:A:130:LEU:HD22	1:A:177:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/200 (74%)	146 (98%)	3 (2%)	0	100	100
1	B	179/200 (90%)	169 (94%)	10 (6%)	0	100	100
2	C	14/21 (67%)	14 (100%)	0	0	100	100
2	D	19/21 (90%)	19 (100%)	0	0	100	100
All	All	361/442 (82%)	348 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/161 (76%)	120 (98%)	2 (2%)	75	93
1	B	143/161 (89%)	136 (95%)	7 (5%)	35	59
2	C	12/16 (75%)	11 (92%)	1 (8%)	16	29
2	D	15/16 (94%)	15 (100%)	0	100	100
All	All	292/354 (82%)	282 (97%)	10 (3%)	49	75

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	158	ILE
1	B	1039	LEU
1	B	1063	SER
1	B	1068	THR
1	B	1149	ARG
1	B	1156	ARG
1	B	1160	TYR
1	B	1185	CYS
2	C	22	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	67	GLN
1	A	75	ASN
1	A	200	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	151/200 (75%)	-0.14	3 (1%) 62 64	26, 41, 57, 64	0
1	B	181/200 (90%)	-0.16	3 (1%) 67 69	26, 44, 65, 79	0
2	C	16/21 (76%)	-0.48	1 (6%) 19 19	26, 32, 44, 53	0
2	D	21/21 (100%)	-0.48	0 100 100	29, 33, 50, 60	0
All	All	369/442 (83%)	-0.19	7 (1%) 64 66	26, 41, 63, 79	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	36	ALA	2.4
1	B	1185	CYS	2.4
1	A	166	GLY	2.4
1	B	1207	SER	2.3
1	B	1071	ALA	2.3
1	A	66	THR	2.1
1	A	71	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	A	901	1/1	0.13	2.64	41,41,41,41	0
3	ZN	B	902	1/1	0.07	-1.49	42,42,42,42	0

## 6.5 Other polymers

There are no such residues in this entry.