



wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 07:49 PM GMT

PDB ID : 3OI0
Title : Structure of the *Thermus thermophilus* 70S ribosome complexed with azithromycin. This file contains the 30S subunit of one 70S ribosome. The entire crystal structure contains two 70S ribosomes.
Authors : Bulkley, D.P.; Innis, C.A.; Blaha, G.; Steitz, T.A.
Deposited on : 2010-08-18
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)
RNA backbone	1838	1070 (3.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	1522	
2	B	256	
3	C	239	
4	D	209	
5	E	162	
6	F	101	
7	G	156	
8	H	138	
9	I	128	
10	J	105	
11	K	129	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
12	L	135	
13	M	126	
14	N	61	
15	O	89	
16	P	88	
17	Q	105	
18	R	88	
19	S	93	
20	T	106	
21	U	27	

2 Entry composition

There are 23 unique types of molecules in this entry. The entry contains 51469 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S rRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1504	Total	C	N	O	P	0	0	0
			32329	14390	5992	10444	1503			

- Molecule 2 is a protein called 30S ribosomal protein S2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	235	Total	C	N	O	S	0	0	1
			1901	1213	342	341	5			

- Molecule 3 is a protein called 30S ribosomal protein S3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	207	Total	C	N	O	S	0	0	1
			1613	1016	315	281	1			

- Molecule 4 is a protein called 30S ribosomal protein S4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	208	Total	C	N	O	S	0	0	0
			1703	1066	339	291	7			

- Molecule 5 is a protein called 30S ribosomal protein S5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	151	Total	C	N	O	S	0	0	1
			1147	724	218	201	4			

- Molecule 6 is a protein called 30S ribosomal protein S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			843	531	155	154	3			

- Molecule 7 is a protein called 30S ribosomal protein S7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	155	Total	C	N	O	S	0	0	0
			1257	781	252	218	6			

- Molecule 8 is a protein called 30S ribosomal protein S8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	138	Total	C	N	O	S	0	0	0
			1116	705	215	193	3			

- Molecule 9 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	127	Total	C	N	O	S	0	0	0
			1011	639	198	174				

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	58	ARG	HIS	CONFLICT	UNP P80374

- Molecule 10 is a protein called 30S ribosomal protein S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	99	Total	C	N	O	S	0	0	1
			795	499	157	138	1			

- Molecule 11 is a protein called 30S ribosomal protein S11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	119	Total	C	N	O	S	0	0	0
			885	549	168	165	3			

- Molecule 12 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	125	Total	C	N	O	S	0	0	1
			971	611	196	163	1			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	2	VAL	-	INSERTION	UNP Q5SHN3
L	3	ALA	-	INSERTION	UNP Q5SHN3
L	4	LEU	-	INSERTION	UNP Q5SHN3

- Molecule 13 is a protein called 30S ribosomal protein S13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	115	Total	C	N	O	S	0	0	0
			921	569	190	160	2			

- Molecule 14 is a protein called 30S ribosomal protein S14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	60	Total	C	N	O	S	0	0	0
			492	312	104	72	4			

- Molecule 15 is a protein called 30S ribosomal protein S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	O	88	Total	C	N	O	S	0	0	0
			734	459	147	126	2			

- Molecule 16 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
16	P	84	Total	C	N	O	S	0	0	1
			701	443	140	117	1			

- Molecule 17 is a protein called 30S ribosomal protein S17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
17	Q	100	Total	C	N	O	S	0	0	1
			824	528	152	142	2			

- Molecule 18 is a protein called 30S ribosomal protein S18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
18	R	70	Total	C	N	O		0	0	0
			574	367	112	95				

- Molecule 19 is a protein called 30S ribosomal protein S19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
19	S	79	Total	C	N	O	S	0	0	1
			630	403	115	110	2			

- Molecule 20 is a protein called 30S ribosomal protein S20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
20	T	99	Total	C	N	O	S	0	0	0
			763	470	162	129	2			

- Molecule 21 is a protein called 30S ribosomal protein Thx.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
21	U	25	Total	C	N	O	0	0	1
			209	128	51	30			


- Molecule 22 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
22	A	48	Total	Mg	0	0
			48	48		

- Molecule 23 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
23	D	1	Total	Zn	0	0
			1	1		
23	N	1	Total	Zn	0	0
			1	1		






• Molecule 7: 30S ribosomal protein S7

Chain G:




• Molecule 8: 30S ribosomal protein S8

Chain H:




• Molecule 9: 30S ribosomal protein S9

Chain I:




• Molecule 10: 30S ribosomal protein S10

Chain J:




• Molecule 11: 30S ribosomal protein S11

Chain K:




- Molecule 12: 30S ribosomal protein S12

Chain L:




- Molecule 13: 30S ribosomal protein S13

Chain M:




- Molecule 14: 30S ribosomal protein S14

Chain N:



- Molecule 15: 30S ribosomal protein S15


Chain O:



- Molecule 16: 30S ribosomal protein S16


Chain P:






- Molecule 17: 30S ribosomal protein S17

Chain Q:




- Molecule 18: 30S ribosomal protein S18

Chain R:





- Molecule 19: 30S ribosomal protein S19

Chain S:




- Molecule 20: 30S ribosomal protein S20

Chain T:



- Molecule 21: 30S ribosomal protein Thx

Chain U:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	210.22Å 450.25Å 623.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.80 – 3.00 49.80 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.80-3.00) 88.7 (49.80-3.00)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 3.01Å)	Xtriage
Refinement program	Phenix	Depositor
R, R_{free}	0.235 , 0.269 0.506 , 0.506	Depositor DCC
R_{free} test set	51892 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	73.9	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 72.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 1035238 reflections	Xtriage
F_o, F_c correlation	0.50	EDS
Total number of atoms	51469	wwPDB-VP
Average B, all atoms (Å ²)	136.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/36190	0.88	40/56486 (0.1%)
2	B	0.29	0/1936	0.50	0/2611
3	C	0.27	0/1637	0.45	0/2207
4	D	0.34	0/1733	0.53	0/2318
5	E	0.34	0/1163	0.55	0/1566
6	F	0.36	0/856	0.54	0/1154
7	G	0.26	0/1276	0.44	0/1709
8	H	0.33	0/1136	0.54	0/1527
9	I	0.27	0/1028	0.44	0/1375
10	J	0.29	0/808	0.48	0/1087
11	K	0.32	0/900	0.52	0/1213
12	L	0.39	0/987	0.62	0/1322
13	M	0.27	0/928	0.47	0/1238
14	N	0.28	0/501	0.44	0/664
15	O	0.33	0/745	0.56	0/992
16	P	0.33	0/717	0.55	0/965
17	Q	0.34	0/837	0.56	0/1119
18	R	0.37	0/579	0.57	0/768
19	S	0.28	0/643	0.46	0/867
20	T	0.34	0/765	0.55	0/1007
21	U	0.28	0/213	0.43	0/279
All	All	0.44	0/55578	0.78	40/82474 (0.0%)

There are no bond length outliers.

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	893	C	C6-N1-C2	8.02	123.51	120.30
1	A	1509	C	C6-N1-C2	7.49	123.30	120.30
1	A	34	C	C6-N1-C2	6.90	123.06	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	895	G	N1-C6-O6	6.57	123.84	119.90
1	A	909	A	C8-N9-C4	6.55	108.42	105.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	32329	0	16318	1374	0
2	B	1901	0	1951	167	0
3	C	1613	0	1677	117	0
4	D	1703	0	1763	160	0
5	E	1147	0	1207	107	0
6	F	843	0	857	86	0
7	G	1257	0	1296	62	0
8	H	1116	0	1177	82	0
9	I	1011	0	1042	85	0
10	J	795	0	840	82	0
11	K	885	0	904	69	0
12	L	971	0	1057	106	0
13	M	921	0	976	61	0
14	N	492	0	529	33	0
15	O	734	0	771	54	0
16	P	701	0	720	91	0
17	Q	824	0	891	49	0
18	R	574	0	644	64	0
19	S	630	0	652	34	0
20	T	763	0	861	75	0
21	U	209	0	221	11	0
22	A	48	0	0	0	0
23	D	1	0	0	0	0
23	N	1	0	0	0	0
All	All	51469	0	36354	2706	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 31.

The worst 5 of 2706 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:111:ARG:HH11	2:B:111:ARG:HG2	1.09	1.14
7:G:113:GLU:HB2	7:G:119:ARG:HG2	1.33	1.11
2:B:185:ILE:HG22	2:B:199:TYR:HB2	1.32	1.09
1:A:59:A:H5"	1:A:60:A:H5"	1.28	1.09
4:D:128:VAL:HG12	4:D:129:ASN:ND2	1.69	1.06

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	233/256 (91%)	177 (76%)	39 (17%)	17 (7%)	2	8
3	C	205/239 (86%)	155 (76%)	37 (18%)	13 (6%)	2	12
4	D	206/209 (99%)	137 (66%)	55 (27%)	14 (7%)	2	10
5	E	149/162 (92%)	103 (69%)	33 (22%)	13 (9%)	1	5
6	F	99/101 (98%)	76 (77%)	14 (14%)	9 (9%)	1	5
7	G	153/156 (98%)	131 (86%)	18 (12%)	4 (3%)	8	39
8	H	136/138 (99%)	98 (72%)	31 (23%)	7 (5%)	3	18
9	I	123/128 (96%)	94 (76%)	22 (18%)	7 (6%)	3	16
10	J	97/105 (92%)	81 (84%)	11 (11%)	5 (5%)	3	18
11	K	117/129 (91%)	86 (74%)	27 (23%)	4 (3%)	6	31
12	L	123/135 (91%)	83 (68%)	29 (24%)	11 (9%)	1	5
13	M	107/126 (85%)	84 (78%)	17 (16%)	6 (6%)	3	16
14	N	58/61 (95%)	44 (76%)	12 (21%)	2 (3%)	6	31
15	O	86/89 (97%)	61 (71%)	21 (24%)	4 (5%)	4	21
16	P	82/88 (93%)	47 (57%)	29 (35%)	6 (7%)	2	8

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
17	Q	98/105 (93%)	73 (74%)	19 (19%)	6 (6%)	2	14
18	R	68/88 (77%)	51 (75%)	13 (19%)	4 (6%)	2	14
19	S	77/93 (83%)	59 (77%)	12 (16%)	6 (8%)	1	7
20	T	97/106 (92%)	65 (67%)	23 (24%)	9 (9%)	1	5
21	U	23/27 (85%)	18 (78%)	4 (17%)	1 (4%)	4	23
All	All	2337/2541 (92%)	1723 (74%)	466 (20%)	148 (6%)	2	12

5 of 148 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	15	VAL
2	B	24	TRP
2	B	84	GLU
2	B	154	LEU
2	B	165	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	202/220 (92%)	176 (87%)	26 (13%)	6	26
3	C	160/188 (85%)	152 (95%)	8 (5%)	34	77
4	D	180/181 (99%)	156 (87%)	24 (13%)	6	25
5	E	115/123 (94%)	100 (87%)	15 (13%)	6	26
6	F	90/90 (100%)	79 (88%)	11 (12%)	7	29
7	G	126/127 (99%)	121 (96%)	5 (4%)	42	84
8	H	119/119 (100%)	107 (90%)	12 (10%)	11	39
9	I	98/99 (99%)	88 (90%)	10 (10%)	11	38
10	J	88/92 (96%)	81 (92%)	7 (8%)	17	53
11	K	90/99 (91%)	80 (89%)	10 (11%)	9	34
12	L	104/111 (94%)	96 (92%)	8 (8%)	18	56
13	M	93/101 (92%)	86 (92%)	7 (8%)	19	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
14	N	49/50 (98%)	47 (96%)	2 (4%)	41	83
15	O	79/80 (99%)	69 (87%)	10 (13%)	6	27
16	P	72/74 (97%)	60 (83%)	12 (17%)	3	16
17	Q	94/97 (97%)	91 (97%)	3 (3%)	51	89
18	R	61/77 (79%)	55 (90%)	6 (10%)	12	41
19	S	69/80 (86%)	62 (90%)	7 (10%)	11	39
20	T	76/82 (93%)	66 (87%)	10 (13%)	6	25
21	U	19/22 (86%)	19 (100%)	0	100	100
All	All	1984/2112 (94%)	1791 (90%)	193 (10%)	12	42

5 of 193 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	G	151	TYR
9	I	114	TYR
19	S	22	LEU
8	H	1	MET
8	H	95	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
6	F	94	GLN
8	H	82	HIS
16	P	82	GLN
7	G	13	GLN
7	G	84	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1503/1522 (98%)	288 (19%)	31 (2%)

5 of 288 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	9	G

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	31	G
1	A	32	A
1	A	39	G
1	A	47	C

5 of 31 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	533	A
1	A	748	C
1	A	1493	A
1	A	560	U
1	A	776	G

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 50 ligands modelled in this entry, 50 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1504/1522 (98%)	0.43	125 (8%) 11 3	61, 125, 191, 194	0
2	B	235/256 (91%)	0.05	5 (2%) 60 12	107, 158, 185, 191	0
3	C	207/239 (86%)	0.22	15 (7%) 15 4	119, 166, 184, 191	0
4	D	208/209 (99%)	0.25	20 (9%) 8 2	82, 131, 168, 182	0
5	E	151/162 (93%)	0.25	14 (9%) 9 2	84, 117, 162, 189	0
6	F	101/101 (100%)	-0.16	3 (2%) 48 9	86, 132, 165, 182	0
7	G	155/156 (99%)	0.40	16 (10%) 7 2	140, 171, 188, 190	0
8	H	138/138 (100%)	-0.01	7 (5%) 27 6	85, 123, 156, 162	0
9	I	127/128 (99%)	0.68	18 (14%) 3 1	143, 183, 190, 191	0
10	J	99/105 (94%)	0.96	20 (20%) 2 1	130, 177, 190, 193	0
11	K	119/129 (92%)	0.16	2 (1%) 67 15	84, 123, 165, 186	0
12	L	125/135 (92%)	0.47	14 (11%) 6 2	82, 109, 164, 189	0
13	M	115/126 (91%)	0.49	13 (11%) 6 2	149, 185, 190, 192	0
14	N	60/61 (98%)	0.62	2 (3%) 44 8	132, 170, 186, 189	0
15	O	88/89 (98%)	-0.11	2 (2%) 57 12	74, 112, 159, 165	0
16	P	84/88 (95%)	0.75	14 (16%) 2 1	89, 116, 160, 180	0
17	Q	100/105 (95%)	0.43	13 (13%) 4 1	85, 110, 153, 159	0
18	R	70/88 (79%)	0.19	3 (4%) 34 7	93, 122, 171, 183	0
19	S	79/93 (84%)	1.24	16 (20%) 1 1	142, 186, 191, 192	0
20	T	99/106 (93%)	0.39	9 (9%) 9 2	84, 119, 157, 179	0
21	U	25/27 (92%)	2.66	14 (56%) 0 0	141, 172, 188, 189	0
All	All	3889/4063 (95%)	0.39	345 (8%) 10 3	61, 141, 190, 194	0

The worst 5 of 345 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1149	C	19.5
1	A	1148	U	17.2
1	A	1126	U	11.1
1	A	1150	U	10.9
1	A	1322	C	10.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
22	MG	A	1550	1/1	0.75	-	72,72,72,72	0
22	MG	A	1571	1/1	0.19	-	66,66,66,66	0
22	MG	A	1574	1/1	0.18	-	77,77,77,77	0
22	MG	A	1578	1/1	1.10	-	87,87,87,87	0
22	MG	A	1572	1/1	0.60	-	75,75,75,75	0
22	MG	A	1585	1/1	1.08	-	87,87,87,87	0
23	ZN	D	210	1/1	0.04	-	93,93,93,93	0
22	MG	A	1580	1/1	1.68	-	79,79,79,79	0
22	MG	A	1573	1/1	0.12	-	82,82,82,82	0
22	MG	A	1545	1/1	0.51	-	83,83,83,83	0
22	MG	A	1577	1/1	0.37	-	87,87,87,87	0
22	MG	A	1583	1/1	0.13	-	64,64,64,64	0
22	MG	A	1556	1/1	0.11	-	77,77,77,77	0
22	MG	A	1568	1/1	0.26	-	65,65,65,65	0
22	MG	A	1563	1/1	0.62	-	75,75,75,75	0
22	MG	A	1582	1/1	0.23	-	71,71,71,71	0
23	ZN	N	62	1/1	0.10	-	157,157,157,157	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
22	MG	A	1575	1/1	0.96	-	70,70,70,70	0
22	MG	A	1579	1/1	0.85	-	86,86,86,86	0
22	MG	A	1592	1/1	0.33	-	79,79,79,79	0
22	MG	A	1551	1/1	0.10	-	82,82,82,82	0
22	MG	A	1576	1/1	0.17	-	79,79,79,79	0
22	MG	A	1581	1/1	0.64	-	80,80,80,80	0
22	MG	A	1569	1/1	0.25	-	74,74,74,74	0
22	MG	A	1548	1/1	0.10	-	86,86,86,86	0
22	MG	A	1553	1/1	0.23	-	94,94,94,94	0
22	MG	A	1587	1/1	0.26	-	62,62,62,62	0
22	MG	A	1557	1/1	0.97	-	80,80,80,80	0
22	MG	A	1547	1/1	0.40	-	63,63,63,63	0
22	MG	A	1562	1/1	0.41	-	62,62,62,62	0
22	MG	A	1567	1/1	0.23	-	67,67,67,67	0
22	MG	A	1554	1/1	0.08	-	61,61,61,61	0
22	MG	A	1561	1/1	0.23	-	61,61,61,61	0
22	MG	A	1555	1/1	0.13	-	81,81,81,81	0
22	MG	A	1549	1/1	0.89	-	61,61,61,61	0
22	MG	A	1560	1/1	0.17	-	73,73,73,73	0
22	MG	A	1590	1/1	0.57	-	80,80,80,80	0
22	MG	A	1552	1/1	0.26	-	51,51,51,51	0
22	MG	A	1564	1/1	0.49	-	66,66,66,66	0
22	MG	A	1558	1/1	1.37	-	76,76,76,76	0
22	MG	A	1589	1/1	0.24	-	97,97,97,97	0
22	MG	A	1584	1/1	0.39	-	68,68,68,68	0
22	MG	A	1570	1/1	0.35	-	78,78,78,78	0
22	MG	A	1591	1/1	0.45	-	84,84,84,84	0
22	MG	A	1559	1/1	0.19	-	67,67,67,67	0
22	MG	A	1566	1/1	0.39	-	78,78,78,78	0
22	MG	A	1586	1/1	0.12	-	62,62,62,62	0
22	MG	A	1546	1/1	2.10	-	70,70,70,70	0
22	MG	A	1565	1/1	0.09	-	68,68,68,68	0
22	MG	A	1588	1/1	0.15	-	74,74,74,74	0

6.5 Other polymers ⓘ

There are no such residues in this entry.