



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2014 – 05:29 PM EDT

PDB ID : 4OI4
Title : Yeast Clp1/Pcf11 complex
Authors : Dikfidan, A.; Loll, B.; Zeymer, C.; Clausen, T.; Meinhart, A.
Deposited on : 2014-01-18
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23161
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23161

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	453	
1	C	453	
2	B	115	
2	D	115	
2	U	115	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	GOL	C	503	-	X
5	GOL	C	504	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7797 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mRNA cleavage and polyadenylation factor CLP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	19	0
			3502	2259	577	654	12			
1	C	427	Total	C	N	O	S	0	14	0
			3476	2240	579	645	12			


- Molecule 2 is a protein called Protein PCF11.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	25	Total	C	N	O	0	1	0
			215	135	36	44			
2	D	24	Total	C	N	O	0	2	0
			214	134	34	46			
2	U	8	Total	C	N	O	0	0	0
			61	37	10	14			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	451	PRO	-	EXPRESSION TAG	UNP P39081
U	452	GLU	-	EXPRESSION TAG	UNP P39081
U	453	PHE	-	EXPRESSION TAG	UNP P39081

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	C	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	C	1	Total Mg 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	120	Total	O	0	0
			120	120		
6	C	90	Total	O	0	0
			90	90		
6	B	20	Total	O	0	0
			20	20		
6	D	15	Total	O	0	0
			15	15		
6	U	2	Total	O	0	0
			2	2		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: mRNA cleavage and polyadenylation factor CLP1

Chain A: 




- Molecule 1: mRNA cleavage and polyadenylation factor CLP1

Chain C: 




- Molecule 2: Protein PCF11

Chain B: 



- Molecule 2: Protein PCF11

Chain D: 



GLU	GLY	SER	VAL	ASP	ASN	THR	LEU	GLY	SER	ASP	ARG	SER	ASN	GLU	LEU	GLU	ILE	ARG	GLY	LYS	TYR	VAL	VAL	PRO	GLU	THR	SER	GLN	ASP	MET	ALA	PHE	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

● Molecule 2: Protein PCF11

Chain U: 

GLY	SER	P451	E452	T457	T458	ALA	ASN	THR	GLY	ILE	SER	ASN	SER	ASN	ASN	LEU	ASN	THR	THR	THR	ARG	LYS	ASN	ILE	GLN	SER	ARG	ASN	TRP	TYR	LEU	SER	ILE	ASP	SER	GLN	TRP	ALA	ALA	PHE	LYS	ASP	ASP	GLU	ILE	THR	SER	THR	LYS	HIS	LYS	LYS	ASN	ASP	TYR	THR	ASP	PRO	HIS	ALA
-----	-----	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

ASN	LYS	ASN	ILE	ASP	LYS	SER	ALA	LEU	ASN	ILE	ILE	HIS	ALA	ASP	GLU	ASN	ASP	GLU	GLY	SER	VAL	ASN	THR	LEU	GLY	SER	ASP	ARG	SER	ASN	GLU	LEU	GLU	ILE	ARG	GLY	LYS	TYR	VAL	VAL	PHE	VAL	PRO	GLU	THR	SER	GLN	ASP	MET	ALA	PHE	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.64Å 95.77Å 181.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 2.40 19.99 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.99-2.40) 99.0 (19.99-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.62 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.221 0.182 , 0.213	Depositor DCC
R_{free} test set	3061 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 61162 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7797	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3636	0.61	0/4947
1	C	0.62	3/3595 (0.1%)	0.58	0/4888
2	B	0.64	0/222	0.64	0/298
2	D	0.56	0/225	0.52	0/305
2	U	0.44	0/62	0.41	0/82
All	All	0.56	3/7740 (0.0%)	0.59	0/10520

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	19	GLY	N-CA	22.62	1.79	1.46
1	C	22	GLU	CD-OE2	10.32	1.36	1.25
1	C	22	GLU	CD-OE1	5.89	1.32	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3502	0	3588	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3476	0	3554	18	0
2	B	215	0	197	1	0
2	D	214	0	189	0	0
2	U	61	0	51	0	0
3	A	31	0	12	0	0
3	C	31	0	12	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	6	0	8	0	0
5	C	12	0	16	2	0
6	A	120	0	0	2	0
6	B	20	0	0	0	0
6	C	90	0	0	0	0
6	D	15	0	0	0	0
6	U	2	0	0	0	0
All	All	7797	0	7627	53	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (53) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:19:GLY:N	1:C:19:GLY:CA	1.80	1.44
1:A:109[B]:LEU:CD2	1:A:278[B]:LEU:HD21	1.86	1.05
1:A:109[B]:LEU:HD22	1:A:278[B]:LEU:HD21	1.48	0.96
1:A:109[B]:LEU:HD21	1:A:278[B]:LEU:CD2	1.98	0.92
1:A:439:THR:HG22	1:A:441:TYR:H	1.39	0.86
1:A:109[B]:LEU:CD2	1:A:278[B]:LEU:CD2	2.54	0.82
1:A:385:ILE:O	1:A:439:THR:HB	1.82	0.80
1:A:164[B]:GLN:HG3	6:A:698:HOH:O	1.90	0.71
1:A:439:THR:CG2	1:A:441:TYR:H	2.04	0.70
1:A:109[B]:LEU:HD21	1:A:278[B]:LEU:HD21	1.63	0.67
1:C:126[B]:VAL:HG12	1:C:276:ILE:HB	1.77	0.67
1:A:109[B]:LEU:HD21	1:A:278[B]:LEU:HD22	1.76	0.67
1:C:109:LEU:HD21	1:C:278[B]:LEU:HD13	1.77	0.66
1:A:359[A]:SER:OG	1:A:370:GLU:HG3	1.96	0.64
1:C:133:GLN:HG3	1:C:315:ALA:HA	1.79	0.64
1:A:109[A]:LEU:HD11	1:A:307:ILE:HD11	1.80	0.63
1:A:128:ILE:O	1:A:252[B]:THR:HG22	2.00	0.62
1:A:55:ILE:HD12	1:A:60:LEU:HD21	1.82	0.60
1:A:126[B]:VAL:HG12	1:A:276:ILE:HB	1.84	0.60
1:A:57:GLY:O	1:A:149:LYS:NZ	2.36	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:164[B]:GLN:HG2	6:A:643:HOH:O	2.04	0.58
1:A:280:LEU:HD23	1:A:307:ILE:HB	1.86	0.56
1:A:19:GLY:O	1:A:20:ASP:HB3	2.05	0.56
1:C:348:VAL:HG11	1:C:431:LEU:HD22	1.88	0.54
1:C:67:THR:HB	5:C:504:GOL:H32	1.90	0.53
1:C:125:ARG:HD2	1:C:273:ASN:O	2.12	0.49
1:A:252[B]:THR:OG1	1:A:253:PRO:HD2	2.13	0.48
1:C:154:GLN:HG2	1:C:154:GLN:O	2.14	0.48
1:A:245[B]:ARG:HD3	1:C:238:HIS:CD2	2.48	0.48
1:A:439:THR:HG22	1:A:441:TYR:N	2.19	0.47
1:A:346:ILE:HG13	1:A:423[B]:VAL:HG13	1.95	0.47
1:C:288:TRP:CZ3	1:C:306:PHE:HB3	2.51	0.46
1:A:167:PHE:CE1	1:A:203:LYS:HG3	2.50	0.46
1:A:18:THR:HB	1:A:91:THR:HG21	1.99	0.45
1:C:278[A]:LEU:HD12	1:C:305:PHE:HB2	2.00	0.44
1:A:427:VAL:HG22	2:B:480:ARG:O	2.18	0.43
1:A:348:VAL:HG11	1:A:431:LEU:HD22	2.00	0.43
1:C:357:LYS:HB2	1:C:374:VAL:CG2	2.48	0.43
1:C:192:SER:HA	1:C:203:LYS:O	2.18	0.43
1:C:67:THR:HB	5:C:504:GOL:C3	2.49	0.43
1:C:237:LEU:CD2	1:C:243:VAL:HG12	2.49	0.43
1:A:109[A]:LEU:CD1	1:A:307:ILE:HD11	2.49	0.42
1:C:188:THR:O	1:C:189:TRP:HB2	2.19	0.42
1:C:19:GLY:C	1:C:19:GLY:N	2.64	0.42
1:A:169:VAL:HB	1:A:170:PRO:HD2	2.01	0.42
1:A:26:LEU:HD13	1:A:95:ILE:HD11	2.02	0.42
1:A:188:THR:O	1:A:189:TRP:HB2	2.20	0.41
1:A:283:GLU:HG3	1:A:306:PHE:CE2	2.55	0.41
1:A:126[A]:VAL:HG11	1:A:278[A]:LEU:HD11	2.03	0.41
1:C:353:LEU:HD13	1:C:355:ILE:HD11	2.02	0.41
1:A:128:ILE:HG12	1:A:278[A]:LEU:HD12	2.03	0.41
1:A:109[B]:LEU:HD22	1:A:278[B]:LEU:CD2	2.30	0.41
1:A:353:LEU:HD13	1:A:355:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/453 (98%)	426 (96%)	18 (4%)	1 (0%)	56	74
1	C	439/453 (97%)	424 (97%)	14 (3%)	1 (0%)	56	74
2	B	23/115 (20%)	23 (100%)	0	0	100	100
2	D	24/115 (21%)	24 (100%)	0	0	100	100
2	U	6/115 (5%)	6 (100%)	0	0	100	100
All	All	937/1251 (75%)	903 (96%)	32 (3%)	2 (0%)	56	74

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	40	ALA
1	A	20	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/407 (99%)	398 (98%)	6 (2%)	76	90
1	C	398/407 (98%)	397 (100%)	1 (0%)	96	99
2	B	24/102 (24%)	24 (100%)	0	100	100
2	D	24/102 (24%)	24 (100%)	0	100	100
2	U	7/102 (7%)	7 (100%)	0	100	100
All	All	857/1120 (76%)	850 (99%)	7 (1%)	91	96

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	VAL
1	A	95	ILE
1	A	259[A]	ASP
1	A	259[B]	ASP
1	A	275	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	SER
1	C	284	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ATP	A	501	4	33,33,33	1.13	2 (6%)	52,52,52	1.86	7 (13%)
5	GOL	A	503	-	5,5,5	0.27	0	5,5,5	0.46	0
3	ATP	C	501	4	33,33,33	1.07	2 (6%)	52,52,52	1.84	6 (11%)
5	GOL	C	503	-	5,5,5	0.29	0	5,5,5	0.42	0
5	GOL	C	504	-	5,5,5	0.31	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	501	4	-	0/22/38/38	0/3/3/3
5	GOL	A	503	-	-	0/4/4/4	0/0/0/0
3	ATP	C	501	4	-	0/22/38/38	0/3/3/3
5	GOL	C	503	-	-	0/4/4/4	0/0/0/0
5	GOL	C	504	-	-	0/4/4/4	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	501	ATP	C5-C4	3.26	1.47	1.40
3	A	501	ATP	C5-C4	3.20	1.47	1.40
3	C	501	ATP	C4-N9	-2.72	1.33	1.37
3	A	501	ATP	C4-N9	-2.70	1.33	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	501	ATP	C5-C4-N3	-7.28	118.88	125.98
3	A	501	ATP	C5-C4-N3	-6.66	119.49	125.98
3	A	501	ATP	N3-C2-N1	-6.61	123.08	128.89
3	C	501	ATP	N3-C2-N1	-5.69	123.88	128.89
3	C	501	ATP	N3-C4-N9	5.45	134.75	125.39
3	A	501	ATP	N3-C4-N9	5.20	134.31	125.39
3	C	501	ATP	C4-C5-N7	-2.99	106.52	109.41
3	A	501	ATP	C4-C5-N7	-2.95	106.56	109.41
3	A	501	ATP	C2-N3-C4	2.58	120.71	113.27
3	C	501	ATP	C2-N3-C4	2.57	120.66	113.27
3	A	501	ATP	C8-N9-C4	2.54	109.02	106.96
3	C	501	ATP	PA-O3A-PB	-2.38	125.34	131.93
3	A	501	ATP	C1'-N9-C4	-2.27	122.71	126.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/453 (94%)	-0.24	11 (2%) 53 51	21, 41, 74, 103	0
1	C	427/453 (94%)	-0.13	13 (3%) 48 45	25, 49, 87, 110	0
2	B	25/115 (21%)	-0.53	1 (4%) 36 34	25, 30, 57, 71	0
2	D	24/115 (20%)	-0.18	3 (12%) 5 4	24, 36, 74, 91	0
2	U	8/115 (6%)	1.56	4 (50%) 0 0	40, 72, 107, 139	0
All	All	912/1251 (72%)	-0.18	32 (3%) 42 40	21, 44, 81, 139	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	284	THR	5.2
1	A	40	ALA	5.0
1	A	309	LYS	4.9
2	U	458	THR	4.8
1	C	40	ALA	4.6
1	A	18	THR	4.4
1	C	199	LEU	4.2
1	C	93	ASN	4.0
1	C	362	PHE	3.9
2	U	451	PRO	3.7
2	D	499	SER	3.7
2	D	476	ASN	3.6
1	A	362	PHE	3.3
1	A	42	GLY	3.2
1	C	41	GLU	3.2
2	B	499[A]	SER	3.0
1	A	41	GLU	3.0
1	A	394	ARG	3.0
2	U	452	GLU	2.9
1	C	284	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	394	ARG	2.8
1	C	282	SER	2.8
1	C	418	ARG	2.8
1	A	417	LYS	2.6
1	C	309	LYS	2.5
1	C	63	ASP	2.4
1	C	94	THR	2.3
2	D	477	ILE	2.3
2	U	457	ASN	2.2
1	A	366	VAL	2.1
1	A	282	SER	2.1
1	C	444	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	GOL	C	504	6/6	0.40	6.35	60,61,61,62	0
5	GOL	C	503	6/6	0.18	2.82	53,55,56,56	0
5	GOL	A	503	6/6	0.15	1.57	46,49,49,49	0
3	ATP	A	501	31/31	0.11	-0.09	34,36,44,45	0
3	ATP	C	501	31/31	0.12	-0.37	37,40,43,44	0
4	MG	A	502	1/1	0.06	-2.03	28,28,28,28	0
4	MG	C	502	1/1	0.05	-3.14	36,36,36,36	0

6.5 Other polymers ⓘ

There are no such residues in this entry.