



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 03:04 AM GMT

PDB ID : 1OJO
Title : SPECIFICITY AND MECHANISM OF STREPTOCOCCUS PNEUMONIAE HYALURONATE LYASE: COMPLEX OF THE TYR408PHE MUTANT WITH 4-SULPHATED CHONDROITIN DISACCHARIDE
Authors : Rigden, D.J.; Jedrzejewski, M.J.
Deposited on : 2003-07-11
Resolution : 1.75 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance


The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1134 (1.76-1.76)
Clashscore	79885	1304 (1.76-1.76)
Ramachandran outliers	78287	1288 (1.76-1.76)
Sidechain outliers	78261	1288 (1.76-1.76)
RSRZ outliers	66119	1135 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	731	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	1201	-	X
3	SO4	A	1203	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6581 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYALURONATE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	722	Total	C	N	O	S	0	23	0
			5995	3770	1006	1196	23			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	THR	ALA	CONFLICT	UNP Q54873
A	196	ASP	GLU	SEE REMARK 999	UNP Q54873
A	223	ILE	THR	CONFLICT	UNP Q54873
A	408	PHE	TYR	ENGINEERED MUTATION	UNP Q54873
A	496	ARG	CYS	CONFLICT	UNP Q54873
A	541	THR	PRO	CONFLICT	UNP Q54873
A	704	SER	GLY	CONFLICT	UNP Q54873
A	736	SER	PHE	CONFLICT	UNP Q54873


- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	2	Total	C	N	O	S	0	0
			30	14	1	14	1		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	THR	ALA	CONFLICT	UNP Q54873
A	196	ASP	GLU	SEE REMARK 999	UNP Q54873
A	223	ILE	THR	CONFLICT	UNP Q54873
A	408	PHE	TYR	ENGINEERED MUTATION	UNP Q54873
A	496	ARG	CYS	CONFLICT	UNP Q54873
A	541	THR	PRO	CONFLICT	UNP Q54873
A	704	SER	GLY	CONFLICT	UNP Q54873
A	736	SER	PHE	CONFLICT	UNP Q54873

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	536	Total	O	0	0
			536	536		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HYALURONATE LYASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.74Å 103.78Å 101.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.96 – 1.75 40.43 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.96-1.75) 86.9 (40.43-1.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 1.50Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.202 , 0.219 0.210 , 0.227	Depositor DCC
R_{free} test set	4429 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.6	EDS
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	3 of 122689 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6581	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ASG, GCD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/6116	0.60	0/8251

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5995	0	5803	90	0
2	A	30	0	19	0	0
3	A	20	0	0	1	0
4	A	536	0	0	9	0
All	All	6581	0	5822	90	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (90) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:613:ASN:H	1:A:698:GLN:HE22	1.03	0.97
1:A:315:ASP:O	1:A:319[B]:LYS:HG3	1.87	0.75
1:A:248:MET:O	1:A:252[A]:VAL:HG23	1.91	0.70
1:A:708:LYS:HE3	1:A:710:TYR:OH	1.95	0.67
1:A:223:ILE:HG13	1:A:224:TYR:CD2	2.31	0.66
1:A:529:LEU:HD23	1:A:529:LEU:C	2.17	0.65
1:A:282:ASN:ND2	1:A:285:LYS:HG2	2.13	0.64
1:A:581:LYS:HB3	1:A:768:GLU:HB2	1.81	0.62
1:A:704[A]:SER:C	1:A:705:ASN:HD22	2.03	0.62
1:A:874:ASN:HD22	1:A:874:ASN:C	2.03	0.62
1:A:839:LYS:HD2	1:A:853:VAL:HG23	1.83	0.59
1:A:863:LYS:HE2	1:A:865[A]:GLY:O	2.01	0.59
1:A:422:VAL:HA	4:A:2007:HOH:O	2.03	0.58
1:A:626:LYS:H	1:A:626:LYS:CD	2.16	0.58
1:A:282:ASN:HD22	1:A:282:ASN:C	2.07	0.58
1:A:704[B]:SER:C	1:A:705:ASN:HD22	2.05	0.58
1:A:865[B]:GLY:C	1:A:867:GLU:H	2.06	0.58
1:A:254:ASN:HD22	1:A:254:ASN:C	2.06	0.57
1:A:492:GLU:OE2	1:A:496:ARG:HD3	2.06	0.55
1:A:502:LYS:HB2	1:A:529:LEU:HD21	1.87	0.55
1:A:440:HIS:CE1	1:A:444[B]:LYS:HD2	2.42	0.55
1:A:254:ASN:ND2	1:A:256:SER:H	2.05	0.55
1:A:626:LYS:H	1:A:626:LYS:HD3	1.72	0.54
1:A:864:GLU:O	1:A:865[B]:GLY:C	2.46	0.54
1:A:243:ARG:O	1:A:247:GLU:HG3	2.07	0.54
1:A:650:ASN:HD21	1:A:832:GLN:HE22	1.55	0.54
1:A:373:THR:O	1:A:377:ILE:HG12	2.08	0.54
1:A:613:ASN:N	1:A:698:GLN:HE22	1.88	0.53
1:A:646:LEU:HD21	1:A:862[A]:ARG:HB2	1.91	0.53
1:A:502:LYS:CB	1:A:529:LEU:HD21	2.39	0.52
1:A:248:MET:O	1:A:252[B]:VAL:HG13	2.08	0.52
1:A:440:HIS:NE2	1:A:444[B]:LYS:HD2	2.25	0.52
1:A:264:THR:HG22	4:A:2075:HOH:O	2.09	0.51
1:A:626:LYS:N	1:A:626:LYS:HD3	2.27	0.50
1:A:252[B]:VAL:HG23	1:A:253:THR:HG23	1.93	0.49
1:A:327:LYS:HD3	1:A:327:LYS:C	2.32	0.49
1:A:331:ASP:OD2	1:A:334:HIS:ND1	2.43	0.49
1:A:822:GLU:CD	1:A:822:GLU:H	2.16	0.49
1:A:584:LYS:HE2	3:A:1200:SO4:O1	2.12	0.49
1:A:848:SER:O	1:A:850:GLN:HG3	2.12	0.49
1:A:864:GLU:O	1:A:867:GLU:HB2	2.12	0.48
1:A:298:THR:HB	1:A:299:PRO:HD3	1.94	0.48
1:A:420:LEU:HD13	1:A:435:MET:CE	2.44	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:374:ILE:O	1:A:378:GLU:HG3	2.14	0.48
1:A:314:SER:OG	1:A:317:GLU:HG3	2.14	0.47
1:A:808[B]:GLN:HG2	4:A:2479:HOH:O	2.14	0.47
1:A:820:ASN:HD22	1:A:825:GLN:HG2	1.78	0.47
1:A:650:ASN:ND2	1:A:862[B]:ARG:HH21	2.13	0.47
1:A:808[B]:GLN:HG3	4:A:2484:HOH:O	2.14	0.47
1:A:811:LYS:HE3	4:A:2481:HOH:O	2.14	0.47
1:A:845:SER:HB2	1:A:853:VAL:HG23	1.97	0.46
1:A:228[B]:LYS:HG2	1:A:228[B]:LYS:O	2.15	0.46
1:A:540[B]:ARG:HG2	1:A:540[B]:ARG:HH11	1.80	0.46
1:A:584:LYS:NZ	1:A:768:GLU:HG2	2.31	0.46
1:A:242:TYR:CD2	1:A:298:THR:HG23	2.52	0.45
1:A:671:PHE:HB2	1:A:678:ALA:HB3	1.99	0.45
1:A:642:GLY:HA2	4:A:2300:HOH:O	2.17	0.45
1:A:705:ASN:N	1:A:705:ASN:HD22	2.12	0.44
1:A:864:GLU:O	1:A:865[B]:GLY:O	2.36	0.44
1:A:626:LYS:H	1:A:626:LYS:HZ3	1.65	0.44
1:A:404:TYR:CE1	1:A:461:GLY:HA3	2.53	0.43
1:A:549[A]:LYS:HE3	4:A:2515:HOH:O	2.19	0.43
1:A:254:ASN:ND2	1:A:254:ASN:C	2.68	0.43
1:A:882:PRO:HB2	1:A:884:GLN:NE2	2.33	0.43
1:A:758[B]:LEU:HD23	1:A:778:GLU:HB2	1.99	0.43
1:A:354:GLY:O	1:A:358:VAL:HB	2.20	0.42
1:A:661:ASN:HD22	1:A:661:ASN:C	2.23	0.42
1:A:870:ILE:HD11	1:A:887:PHE:CD2	2.55	0.42
1:A:341:ASN:N	1:A:342:PRO:HD3	2.34	0.42
1:A:483:HIS:NE2	1:A:529:LEU:HD12	2.35	0.42
1:A:671:PHE:N	1:A:671:PHE:CD1	2.87	0.42
1:A:420:LEU:HD13	1:A:435:MET:HE3	2.00	0.42
1:A:617:MET:HA	1:A:618[B]:PRO:HD3	1.95	0.42
1:A:862[A]:ARG:HD3	1:A:864:GLU:OE1	2.20	0.41
1:A:865[B]:GLY:O	1:A:867:GLU:N	2.53	0.41
1:A:235:SER:HB2	1:A:293:ASP:HB2	2.02	0.41
1:A:882:PRO:HB2	1:A:884:GLN:HE22	1.86	0.41
1:A:202:ASN:ND2	1:A:251:GLN:HE22	2.16	0.41
1:A:650:ASN:ND2	1:A:862[B]:ARG:NH2	2.69	0.41
1:A:177:ARG:NE	1:A:177:ARG:HA	2.36	0.41
1:A:529:LEU:C	1:A:529:LEU:CD2	2.87	0.41
1:A:368:GLN:HE21	1:A:368:GLN:HB3	1.71	0.40
1:A:664:LEU:C	1:A:664:LEU:HD23	2.42	0.40
1:A:865[B]:GLY:C	1:A:867:GLU:N	2.73	0.40
1:A:808[A]:GLN:HG3	4:A:2485:HOH:O	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:263[A]:GLU:HG3	1:A:263[A]:GLU:H	1.72	0.40
1:A:640:PHE:CD1	1:A:875:PRO:HG2	2.56	0.40
1:A:404:TYR:CD1	1:A:407:ALA:HB3	2.57	0.40
1:A:681:GLY:O	1:A:792:SER:HB2	2.21	0.40
1:A:375:ARG:HD3	4:A:2166:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	743/731 (102%)	714 (96%)	26 (4%)	3 (0%)	43 21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	866	ASP
1	A	865[A]	GLY
1	A	865[B]	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	663/650 (102%)	648 (98%)	15 (2%)	63 36

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	218	SER
1	A	219	GLN
1	A	254	ASN
1	A	282	ASN
1	A	327	LYS
1	A	368	GLN
1	A	404	TYR
1	A	433[A]	ASP
1	A	433[B]	ASP
1	A	626	LYS
1	A	661	ASN
1	A	705	ASN
1	A	773	LYS
1	A	822	GLU
1	A	874	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	231	ASN
1	A	237	ASN
1	A	254	ASN
1	A	261	GLN
1	A	277	HIS
1	A	282	ASN
1	A	341	ASN
1	A	368	GLN
1	A	386	GLN
1	A	392	GLN
1	A	418	GLN
1	A	650	ASN
1	A	661	ASN
1	A	667	HIS
1	A	683	ASN
1	A	698	GLN
1	A	759	GLN
1	A	788	GLN
1	A	820	ASN
1	A	825	GLN
1	A	874	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GCD	A	900	2	10,11,12	5.09	3 (30%)	11,15,17	3.75	9 (81%)
2	ASG	A	901	2	19,19,19	2.64	7 (36%)	28,28,28	1.22	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GCD	A	900	2	-	0/4/17/20	0/1/1/1
2	ASG	A	901	2	-	0/11/31/31	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	GCD	O5-C5	15.12	1.54	1.37
2	A	901	ASG	O4-S	-6.60	1.49	1.60
2	A	901	ASG	O5-C1	4.76	1.52	1.43
2	A	901	ASG	O4-C4	-4.12	1.39	1.46
2	A	901	ASG	O5-C5	4.11	1.54	1.44
2	A	900	GCD	C3-C4	3.98	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	ASG	C2-N	3.59	1.52	1.45
2	A	901	ASG	C3-C4	2.54	1.59	1.52
2	A	900	GCD	C4-C5	2.45	1.38	1.33
2	A	901	ASG	C7-N	2.11	1.42	1.34

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	900	GCD	O3-C3-C2	-7.35	101.33	111.03
2	A	900	GCD	C2-C3-C4	4.47	119.91	111.68
2	A	900	GCD	O3-C3-C4	4.47	120.28	109.67
2	A	900	GCD	O6A-C6-C5	-3.87	111.04	120.19
2	A	900	GCD	O5-C5-C4	-3.49	118.51	122.91
2	A	901	ASG	O4-C4-C5	3.31	115.62	108.59
2	A	900	GCD	O6B-C6-O6A	3.24	131.29	123.62
2	A	900	GCD	O5-C5-C6	2.77	116.12	111.75
2	A	900	GCD	C3-C4-C5	-2.64	117.35	121.82
2	A	900	GCD	O2-C2-C3	-2.57	107.63	111.03
2	A	901	ASG	C8-C7-N	-2.27	111.66	116.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1200	-	4,4,4	0.31	0	6,6,6	0.07	0
3	SO4	A	1201	-	4,4,4	0.28	0	6,6,6	0.08	0
3	SO4	A	1202	-	4,4,4	0.39	0	6,6,6	0.08	0
3	SO4	A	1203	-	4,4,4	0.40	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	1200	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1201	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1202	-	-	0/0/0/0	0/0/0/0
3	SO4	A	1203	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	722/731 (98%)	0.16	37 (5%)	27 34	12, 23, 41, 58	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	VAL	8.8
1	A	221	ASP	6.1
1	A	427	LYS	4.8
1	A	224	TYR	4.8
1	A	171	LYS	4.3
1	A	223	ILE	4.2
1	A	891	GLU	4.2
1	A	340[A]	ASP	4.2
1	A	233	LYS	4.1
1	A	279	HIS	4.0
1	A	375	ARG	4.0
1	A	865[A]	GLY	3.8
1	A	232	TYR	3.4
1	A	866	ASP	3.4
1	A	316[A]	GLU	3.3
1	A	341	ASN	3.2
1	A	230	SER	2.9
1	A	338	THR	2.7
1	A	433[A]	ASP	2.7
1	A	890	LEU	2.6
1	A	174	TYR	2.5
1	A	849	ASN	2.5
1	A	689	THR	2.4
1	A	495	GLN	2.3
1	A	218	SER	2.3
1	A	867	GLU	2.3
1	A	228[A]	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	431	ASP	2.2
1	A	773	LYS	2.2
1	A	428	ASN	2.2
1	A	196	ASP	2.2
1	A	227	GLU	2.1
1	A	704[A]	SER	2.1
1	A	411	VAL	2.1
1	A	234	THR	2.1
1	A	721	GLN	2.1
1	A	219	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ASG	A	901	19/19	0.11	-0.17	21,27,41,43	0
2	GCD	A	900	11/12	0.10	-0.89	24,30,33,33	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	A	1201	5/5	0.12	9.61	34,34,44,45	0
3	SO4	A	1203	5/5	0.20	2.61	58,60,62,63	0
3	SO4	A	1200	5/5	0.16	0.99	34,39,45,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	1202	5/5	0.13	0.19	53,54,55,55	0

6.5 Other polymers ⓘ

There are no such residues in this entry.