



Full wwPDB X-ray Structure Validation Report

May 8, 2014 – 01:32 AM EDT

PDB ID : 4OJD
Title : Crystal structure of a C-terminally truncated trimeric ectodomain of the C. elegans fusion protein EFF-1 G260A/D321E/D322E mutant
Authors : Krey, T.; Rey, F.A.
Deposited on : 2014-01-21
Resolution : 2.26 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable22978
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22978

1 Overall quality at a glance


The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	H	526	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NAG	H	1104	-	X
3	NAG	H	1105	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3684 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EFF-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	444	Total	C	N	O	S	0	1	0
			3528	2213	625	664	26			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	21	ARG	-	EXPRESSION TAG	UNP G5ECA1
H	22	SER	-	EXPRESSION TAG	UNP G5ECA1
H	260	ALA	GLY	ENGINEERED MUTATION	UNP G5ECA1
H	321	GLU	ASP	ENGINEERED MUTATION	UNP G5ECA1
H	322	GLU	ASP	ENGINEERED MUTATION	UNP G5ECA1
H	510	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	511	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	512	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	513	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	514	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	515	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	516	LYS	-	EXPRESSION TAG	UNP G5ECA1
H	517	ALA	-	EXPRESSION TAG	UNP G5ECA1
H	518	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	519	TRP	-	EXPRESSION TAG	UNP G5ECA1
H	520	SER	-	EXPRESSION TAG	UNP G5ECA1
H	521	HIS	-	EXPRESSION TAG	UNP G5ECA1
H	522	PRO	-	EXPRESSION TAG	UNP G5ECA1
H	523	GLN	-	EXPRESSION TAG	UNP G5ECA1
H	524	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	525	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	526	LYS	-	EXPRESSION TAG	UNP G5ECA1
H	527	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	528	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	529	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	530	SER	-	EXPRESSION TAG	UNP G5ECA1
H	531	GLY	-	EXPRESSION TAG	UNP G5ECA1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	532	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	533	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	534	SER	-	EXPRESSION TAG	UNP G5ECA1
H	535	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	536	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	537	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	538	SER	-	EXPRESSION TAG	UNP G5ECA1
H	539	TRP	-	EXPRESSION TAG	UNP G5ECA1
H	540	SER	-	EXPRESSION TAG	UNP G5ECA1
H	541	HIS	-	EXPRESSION TAG	UNP G5ECA1
H	542	PRO	-	EXPRESSION TAG	UNP G5ECA1
H	543	GLN	-	EXPRESSION TAG	UNP G5ECA1
H	544	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	545	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	546	LYS	-	EXPRESSION TAG	UNP G5ECA1

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	H	3	Total	C	N	O	0	0
			39	22	2	15		

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
H	21	ARG	-	EXPRESSION TAG	UNP G5ECA1
H	22	SER	-	EXPRESSION TAG	UNP G5ECA1
H	260	ALA	GLY	ENGINEERED MUTATION	UNP G5ECA1
H	321	GLU	ASP	ENGINEERED MUTATION	UNP G5ECA1
H	322	GLU	ASP	ENGINEERED MUTATION	UNP G5ECA1
H	510	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	511	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	512	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	513	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	514	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	515	ASP	-	EXPRESSION TAG	UNP G5ECA1
H	516	LYS	-	EXPRESSION TAG	UNP G5ECA1
H	517	ALA	-	EXPRESSION TAG	UNP G5ECA1
H	518	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	519	TRP	-	EXPRESSION TAG	UNP G5ECA1
H	520	SER	-	EXPRESSION TAG	UNP G5ECA1
H	521	HIS	-	EXPRESSION TAG	UNP G5ECA1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	522	PRO	-	EXPRESSION TAG	UNP G5ECA1
H	523	GLN	-	EXPRESSION TAG	UNP G5ECA1
H	524	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	525	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	526	LYS	-	EXPRESSION TAG	UNP G5ECA1
H	527	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	528	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	529	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	530	SER	-	EXPRESSION TAG	UNP G5ECA1
H	531	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	532	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	533	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	534	SER	-	EXPRESSION TAG	UNP G5ECA1
H	535	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	536	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	537	GLY	-	EXPRESSION TAG	UNP G5ECA1
H	538	SER	-	EXPRESSION TAG	UNP G5ECA1
H	539	TRP	-	EXPRESSION TAG	UNP G5ECA1
H	540	SER	-	EXPRESSION TAG	UNP G5ECA1
H	541	HIS	-	EXPRESSION TAG	UNP G5ECA1
H	542	PRO	-	EXPRESSION TAG	UNP G5ECA1
H	543	GLN	-	EXPRESSION TAG	UNP G5ECA1
H	544	PHE	-	EXPRESSION TAG	UNP G5ECA1
H	545	GLU	-	EXPRESSION TAG	UNP G5ECA1
H	546	LYS	-	EXPRESSION TAG	UNP G5ECA1

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	H	1	Total	C	N	O	0	0
			14	8	1	5		
3	H	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	89	Total	O	0	0
			89	89		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EFF-1A

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	132.44Å 132.44Å 132.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.83 – 2.26 46.83 – 2.26	Depositor EDS
% Data completeness (in resolution range)	96.8 (46.83-2.26) 96.9 (46.83-2.26)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.198 , 0.222 0.200 , 0.227	Depositor DCC
R_{free} test set	1776 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.4	EDS
Estimated twinning fraction	0.037 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 35346 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3684	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.51	0/3610	0.74	0/4879

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	194	TYR	Mainchain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	3528	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	39	0	34	0	0
3	H	28	0	26	0	0
4	H	89	0	0	0	0
All	All	3684	0	60	1	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 0.

All (1) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:H:194:TYR:O	1:H:196:ASN:N	2.45	0.49

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	435/526 (83%)	417 (96%)	17 (4%)	1 (0%)	56 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	54	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	H	392/457 (86%)	378 (96%)	14 (4%)	47 54

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	34	ARG
1	H	52	GLN
1	H	55	MET
1	H	101	LEU
1	H	102	LEU
1	H	192	LYS
1	H	221	VAL
1	H	230	LYS
1	H	244	ARG
1	H	325	LYS
1	H	339	ILE
1	H	375	GLN
1	H	426	LEU
1	H	474	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	H	1101	1,2	12,14,15	0.40	0	15,19,21	0.57	0
2	NAG	H	1102	2	12,14,15	0.46	0	15,19,21	0.78	0
2	BMA	H	1103	2	10,11,12	0.40	0	11,15,17	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	H	1101	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	1102	2	-	0/6/23/26	0/1/1/1
2	BMA	H	1103	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	H	1104	1	12,14,15	0.35	0	15,19,21	0.40	0
3	NAG	H	1105	1	12,14,15	0.34	0	15,19,21	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	H	1104	1	-	0/6/23/26	0/1/1/1
3	NAG	H	1105	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	444/526 (84%)	0.02	18 (4%) 35 40	34, 45, 80, 134	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	55	MET	6.3
1	H	398	SER	5.6
1	H	277	ASN	4.5
1	H	397	GLU	4.3
1	H	510	PHE	3.4
1	H	495	ILE	3.1
1	H	35	ALA	2.9
1	H	194	TYR	2.7
1	H	278	GLY	2.7
1	H	276	SER	2.6
1	H	260	ALA	2.6
1	H	399	GLN	2.5
1	H	54	ALA	2.5
1	H	161	THR	2.5
1	H	195	LYS	2.5
1	H	261	ARG	2.3
1	H	159	ASP	2.2
1	H	418	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	H	1102	14/15	0.15	0.54	58,66,71,79	0
2	NAG	H	1101	14/15	0.16	0.39	54,57,62,65	0
2	BMA	H	1103	11/12	0.24	-	80,85,86,88	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	H	1105	14/15	0.47	6.71	104,107,112,112	0
3	NAG	H	1104	14/15	0.41	5.12	109,113,115,115	0

6.5 Other polymers

There are no such residues in this entry.