



wwPDB X-ray Structure Validation Summary Report i

Mar 31, 2014 – 03:13 PM BST

PDB ID : 3OQC
Title : Ubiquitin-fold modifier 1 Specific Protease, UfSP2
Authors : Ha, B.H.; Chung, C.H.; Kim, E.E.
Deposited on : 2010-09-02
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable23004
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable23004

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	481	
1	B	481	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6716 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ufm1-specific protease 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	0	0
			3328	2131	573	606	18			
1	B	411	Total	C	N	O	S	0	0	0
			3281	2101	561	601	18			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q99K23
A	-18	GLY	-	EXPRESSION TAG	UNP Q99K23
A	-17	SER	-	EXPRESSION TAG	UNP Q99K23
A	-16	SER	-	EXPRESSION TAG	UNP Q99K23
A	-15	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-14	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-13	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-12	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-11	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-10	HIS	-	EXPRESSION TAG	UNP Q99K23
A	-9	SER	-	EXPRESSION TAG	UNP Q99K23
A	-8	SER	-	EXPRESSION TAG	UNP Q99K23
A	-7	GLY	-	EXPRESSION TAG	UNP Q99K23
A	-6	LEU	-	EXPRESSION TAG	UNP Q99K23
A	-5	VAL	-	EXPRESSION TAG	UNP Q99K23
A	-4	PRO	-	EXPRESSION TAG	UNP Q99K23
A	-3	ARG	-	EXPRESSION TAG	UNP Q99K23
A	-2	GLY	-	EXPRESSION TAG	UNP Q99K23
A	-1	SER	-	EXPRESSION TAG	UNP Q99K23
A	0	HIS	-	EXPRESSION TAG	UNP Q99K23
A	94	ARG	LYS	ENGINEERED MUTATION	UNP Q99K23
A	128	ALA	ARG	ENGINEERED MUTATION	UNP Q99K23
A	294	SER	CYS	ENGINEERED MUTATION	UNP Q99K23
B	-19	MET	-	EXPRESSION TAG	UNP Q99K23
B	-18	GLY	-	EXPRESSION TAG	UNP Q99K23

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	SER	-	EXPRESSION TAG	UNP Q99K23
B	-16	SER	-	EXPRESSION TAG	UNP Q99K23
B	-15	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-14	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-13	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-12	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-11	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-10	HIS	-	EXPRESSION TAG	UNP Q99K23
B	-9	SER	-	EXPRESSION TAG	UNP Q99K23
B	-8	SER	-	EXPRESSION TAG	UNP Q99K23
B	-7	GLY	-	EXPRESSION TAG	UNP Q99K23
B	-6	LEU	-	EXPRESSION TAG	UNP Q99K23
B	-5	VAL	-	EXPRESSION TAG	UNP Q99K23
B	-4	PRO	-	EXPRESSION TAG	UNP Q99K23
B	-3	ARG	-	EXPRESSION TAG	UNP Q99K23
B	-2	GLY	-	EXPRESSION TAG	UNP Q99K23
B	-1	SER	-	EXPRESSION TAG	UNP Q99K23
B	0	HIS	-	EXPRESSION TAG	UNP Q99K23
B	94	ARG	LYS	ENGINEERED MUTATION	UNP Q99K23
B	128	ALA	ARG	ENGINEERED MUTATION	UNP Q99K23
B	294	SER	CYS	ENGINEERED MUTATION	UNP Q99K23

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	53	Total O 53 53	0	0
2	B	54	Total O 54 54	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ufm1-specific protease 2

Chain A: 



- Molecule 1: Ufm1-specific protease 2

Chain B: 



T312	E313	R314	T318	H319	A325	L326	A329	K332	F333	A334	T335	F336	S339	R340	Q341	L355	I356	G357	V358	T359	S360	K361	L363	F364	V365	Q367	G368	S369	E370	S373	Q374	V385	G386	T387	K390	V391	L396	I400	L401	N406	E407	T408	T409	G410	Q411
I416	L417	H420	Y421	T422	G423	A424	E425	D426	L431	G437	W438	K439	S440	P441	D442	F443	W444	N445	K446	D447	A448	Y449	L452	C453	L454	R457	P458	N459	A460	L461															

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	184.53Å 56.04Å 143.27Å 90.00° 128.01° 90.00°	Depositor
Resolution (Å)	46.10 – 2.60 46.10 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.4 (46.10-2.60) 98.4 (46.10-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.238 , 0.298 0.233 , 0.292	Depositor DCC
R_{free} test set	1776 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 23.1	EDS
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 35390 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6716	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	1/3409 (0.0%)	0.85	1/4627 (0.0%)
1	B	0.71	0/3361	0.84	4/4564 (0.1%)
All	All	0.74	1/6770 (0.0%)	0.84	5/9191 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	GLY	N-CA	6.91	1.56	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	8	ARG	NE-CZ-NH2	5.79	123.19	120.30
1	B	396	LEU	CA-CB-CG	5.44	127.80	115.30
1	B	8	ARG	NE-CZ-NH1	-5.38	117.61	120.30
1	B	201	LEU	CA-CB-CG	5.22	127.30	115.30
1	A	457	ARG	NE-CZ-NH1	-5.02	117.79	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	262	PRO	Peptide
1	A	57	TRP	Peptide
1	B	57	TRP	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3328	0	3297	167	1
1	B	3281	0	3243	126	0
2	A	53	0	0	15	0
2	B	54	0	0	10	1
All	All	6716	0	6540	293	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 22.

The worst 5 of 293 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:65:GLY:HA2	1:B:158:ARG:HH12	0.99	1.09
1:B:409:THR:HG21	2:B:494:HOH:O	1.50	1.08
1:B:260:LEU:HD11	1:B:437:GLY:HA2	1.37	1.07
1:A:216:GLN:HB2	2:A:476:HOH:O	1.58	1.03
1:B:362:ILE:HD11	1:B:364:PHE:CZ	1.94	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:181:LYS:O	2:B:463:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/481 (84%)	352 (87%)	45 (11%)	9 (2%)	10	18
1	B	401/481 (83%)	346 (86%)	45 (11%)	10 (2%)	9	14
All	All	807/962 (84%)	698 (86%)	90 (11%)	19 (2%)	9	16

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	58	PRO
1	A	73	CYS
1	A	182	GLY
1	B	208	SER
1	B	212	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/424 (86%)	291 (80%)	75 (20%)	2	3
1	B	360/424 (85%)	306 (85%)	54 (15%)	4	7
All	All	726/848 (86%)	597 (82%)	129 (18%)	2	4

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	363	LEU
1	A	452	LEU
1	B	390	MET
1	A	384	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	409	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	31	GLN
1	B	200	ASN
1	B	445	ASN
1	B	104	GLN
1	B	194	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/481 (86%)	-0.28	8 (1%) 64 61	5, 36, 81, 119	0
1	B	411/481 (85%)	-0.23	6 (1%) 70 71	5, 40, 83, 118	0
All	All	827/962 (85%)	-0.26	14 (1%) 67 66	5, 38, 82, 119	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	PRO	5.0
1	A	68	THR	4.6
1	B	115	ILE	4.4
1	B	135	TYR	3.7
1	A	59	ASN	3.3

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.