



Full wwPDB X-ray Structure Validation Report

Nov 11, 2014 – 05:25 PM EST

PDB ID : 4OQY
Title : Streptomyces sp. GF3546 imine reductase
Authors : Schneider, L.K.; Huber, T.; Gerhardt, S.; Mller, M.; Einsle, O.
Deposited on : 2014-02-10
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable24103
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.1.3
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable24103

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	290	
1	B	290	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NDP	A	1000	-	X

2 Entry composition ⓘ


There are 3 unique types of molecules in this entry. The entry contains 4264 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (S)-imine reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	0	0
			2006	1274	332	389	11			
1	B	274	Total	C	N	O	S	0	0	0
			2058	1310	339	398	11			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	77	Total	O	0	0
			77	77		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	75	Total	O	0	0
			75	75		

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (S)-imine reductase

Chain A: 



- Molecule 1: (S)-imine reductase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.32Å 68.24Å 181.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.53 – 1.90 54.53 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.6 (54.53-1.90) 96.6 (54.53-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.216 , 0.241 0.223 , 0.249	Depositor DCC
R_{free} test set	2198 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.802	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 31.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 43789 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4264	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2047	0.73	3/2785 (0.1%)
1	B	0.53	0/2101	0.69	1/2859 (0.0%)
All	All	0.52	0/4148	0.71	4/5644 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ARG	NE-CZ-NH1	6.91	123.76	120.30
1	A	158	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	A	161	ASP	CB-CG-OD1	5.32	123.08	118.30
1	B	158	ARG	NE-CZ-NH1	5.27	122.94	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2006	0	1991	18	0
1	B	2058	0	2046	17	0
2	A	48	0	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	77	0	0	2	0
3	B	75	0	0	0	0
All	All	4264	0	4063	30	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (30) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:191:GLN:OE1	1:B:286:LYS:HE2	1.73	0.88
1:A:191:GLN:OE1	1:B:286:LYS:CE	2.37	0.72
1:A:20:ASN:HD22	1:A:45:ARG:HH11	1.41	0.66
1:A:213:HIS:O	1:A:215:LEU:N	2.30	0.65
1:B:20:ASN:HD22	1:B:45:ARG:HH11	1.47	0.62
1:A:172:GLN:OE1	3:A:1164:HOH:O	2.17	0.58
1:B:232:ASP:OD2	1:B:277:SER:HB2	2.05	0.57
1:B:92:LEU:HD22	1:B:117:GLY:HA3	1.87	0.56
1:B:282:ILE:O	1:B:286:LYS:HG3	2.06	0.55
1:B:180:MET:HE3	1:B:184:TYR:HE2	1.72	0.55
1:B:4:GLN:O	1:B:27:HIS:HD2	1.90	0.55
1:A:4:GLN:O	1:A:27:HIS:HD2	1.91	0.53
1:B:5:SER:HB3	1:B:28:GLU:HB2	1.91	0.53
1:A:3:LYS:HB3	1:A:3:LYS:NZ	2.25	0.51
1:B:24:ASP:OD1	1:B:45:ARG:NH2	2.44	0.50
1:A:211:MET:HE3	1:A:215:LEU:HD11	1.94	0.50
1:A:282:ILE:O	1:A:286:LYS:HG3	2.12	0.50
1:A:248:HIS:HE1	1:B:174:THR:OG1	1.99	0.45
1:A:286:LYS:HD3	1:B:191:GLN:OE1	2.19	0.43
1:A:20:ASN:ND2	1:A:45:ARG:HH11	2.13	0.43
1:B:158:ARG:HD3	1:B:168:TYR:CZ	2.55	0.42
1:B:11:LEU:H	1:B:33:ASN:ND2	2.18	0.41
1:A:178:THR:HG22	1:B:178:THR:HG22	2.03	0.41
1:B:246:HIS:HD2	1:B:249:GLN:OE1	2.04	0.41
1:A:24:ASP:OD1	1:A:45:ARG:NH2	2.52	0.41
1:B:287:LYS:N	1:B:287:LYS:HD2	2.36	0.41
1:A:121:VAL:O	2:A:1000:NDP:H42N	2.21	0.41
1:A:214:PHE:CD1	1:A:214:PHE:O	2.74	0.40
1:A:116:THR:O	1:A:135:TYR:HA	2.22	0.40
1:A:114:HIS:HD2	3:A:1136:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/290 (91%)	258 (98%)	4 (2%)	2 (1%)	27	12
1	B	270/290 (93%)	267 (99%)	3 (1%)	0	100	100
All	All	534/580 (92%)	525 (98%)	7 (1%)	2 (0%)	43	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	214	PHE
1	A	213	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/229 (92%)	205 (97%)	6 (3%)	56	45
1	B	217/229 (95%)	214 (99%)	3 (1%)	78	75
All	All	428/458 (93%)	419 (98%)	9 (2%)	66	59

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	LEU
1	A	139	LYS
1	A	198	LYS
1	A	206	MET
1	A	210	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	277	SER
1	B	206	MET
1	B	217	LEU
1	B	277	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	25	ASN
1	A	27	HIS
1	A	60	ASN
1	A	114	HIS
1	A	248	HIS
1	B	20	ASN
1	B	25	ASN
1	B	27	HIS
1	B	33	ASN
1	B	60	ASN
1	B	170	GLN
1	B	213	HIS
1	B	243	HIS
1	B	246	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	1000	-	52,52,52	1.26	6 (11%)	80,80,80	1.73	16 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1000	-	-	0/35/77/77	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	NDP	C6N-C5N	3.73	1.41	1.33
2	A	1000	NDP	C5A-C4A	3.60	1.48	1.40
2	A	1000	NDP	C4A-N9A	-3.43	1.32	1.37
2	A	1000	NDP	C2N-C3N	2.45	1.39	1.34
2	A	1000	NDP	PA-O3	2.24	1.63	1.59
2	A	1000	NDP	C4N-C3N	2.10	1.54	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	NDP	N3A-C2A-N1A	-6.76	122.94	128.89
2	A	1000	NDP	C5A-C4A-N3A	-4.61	121.49	125.98
2	A	1000	NDP	N3A-C4A-N9A	4.36	132.88	125.39
2	A	1000	NDP	C1B-N9A-C4A	-4.01	119.70	126.64
2	A	1000	NDP	C8A-N9A-C4A	3.62	109.90	106.96
2	A	1000	NDP	C4B-O4B-C1B	-2.89	106.55	109.72
2	A	1000	NDP	C2A-N1A-C6A	2.62	123.42	118.76
2	A	1000	NDP	O2B-P2B-O1X	-2.46	99.75	106.80
2	A	1000	NDP	C1D-N1N-C2N	-2.40	116.88	121.03
2	A	1000	NDP	O3X-P2B-O2X	2.36	116.24	107.38
2	A	1000	NDP	O3X-P2B-O2B	-2.24	100.64	107.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	NDP	C8A-N9A-C1B	2.23	130.36	126.15
2	A	1000	NDP	O2D-C2D-C3D	2.20	118.94	111.83
2	A	1000	NDP	C4A-C5A-N7A	-2.17	107.31	109.41
2	A	1000	NDP	C3B-C2B-C1B	-2.13	98.60	102.70
2	A	1000	NDP	C3N-C7N-N7N	2.11	122.14	117.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/290 (92%)	0.12	6 (2%) 59 60	17, 25, 38, 60	0
1	B	274/290 (94%)	0.09	2 (0%) 84 87	18, 25, 37, 44	0
All	All	542/580 (93%)	0.10	8 (1%) 70 72	17, 25, 38, 60	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	214	PHE	4.2
1	B	231	VAL	2.9
1	A	213	HIS	2.8
1	B	130	GLU	2.3
1	A	210	MET	2.3
1	A	130	GLU	2.2
1	A	232	ASP	2.1
1	A	201	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NDP	A	1000	48/48	0.23	4.36	15,20,23,24	0

6.5 Other polymers ⓘ

There are no such residues in this entry.