



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 10:00 PM GMT

PDB ID : 1ORJ
Title : FLAGELLAR EXPORT CHAPERONE
Authors : Evdokimov, A.G.; Phan, J.; Tropea, J.E.; Routzahn, K.M.; Peters III, H.K.;
Pokross, M.; Waugh, D.S.
Deposited on : 2003-03-13
Resolution : 2.25 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	130	
1	B	130	
1	C	130	
1	D	130	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4130 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called flagellar protein FliS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	0	0
			1054	676	175	201	2			
1	B	105	Total	C	N	O	S	0	0	0
			879	567	141	170	1			
1	C	104	Total	C	N	O	S	0	0	0
			870	561	139	169	1			
1	D	125	Total	C	N	O	S	0	0	0
			1044	670	172	200	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1125	HIS	-	EXPRESSION TAG	UNP O67806
A	1126	HIS	-	EXPRESSION TAG	UNP O67806
A	1127	HIS	-	EXPRESSION TAG	UNP O67806
A	1128	HIS	-	EXPRESSION TAG	UNP O67806
A	1129	HIS	-	EXPRESSION TAG	UNP O67806
A	1130	HIS	-	EXPRESSION TAG	UNP O67806
B	2125	HIS	-	EXPRESSION TAG	UNP O67806
B	2126	HIS	-	EXPRESSION TAG	UNP O67806
B	2127	HIS	-	EXPRESSION TAG	UNP O67806
B	2128	HIS	-	EXPRESSION TAG	UNP O67806
B	2129	HIS	-	EXPRESSION TAG	UNP O67806
B	2130	HIS	-	EXPRESSION TAG	UNP O67806
C	3125	HIS	-	EXPRESSION TAG	UNP O67806
C	3126	HIS	-	EXPRESSION TAG	UNP O67806
C	3127	HIS	-	EXPRESSION TAG	UNP O67806
C	3128	HIS	-	EXPRESSION TAG	UNP O67806
C	3129	HIS	-	EXPRESSION TAG	UNP O67806
C	3130	HIS	-	EXPRESSION TAG	UNP O67806
D	4125	HIS	-	EXPRESSION TAG	UNP O67806
D	4126	HIS	-	EXPRESSION TAG	UNP O67806
D	4127	HIS	-	EXPRESSION TAG	UNP O67806

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	4128	HIS	-	EXPRESSION TAG	UNP O67806
D	4129	HIS	-	EXPRESSION TAG	UNP O67806
D	4130	HIS	-	EXPRESSION TAG	UNP O67806

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	77	Total 77	O 77	0	0
2	B	69	Total 69	O 69	0	0
2	C	56	Total 56	O 56	0	0
2	D	81	Total 81	O 81	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: flagellar protein FliS

Chain A: 




- Molecule 1: flagellar protein FliS

Chain B: 



- Molecule 1: flagellar protein FliS


Chain C: 



- Molecule 1: flagellar protein FliS

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.07Å 131.76Å 74.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.00 – 2.25 43.01 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.2 (43.00-2.25) 95.0 (43.01-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.1.19, SHELXL	Depositor
R, R_{free}	0.189 , 0.247 0.204 , 0.222	Depositor DCC
R_{free} test set	2120 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.634	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 85.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 42213 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4130	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1068	0.85	0/1437
1	B	0.35	0/888	0.99	4/1192 (0.3%)
1	C	0.36	0/879	0.92	2/1181 (0.2%)
1	D	0.33	0/1057	0.88	2/1422 (0.1%)
All	All	0.35	0/3892	0.91	8/5232 (0.2%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2049	ARG	NE-CZ-NH2	-11.50	114.55	120.30
1	B	2049	ARG	CD-NE-CZ	7.31	133.83	123.60
1	B	2049	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	C	3074	LYS	C-N-CA	-5.86	109.99	122.30
1	D	4040	TYR	CB-CG-CD1	5.75	124.45	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1054	0	1084	27	0
1	B	879	0	921	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	870	0	908	22	1
1	D	1044	0	1077	36	0
2	A	77	0	0	4	0
2	B	69	0	0	3	1
2	C	56	0	0	3	0
2	D	81	0	0	9	0
All	All	4130	0	3990	101	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1114:ARG:HD3	1:A:1118:GLU:OE2	1.80	0.81
1:D:4016:ALA:O	1:D:4020:GLU:HG3	1.85	0.78
1:B:2049:ARG:HD3	2:B:5059:HOH:O	1.84	0.76
1:D:4044:ASN:H	1:D:4044:ASN:HD22	1.35	0.75
1:A:1109:ILE:O	1:A:1112:ASP:OD1	2.04	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:3077:GLU:OE2	2:B:5174:HOH:O[2_554]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/130 (95%)	118 (95%)	3 (2%)	3 (2%)	9	4
1	B	103/130 (79%)	100 (97%)	3 (3%)	0	100	100
1	C	102/130 (78%)	99 (97%)	3 (3%)	0	100	100
1	D	123/130 (95%)	119 (97%)	3 (2%)	1 (1%)	27	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	452/520 (87%)	436 (96%)	12 (3%)	4 (1%)	25	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1003	ASN
1	A	1011	ASN
1	A	1045	GLU
1	D	4003	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/121 (97%)	101 (86%)	16 (14%)	5	3
1	B	99/121 (82%)	90 (91%)	9 (9%)	14	10
1	C	98/121 (81%)	88 (90%)	10 (10%)	11	7
1	D	116/121 (96%)	105 (90%)	11 (10%)	12	9
All	All	430/484 (89%)	384 (89%)	46 (11%)	10	6

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2097	ASP
1	C	3019	LEU
1	D	4114	ARG
1	B	2101	GLU
1	B	2121	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	4010	GLN
1	D	4081	ASN
1	D	4044	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1072	HIS
1	D	4072	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/130 (96%)	-0.05	5 (3%) 36 42	20, 42, 77, 117	0
1	B	105/130 (80%)	-0.28	2 (1%) 64 70	22, 38, 72, 108	0
1	C	104/130 (80%)	-0.30	1 (0%) 79 84	20, 41, 73, 89	0
1	D	125/130 (96%)	-0.16	2 (1%) 68 75	21, 38, 63, 110	0
All	All	460/520 (88%)	-0.19	10 (2%) 59 65	20, 40, 73, 117	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1127	HIS	3.5
1	A	1124	VAL	3.0
1	B	2121	LYS	3.0
1	C	3120	VAL	2.9
1	A	1126	HIS	2.7

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.