



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 11:23 AM GMT

PDB ID : 2OSY  
Title : Endo-glycoceramidase II from Rhodococcus sp.: Lactosyl-Enzyme Intermediate  
Authors : Caines, M.E.C.; Strynadka, N.C.J.  
Deposited on : 2007-02-06  
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	481	
1	B	481	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7041 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoglycoceramidase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	432	Total	C	N	O	S	37	1	0
			3336	2110	572	643	11			
1	B	435	Total	C	N	O	S	49	1	0
			3358	2125	575	646	12			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	CLONING ARTIFACT	UNP O33853
A	11	GLY	-	CLONING ARTIFACT	UNP O33853
A	12	SER	-	CLONING ARTIFACT	UNP O33853
A	13	SER	-	CLONING ARTIFACT	UNP O33853
A	14	HIS	-	EXPRESSION TAG	UNP O33853
A	15	HIS	-	EXPRESSION TAG	UNP O33853
A	16	HIS	-	EXPRESSION TAG	UNP O33853
A	17	HIS	-	EXPRESSION TAG	UNP O33853
A	18	HIS	-	EXPRESSION TAG	UNP O33853
A	19	HIS	-	EXPRESSION TAG	UNP O33853
A	20	SER	-	CLONING ARTIFACT	UNP O33853
A	21	SER	-	CLONING ARTIFACT	UNP O33853
A	22	GLY	-	CLONING ARTIFACT	UNP O33853
A	23	LEU	-	CLONING ARTIFACT	UNP O33853
A	24	VAL	-	CLONING ARTIFACT	UNP O33853
A	25	PRO	-	CLONING ARTIFACT	UNP O33853
A	26	ARG	-	CLONING ARTIFACT	UNP O33853
A	27	GLY	-	CLONING ARTIFACT	UNP O33853
A	28	SER	-	CLONING ARTIFACT	UNP O33853
A	29	HIS	-	CLONING ARTIFACT	UNP O33853
A	30	MET	-	CLONING ARTIFACT	UNP O33853
A	233	ALA	GLU	ENGINEERED	UNP O33853
B	10	MET	-	CLONING ARTIFACT	UNP O33853
B	11	GLY	-	CLONING ARTIFACT	UNP O33853
B	12	SER	-	CLONING ARTIFACT	UNP O33853

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	13	SER	-	CLONING ARTIFACT	UNP O33853
B	14	HIS	-	EXPRESSION TAG	UNP O33853
B	15	HIS	-	EXPRESSION TAG	UNP O33853
B	16	HIS	-	EXPRESSION TAG	UNP O33853
B	17	HIS	-	EXPRESSION TAG	UNP O33853
B	18	HIS	-	EXPRESSION TAG	UNP O33853
B	19	HIS	-	EXPRESSION TAG	UNP O33853
B	20	SER	-	CLONING ARTIFACT	UNP O33853
B	21	SER	-	CLONING ARTIFACT	UNP O33853
B	22	GLY	-	CLONING ARTIFACT	UNP O33853
B	23	LEU	-	CLONING ARTIFACT	UNP O33853
B	24	VAL	-	CLONING ARTIFACT	UNP O33853
B	25	PRO	-	CLONING ARTIFACT	UNP O33853
B	26	ARG	-	CLONING ARTIFACT	UNP O33853
B	27	GLY	-	CLONING ARTIFACT	UNP O33853
B	28	SER	-	CLONING ARTIFACT	UNP O33853
B	29	HIS	-	CLONING ARTIFACT	UNP O33853
B	30	MET	-	CLONING ARTIFACT	UNP O33853
B	233	ALA	GLU	ENGINEERED	UNP O33853

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	2	Total	C	O	0	0
			22	12	10		
2	B	2	Total	C	O	0	0
			22	12	10		

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	CLONING ARTIFACT	UNP O33853
A	11	GLY	-	CLONING ARTIFACT	UNP O33853
A	12	SER	-	CLONING ARTIFACT	UNP O33853
A	13	SER	-	CLONING ARTIFACT	UNP O33853
A	14	HIS	-	EXPRESSION TAG	UNP O33853
A	15	HIS	-	EXPRESSION TAG	UNP O33853
A	16	HIS	-	EXPRESSION TAG	UNP O33853
A	17	HIS	-	EXPRESSION TAG	UNP O33853
A	18	HIS	-	EXPRESSION TAG	UNP O33853
A	19	HIS	-	EXPRESSION TAG	UNP O33853
A	20	SER	-	CLONING ARTIFACT	UNP O33853

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	SER	-	CLONING ARTIFACT	UNP O33853
A	22	GLY	-	CLONING ARTIFACT	UNP O33853
A	23	LEU	-	CLONING ARTIFACT	UNP O33853
A	24	VAL	-	CLONING ARTIFACT	UNP O33853
A	25	PRO	-	CLONING ARTIFACT	UNP O33853
A	26	ARG	-	CLONING ARTIFACT	UNP O33853
A	27	GLY	-	CLONING ARTIFACT	UNP O33853
A	28	SER	-	CLONING ARTIFACT	UNP O33853
A	29	HIS	-	CLONING ARTIFACT	UNP O33853
A	30	MET	-	CLONING ARTIFACT	UNP O33853
A	233	ALA	GLU	ENGINEERED	UNP O33853
B	10	MET	-	CLONING ARTIFACT	UNP O33853
B	11	GLY	-	CLONING ARTIFACT	UNP O33853
B	12	SER	-	CLONING ARTIFACT	UNP O33853
B	13	SER	-	CLONING ARTIFACT	UNP O33853
B	14	HIS	-	EXPRESSION TAG	UNP O33853
B	15	HIS	-	EXPRESSION TAG	UNP O33853
B	16	HIS	-	EXPRESSION TAG	UNP O33853
B	17	HIS	-	EXPRESSION TAG	UNP O33853
B	18	HIS	-	EXPRESSION TAG	UNP O33853
B	19	HIS	-	EXPRESSION TAG	UNP O33853
B	20	SER	-	CLONING ARTIFACT	UNP O33853
B	21	SER	-	CLONING ARTIFACT	UNP O33853
B	22	GLY	-	CLONING ARTIFACT	UNP O33853
B	23	LEU	-	CLONING ARTIFACT	UNP O33853
B	24	VAL	-	CLONING ARTIFACT	UNP O33853
B	25	PRO	-	CLONING ARTIFACT	UNP O33853
B	26	ARG	-	CLONING ARTIFACT	UNP O33853
B	27	GLY	-	CLONING ARTIFACT	UNP O33853
B	28	SER	-	CLONING ARTIFACT	UNP O33853
B	29	HIS	-	CLONING ARTIFACT	UNP O33853
B	30	MET	-	CLONING ARTIFACT	UNP O33853
B	233	ALA	GLU	ENGINEERED	UNP O33853

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	A	1	Total Na 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	168	Total 168	O 168	0	0
4	B	133	Total 133	O 133	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.95Å 93.69Å 94.46Å 90.00° 98.12° 90.00°	Depositor
Resolution (Å)	18.73 – 2.10 18.72 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (18.73-2.10) 99.4 (18.72-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.97 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.205 , 0.251 0.210 , 0.257	Depositor DCC
$R_{free}$ test set	2757 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtriage
Anisotropy	0.199	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 29.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 54001 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7041	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GLC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	2/3433 (0.1%)	0.74	3/4700 (0.1%)
1	B	0.72	5/3456 (0.1%)	0.77	4/4733 (0.1%)
All	All	0.71	7/6889 (0.1%)	0.75	7/9433 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	146	GLU	CB-CG	-10.63	1.31	1.52
1	A	255	ARG	CB-CG	-9.29	1.27	1.52
1	B	206	GLU	CB-CG	-7.97	1.37	1.52
1	B	296	GLN	CB-CG	-7.16	1.33	1.52
1	B	177	ARG	CB-CG	7.15	1.71	1.52
1	A	419	GLU	CB-CG	5.90	1.63	1.52
1	B	210	LYS	CB-CG	-5.15	1.38	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	453	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	255	ARG	CA-CB-CG	6.41	127.50	113.40
1	B	400	LEU	CA-CB-CG	5.73	128.47	115.30
1	A	223	ASP	CB-CG-OD1	-5.62	113.24	118.30
1	A	223	ASP	CB-CG-OD2	5.37	123.13	118.30
1	B	114	ASP	CB-CG-OD2	-5.36	113.48	118.30
1	B	114	ASP	CB-CG-OD1	5.20	122.98	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3336	0	3180	12	0
1	B	3358	0	3205	27	0
2	A	22	0	19	3	0
2	B	22	0	19	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	168	0	0	0	0
4	B	133	0	0	0	0
All	All	7041	0	6423	39	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (39) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:270:CYS:HB3	1:B:302[B]:CYS:SG	2.04	0.98
1:B:351:GLU:CD	2:B:600:GLC:C1	2.52	0.77
1:A:351:GLU:CD	2:A:600:GLC:C1	2.52	0.77
1:B:277:GLY:O	1:B:282:LEU:HD13	1.98	0.64
1:B:417:PRO:HB3	1:B:429:LEU:HD11	1.83	0.59
1:B:306:TYR:CE2	1:B:351:GLU:OE1	2.55	0.59
1:B:434:ASP:HB3	1:B:437:ILE:HD12	1.86	0.58
1:B:322:THR:O	1:B:326:VAL:HG13	2.05	0.57
1:B:351:GLU:OE2	2:B:600:GLC:C2	2.50	0.55
1:B:270:CYS:CB	1:B:302[B]:CYS:SG	2.89	0.55
1:B:373:ARG:HD2	1:B:420:TRP:O	2.08	0.54
1:B:351:GLU:OE2	2:B:600:GLC:O5	2.25	0.52
1:A:351:GLU:OE2	2:A:600:GLC:C2	2.55	0.51
1:B:337:HIS:O	1:B:341:VAL:HG23	2.11	0.50
1:B:351:GLU:OE1	2:B:600:GLC:C1	2.59	0.50
1:B:131:MET:CE	1:B:270:CYS:SG	3.01	0.48
1:B:131:MET:HE3	1:B:227:ALA:CB	2.44	0.47
1:B:55:ILE:HB	1:B:410:PRO:O	2.14	0.47
1:A:139:TYR:HB2	1:A:191:PHE:CE2	2.50	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:281:GLY:O	1:A:334:ASN:ND2	2.48	0.47
1:A:352:PHE:HE1	1:A:364:TYR:CE2	2.34	0.46
1:A:58:GLY:HA3	1:A:91:PHE:O	2.16	0.46
1:B:356:THR:HB	1:B:400:LEU:HD22	2.00	0.44
1:B:124:ALA:HB2	1:B:224:ALA:HB2	1.99	0.44
1:A:304:HIS:CG	1:A:351:GLU:HB3	2.52	0.44
1:A:55:ILE:HB	1:A:410:PRO:O	2.17	0.44
1:B:306:TYR:HE2	1:B:351:GLU:OE1	1.97	0.44
1:A:352:PHE:CE1	1:A:368:VAL:HG21	2.54	0.42
1:B:282:LEU:HD12	1:B:282:LEU:N	2.34	0.42
1:B:92:VAL:CG1	1:B:130:VAL:HG22	2.50	0.42
1:B:259:ALA:O	1:B:262:GLN:HB2	2.20	0.41
1:B:304:HIS:CG	1:B:351:GLU:HB3	2.55	0.41
1:A:307:PRO:HB3	1:A:327:THR:HG21	2.03	0.41
1:A:351:GLU:OE2	2:A:600:GLC:O5	2.32	0.41
1:A:354:LEU:HD13	1:A:364:TYR:CD2	2.56	0.41
1:B:459:ALA:HB2	1:B:475:THR:HG22	2.02	0.41
1:B:355:ASP:OD1	1:B:388:PRO:HB2	2.20	0.41
1:B:131:MET:HE2	1:B:270:CYS:SG	2.61	0.41
1:B:216:ALA:O	1:B:220:ALA:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/481 (89%)	417 (98%)	10 (2%)	0	100	100
1	B	430/481 (89%)	417 (97%)	13 (3%)	0	100	100
All	All	857/962 (89%)	834 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/379 (90%)	339 (99%)	4 (1%)	82	87
1	B	346/379 (91%)	344 (99%)	2 (1%)	92	95
All	All	689/758 (91%)	683 (99%)	6 (1%)	87	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	ASP
1	A	352	PHE
1	A	354	LEU
1	A	400	LEU
1	B	326	VAL
1	B	352	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	A	600	1,2	10,11,12	0.72	0	11,15,17	1.06	0
2	GAL	A	601	2	10,11,12	0.95	1 (10%)	11,15,17	1.33	1 (9%)
2	GLC	B	600	1,2	10,11,12	0.64	0	11,15,17	0.93	1 (9%)
2	GAL	B	601	2	10,11,12	0.68	0	11,15,17	1.06	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	600	1,2	-	0/2/19/22	0/1/1/1
2	GAL	A	601	2	-	0/2/19/22	0/1/1/1
2	GLC	B	600	1,2	-	0/2/19/22	0/1/1/1
2	GAL	B	601	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	GAL	O5-C5	-2.72	1.40	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GAL	O5-C5-C6	2.77	109.89	106.98
2	B	601	GAL	O5-C5-C6	2.55	109.66	106.98
2	B	600	GLC	O5-C5-C6	2.03	109.11	106.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	432/481 (89%)	-0.05	7 (1%) 68 72	17, 26, 36, 50	10 (2%)
1	B	435/481 (90%)	-0.02	8 (1%) 65 69	18, 27, 37, 49	12 (2%)
All	All	867/962 (90%)	-0.04	15 (1%) 67 71	17, 26, 36, 50	22 (2%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	490	ALA	5.6
1	A	319	LEU	4.9
1	A	320	ALA	4.2
1	A	145	PRO	3.6
1	B	490	ALA	3.4
1	B	294	ALA	3.3
1	A	318	GLY	3.3
1	B	308	LEU	3.2
1	B	295	GLY	2.6
1	B	318	GLY	2.6
1	A	306	TYR	2.6
1	B	396	GLY	2.5
1	A	221	ASP	2.4
1	B	479	SER	2.4
1	B	395	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GAL	B	601	11/12	0.10	-0.39	24,26,28,29	0
2	GLC	B	600	11/12	0.09	-0.46	22,23,26,28	0
2	GLC	A	600	11/12	0.07	-0.87	22,23,26,26	0
2	GAL	A	601	11/12	0.08	-1.52	19,22,24,25	0

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	B	602	1/1	0.05	-1.17	27,27,27,27	0
3	NA	A	602	1/1	0.06	-1.68	23,23,23,23	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.