



wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 06:27 PM GMT

PDB ID : 1OTY
Title : Native PNP +ALLO
Authors : Ealick, S.E.; Bennett, E.M.; Anand, R.; Secrist, J.A.; Parker, W.B.; Hassan, A.E.; Allan, P.W.; McPherson, D.T.; Sorscher, E.J.
Deposited on : 2003-03-23
Resolution : 2.50 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	
1	C	238	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	6MP	A	247	X	-
3	6MP	B	246	X	-
3	6MP	C	245	X	-

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 5598 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purine nucleoside phosphorylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	Total 1793	C 1132	N 307	O 339	S 15	0	0	0
1	B	237	Total 1793	C 1132	N 307	O 339	S 15	0	0	0
1	C	237	Total 1793	C 1132	N 307	O 339	S 15	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0
2	C	1	Total 5	O 4	P 1	0	0

- Molecule 3 is 6-METHYLPURINE (three-letter code: 6MP) (formula: C₆H₆N₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C N 10 6 4	0	0
3	B	1	Total C N 10 6 4	0	0
3	A	1	Total C N 10 6 4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	51	Total O 51 51	0	0
4	B	74	Total O 74 74	0	0
4	C	49	Total O 49 49	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.97Å 119.97Å 240.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.20 – 2.50 26.20 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (26.20-2.50) 100.0 (26.20-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.56 (at 2.50Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.203 , 0.236 0.204 , 0.237	Depositor DCC
R_{free} test set	3603 reflections (9.98%)	DCC
Wilson B-factor (Å ²)	31.3	Xtrriage
Anisotropy	0.309	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 28.7	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 36118 reflections	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5598	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, 6MP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1822	0.60	0/2457
1	B	0.37	0/1822	0.62	0/2457
1	C	0.33	0/1822	0.59	0/2457
All	All	0.35	0/5466	0.60	0/7371

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1793	0	1791	47	0
1	B	1793	0	1791	40	0
1	C	1793	0	1791	51	1
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	10	0	6	0	0
3	B	10	0	6	2	0
3	C	10	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	51	0	0	1	0
4	B	74	0	0	0	0
4	C	49	0	0	0	0
All	All	5598	0	5391	137	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:21:ASP:HB3	1:A:24:ARG:HD3	1.59	0.84
1:A:142:LYS:HE3	1:A:142:LYS:HA	1.59	0.84
1:A:210:GLU:HG2	1:A:211:GLN:H	1.44	0.80
1:C:94:VAL:HG12	1:C:171:GLU:HB2	1.65	0.79
1:B:101:ARG:HA	1:B:220:THR:HG21	1.66	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:107:MET:CE	1:C:107:MET:CE[12_555]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/238 (99%)	220 (94%)	13 (6%)	2 (1%)	25	42
1	B	235/238 (99%)	220 (94%)	13 (6%)	2 (1%)	25	42
1	C	235/238 (99%)	218 (93%)	17 (7%)	0	100	100
All	All	705/714 (99%)	658 (93%)	43 (6%)	4 (1%)	33	55

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	207	ARG
1	A	210	GLU
1	B	211	GLN
1	B	214	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	187/189 (99%)	183 (98%)	4 (2%)	66 88
1	B	187/189 (99%)	176 (94%)	11 (6%)	28 48
1	C	187/189 (99%)	180 (96%)	7 (4%)	45 72
All	All	561/567 (99%)	539 (96%)	22 (4%)	43 70

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	120	PHE
1	B	207	ARG
1	C	120	PHE
1	B	152	ASN
1	B	179	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	40	ASN
1	B	152	ASN
1	C	116	ASN
1	A	123	HIS
1	C	6	ASN

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	6MP	A	247	-	11,11,11	5.01	8 (72%)	12,15,15	3.18	5 (41%)
2	PO4	A	248	-	4,4,4	0.92	0	6,6,6	0.31	0
3	6MP	B	246	-	11,11,11	5.05	7 (63%)	12,15,15	3.23	5 (41%)
2	PO4	B	249	-	4,4,4	1.12	0	6,6,6	0.31	0
3	6MP	C	245	-	11,11,11	4.77	8 (72%)	12,15,15	3.19	5 (41%)
2	PO4	C	250	-	4,4,4	0.97	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6MP	A	247	-	-	0/0/0/0	0/0/2/2
2	PO4	A	248	-	-	0/0/0/0	0/0/0/0
3	6MP	B	246	-	-	0/0/0/0	0/0/2/2
2	PO4	B	249	-	-	0/0/0/0	0/0/0/0
3	6MP	C	245	-	-	0/0/0/0	0/0/2/2
2	PO4	C	250	-	-	0/0/0/0	0/0/0/0

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	246	6MP	C6-N1	9.74	1.41	1.32
3	B	246	6MP	C4-N9	9.67	1.45	1.36
3	A	247	6MP	C6-N1	9.64	1.41	1.32
3	A	247	6MP	C4-N9	9.47	1.44	1.36
3	C	245	6MP	C4-N9	9.36	1.44	1.36

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	246	6MP	C8-N7-C5	-7.67	103.06	108.52
3	A	247	6MP	C8-N7-C5	-7.50	103.18	108.52
3	C	245	6MP	C8-N7-C5	-7.48	103.20	108.52
3	B	246	6MP	C4-C5-N7	4.85	111.28	106.02
3	C	245	6MP	C4-C5-N7	4.69	111.10	106.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/238 (99%)	0.26	21 (8%) 10 9	13, 32, 73, 119	0
1	B	237/238 (99%)	0.24	15 (6%) 19 19	13, 26, 65, 133	0
1	C	237/238 (99%)	0.25	24 (10%) 7 7	17, 32, 66, 114	0
All	All	711/714 (99%)	0.25	60 (8%) 11 10	13, 29, 71, 133	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	212	THR	19.2
1	B	212	THR	16.9
1	B	209	HIS	9.7
1	B	211	GLN	8.7
1	B	213	THR	8.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	6MP	A	247	10/10	0.24	1.37	57,60,62,62	0
3	6MP	C	245	10/10	0.17	1.07	37,41,43,44	0
3	6MP	B	246	10/10	0.18	-0.02	55,56,58,61	0
2	PO4	C	250	5/5	0.13	-0.51	36,36,37,37	0
2	PO4	B	249	5/5	0.10	-0.88	27,28,29,29	0
2	PO4	A	248	5/5	0.12	-1.09	58,58,60,60	0

6.5 Other polymers

There are no such residues in this entry.