



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 05:38 AM GMT

PDB ID : 3PAW  
Title : Low resolution X-ray crystal structure of Yeast Rnr1p with dATP bound in the A-site  
Authors : Fairman, J.W.; Wijerathna, S.R.; Dealwis, C.G.  
Deposited on : 2010-10-19  
Resolution : 6.61 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 6.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1097 (9.50-3.50)
Clashscore	79885	1036 (9.50-3.52)
Ramachandran outliers	78287	1291 (9.50-3.50)
RSRZ outliers	66119	1096 (9.50-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	888	
1	B	888	
1	C	888	
1	D	888	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 14596 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ribonucleoside-diphosphatereductase large chain 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	739	Total 3649	C 2171	N 739	O 739	0	0	0
1	B	739	Total 3649	C 2171	N 739	O 739	0	0	0
1	C	739	Total 3649	C 2171	N 739	O 739	0	0	0
1	D	739	Total 3649	C 2171	N 739	O 739	0	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonucleoside-diphosphatereductase large chain 1






GLU	SER	LYS	VAL	GLU	VAL	PRO	GLU	VAL	PRO	ALA	PRO	THR	LYS	ASN	GLU	GLU	LYS	ALA	ALA	PRO	ILE	VAL	ASP	ASP	GLU	GLU	THR	GLU	PHE	ASP	ILE	TYR	ASN	SER	LYS	VAL	ILE	ALA	CYS	ALA	ILE	ASP	ASN	PRO	GLU	ALA	CYS	GLU	GLY
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

• Molecule 1: Ribonucleoside-diphosphatereductase large chain 1

Chain C: 



• Molecule 1: Ribonucleoside-diphosphatereductase large chain 1

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.51Å 166.51Å 381.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	192.45 – 6.61 36.56 – 6.61	Depositor EDS
% Data completeness (in resolution range)	87.8 (192.45-6.61) 88.3 (36.56-6.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 6.63Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.391 , 0.442 0.390 , 0.442	Depositor DCC
$R_{free}$ test set	469 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	225.0	Xtriage
Anisotropy	0.669	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 196.4	EDS
Estimated twinning fraction	0.437 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.20$	Xtriage
Outliers	0 of 9847 reflections	Xtriage
$F_o, F_c$ correlation	0.65	EDS
Total number of atoms	14596	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/3647	0.65	0/5075
1	B	0.52	0/3647	0.65	0/5075
1	C	0.53	0/3647	0.66	0/5075
1	D	0.53	0/3647	0.63	0/5075
All	All	0.52	0/14588	0.65	0/20300

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3649	0	1652	61	0
1	B	3649	0	1652	55	0
1	C	3649	0	1652	58	0
1	D	3649	0	1652	44	0
All	All	14596	0	6608	217	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:147:GLY:HA3	1:A:614:ILE:HA	1.42	0.99
1:D:147:GLY:HA2	1:D:614:ILE:HA	1.48	0.95
1:C:603:MET:CB	1:C:706:ILE:HA	1.99	0.92
1:A:147:GLY:CA	1:A:614:ILE:HA	2.02	0.90
1:D:147:GLY:CA	1:D:614:ILE:HA	2.07	0.85

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	735/888 (83%)	533 (72%)	157 (21%)	45 (6%)	2	37
1	B	735/888 (83%)	518 (70%)	180 (24%)	37 (5%)	3	42
1	C	735/888 (83%)	536 (73%)	154 (21%)	45 (6%)	2	37
1	D	735/888 (83%)	542 (74%)	148 (20%)	45 (6%)	2	37
All	All	2940/3552 (83%)	2129 (72%)	639 (22%)	172 (6%)	2	38

5 of 172 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	GLU
1	A	227	SER
1	A	288	GLN
1	A	461	LYS
1	A	486	ASP

### 5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	739/888 (83%)	3.95	536 (72%) <b>0</b> <b>3</b>	39, 81, 170, 183	0
1	B	739/888 (83%)	3.46	469 (63%) <b>0</b> <b>3</b>	52, 74, 175, 188	0
1	C	739/888 (83%)	3.70	468 (63%) <b>0</b> <b>3</b>	51, 73, 173, 194	0
1	D	739/888 (83%)	4.18	540 (73%) <b>0</b> <b>3</b>	58, 80, 175, 186	0
All	All	2956/3552 (83%)	3.82	2013 (68%) <b>0</b> <b>3</b>	39, 78, 173, 194	0

The worst 5 of 2013 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	218	CYS	25.3
1	C	433	GLU	22.4
1	D	443	CYS	20.8
1	C	215	MET	19.8
1	B	220	LEU	19.6

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.