



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 02:50 AM GMT

PDB ID : 1PDA
Title : STRUCTURE OF PORPHOBILINOGEN DEAMINASE REVEALS A FLEXIBLE MULTIDOMAIN POLYMERASE WITH A SINGLE CATALYTIC SITE
Authors : Louie, G.V.; Brownlie, P.D.; Lambert, R.; Cooper, J.B.; Blundell, T.L.; Wood, S.P.; Warren, M.J.; Woodcock, S.C.; Jordan, P.M.
Deposited on : 1992-11-17
Resolution : 1.76 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1304 (1.76-1.76)
Ramachandran outliers	78287	1288 (1.76-1.76)
Sidechain outliers	78261	1288 (1.76-1.76)
RSRZ outliers	66119	1135 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	313	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	ACY	A	315	X	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2527 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PORPHOBILINOGEN DEAMINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	2244	1395	412	428	9	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	241	ALA	GLY	CONFLICT	UNP P06983
A	261	GLY	ALA	CONFLICT	UNP P06983

- Molecule 2 is 3-[5-{[3-(2-CARBOXYETHYL)-4-(CARBOXYMETHYL)-5-METHYL-1H-PYRROL-2-YL]METHYL}-4-(CARBOXYMETHYL)-1H-PYRROL-3-YL]PROPANOIC ACID (three-letter code: DPM) (formula: C₂₀H₂₄N₂O₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	30	20	2	8	0	0

- Molecule 3 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	249	Total	O	0	0
			249	249		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

● Molecule 1: PORPHOBILINOGEN DEAMINASE

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.00Å 75.90Å 50.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.76 10.00 – 1.76	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.76) 99.3 (10.00-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 1.76Å)	Xtriage
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.188 , (Not available) 0.172 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 69.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 67778 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2527	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, DPM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	2/2273 (0.1%)	2.14	76/3079 (2.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	293	GLU	CD-OE1	-5.29	1.19	1.25
1	A	273	ARG	CZ-NH2	5.04	1.39	1.33

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	182	ARG	NE-CZ-NH2	-19.53	110.53	120.30
1	A	237	ARG	NE-CZ-NH1	16.50	128.55	120.30
1	A	11	ARG	NE-CZ-NH2	-15.95	112.33	120.30
1	A	182	ARG	NE-CZ-NH1	15.62	128.11	120.30
1	A	74	ARG	NE-CZ-NH1	14.97	127.78	120.30
1	A	277	ARG	NE-CZ-NH1	-13.58	113.51	120.30
1	A	88	GLU	OE1-CD-OE2	12.81	138.68	123.30
1	A	277	ARG	NE-CZ-NH2	12.40	126.50	120.30
1	A	131	ARG	NE-CZ-NH1	11.79	126.20	120.30
1	A	164	TYR	CB-CG-CD1	11.50	127.90	121.00
1	A	84	ASP	CB-CG-OD1	11.36	128.53	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	176	ARG	NE-CZ-NH2	-11.27	114.66	120.30
1	A	155	ARG	NE-CZ-NH2	10.28	125.44	120.30
1	A	164	TYR	CG-CD1-CE1	9.82	129.15	121.30
1	A	206	ARG	NE-CZ-NH1	-9.60	115.50	120.30
1	A	132	ARG	NE-CZ-NH1	-9.59	115.51	120.30
1	A	276	ARG	NE-CZ-NH2	9.44	125.02	120.30
1	A	251	GLU	OE1-CD-OE2	9.14	134.27	123.30
1	A	28	MET	CG-SD-CE	-8.78	86.14	100.20
1	A	160	ASP	CB-CG-OD1	8.76	126.19	118.30
1	A	28	MET	CA-CB-CG	-8.67	98.56	113.30
1	A	164	TYR	CZ-CE2-CD2	8.34	127.30	119.80
1	A	84	ASP	CB-CG-OD2	-8.26	110.86	118.30
1	A	103	ASP	CB-CG-OD1	8.15	125.63	118.30
1	A	74	ARG	NE-CZ-NH2	-7.92	116.34	120.30
1	A	106	ASP	CB-CG-OD1	7.35	124.91	118.30
1	A	276	ARG	NE-CZ-NH1	-7.19	116.70	120.30
1	A	235	ASN	OD1-CG-ND2	7.17	138.40	121.90
1	A	307	TYR	CA-CB-CG	7.16	127.00	113.40
1	A	297	ASN	CB-CG-OD1	-6.96	107.69	121.60
1	A	232	ARG	CD-NE-CZ	6.90	133.26	123.60
1	A	105	ARG	NE-CZ-NH2	6.77	123.68	120.30
1	A	59	LYS	N-CA-CB	6.72	122.69	110.60
1	A	88	GLU	CG-CD-OE1	-6.72	104.86	118.30
1	A	297	ASN	OD1-CG-ND2	6.71	137.34	121.90
1	A	114	ASP	CB-CG-OD1	6.67	124.30	118.30
1	A	223	GLU	OE1-CD-OE2	6.62	131.25	123.30
1	A	139	ARG	NE-CZ-NH1	-6.49	117.06	120.30
1	A	103	ASP	CB-CG-OD2	-6.47	112.48	118.30
1	A	211	ARG	CD-NE-CZ	6.36	132.50	123.60
1	A	237	ARG	NH1-CZ-NH2	-6.31	112.46	119.40
1	A	305	GLU	CA-CB-CG	6.29	127.23	113.40
1	A	91	GLN	N-CA-CB	-6.26	99.33	110.60
1	A	160	ASP	CB-CG-OD2	-6.15	112.76	118.30
1	A	234	MET	CA-CB-CG	6.15	123.75	113.30
1	A	184	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	A	106	ASP	CB-CG-OD2	-6.12	112.80	118.30
1	A	113	TYR	CB-CG-CD1	-6.08	117.35	121.00
1	A	213	ARG	NE-CZ-NH1	-6.08	117.26	120.30
1	A	149	ARG	NE-CZ-NH2	6.04	123.32	120.30
1	A	277	ARG	CD-NE-CZ	-6.04	115.14	123.60
1	A	132	ARG	CD-NE-CZ	5.89	131.85	123.60
1	A	155	ARG	NE-CZ-NH1	-5.83	117.38	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	NE-CZ-NH2	-5.83	117.39	120.30
1	A	88	GLU	O-C-N	5.79	131.97	122.70
1	A	139	ARG	NE-CZ-NH2	5.75	123.18	120.30
1	A	235	ASN	CB-CG-OD1	-5.74	110.13	121.60
1	A	154	THR	CA-CB-OG1	-5.65	97.13	109.00
1	A	184	ARG	CG-CD-NE	-5.64	99.95	111.80
1	A	289	SER	N-CA-CB	5.62	118.93	110.50
1	A	7	ARG	NE-CZ-NH2	5.61	123.11	120.30
1	A	176	ARG	NH1-CZ-NH2	5.60	125.56	119.40
1	A	276	ARG	CD-NE-CZ	-5.55	115.83	123.60
1	A	95	LEU	CB-CG-CD1	-5.54	101.58	111.00
1	A	11	ARG	NH1-CZ-NH2	5.51	125.46	119.40
1	A	202	GLY	O-C-N	5.43	131.39	122.70
1	A	76	ASP	O-C-N	5.43	131.38	122.70
1	A	262	LEU	CA-CB-CG	5.38	127.68	115.30
1	A	89	PHE	CB-CG-CD1	5.36	124.56	120.80
1	A	286	MET	CA-CB-CG	5.36	122.42	113.30
1	A	293	GLU	O-C-N	-5.33	114.17	122.70
1	A	268	GLY	O-C-N	5.24	131.08	122.70
1	A	142	ASP	CB-CG-OD1	5.07	122.86	118.30
1	A	146	ARG	N-CA-CB	-5.03	101.55	110.60
1	A	130	LEU	O-C-N	5.02	130.74	122.70
1	A	131	ARG	NH1-CZ-NH2	-5.01	113.89	119.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	245	PRO	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2244	0	2267	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	30	0	18	0	0
3	A	4	0	3	0	0
4	A	249	0	0	6	1
All	All	2527	0	2288	28	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (28) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:180:GLU:HB2	4:A:557:HOH:O	1.75	0.85
1:A:196:VAL:HG21	1:A:235:ASN:ND2	1.91	0.84
1:A:90:PRO:HG2	1:A:93:LEU:HD12	1.72	0.70
1:A:196:VAL:CG2	1:A:235:ASN:ND2	2.54	0.70
1:A:196:VAL:HG22	1:A:248:SER:HB3	1.78	0.65
1:A:244:VAL:HG11	1:A:307:TYR:CE1	2.34	0.63
1:A:133:GLN:NE2	4:A:561:HOH:O	2.40	0.55
1:A:196:VAL:HG21	1:A:235:ASN:CG	2.27	0.55
1:A:223:GLU:HG3	1:A:252:LEU:HD22	1.88	0.55
1:A:12:GLN:HG2	1:A:40:PRO:HB2	1.88	0.54
1:A:21:HIS:HD2	4:A:541:HOH:O	1.92	0.52
1:A:226:LEU:HD13	1:A:280:PRO:HB3	1.93	0.51
1:A:279:ALA:HB1	1:A:281:GLN:HE22	1.75	0.50
1:A:210:SER:O	1:A:214:GLU:HG3	2.13	0.49
1:A:223:GLU:HG3	1:A:252:LEU:CD2	2.42	0.49
1:A:64:LYS:HD3	1:A:64:LYS:HA	1.52	0.47
1:A:277:ARG:HD3	4:A:592:HOH:O	2.13	0.47
1:A:196:VAL:CG2	1:A:248:SER:HB3	2.46	0.45
1:A:277:ARG:HD3	1:A:277:ARG:HH11	1.32	0.45
1:A:281:GLN:H	1:A:281:GLN:CD	2.21	0.44
1:A:176:ARG:NH1	4:A:627:HOH:O	2.34	0.44
1:A:41:MET:SD	1:A:66:LEU:HD23	2.59	0.43
1:A:24:LYS:HE3	1:A:28:MET:CE	2.48	0.43
1:A:279:ALA:CB	1:A:281:GLN:HE22	2.32	0.42
1:A:68:VAL:O	1:A:72:GLU:HG2	2.19	0.42
1:A:281:GLN:NE2	4:A:586:HOH:O	2.52	0.42
1:A:102:GLU:HG3	1:A:103:ASP:N	2.35	0.41
1:A:133:GLN:HG3	1:A:145:ILE:HG21	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:413:HOH:O	4:A:506:HOH:O[3_546]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	292/313 (93%)	288 (99%)	3 (1%)	1 (0%)	50 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	236/251 (94%)	229 (97%)	7 (3%)	53 25

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	PRO
1	A	47	VAL
1	A	48	ILE
1	A	73	ASN
1	A	128	SER
1	A	243	GLN
1	A	281	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	133	GLN
1	A	235	ASN
1	A	243	GLN
1	A	281	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DPM	A	314	1	31,31,31	3.02	12 (38%)	42,43,43	2.97	19 (45%)
3	ACY	A	315	-	3,3,3	2.37	1 (33%)	3,3,3	2.52	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DPM	A	314	1	1/1/4/4	0/20/22/22	0/2/2/2
3	ACY	A	315	-	-	0/0/0/0	0/0/0/0

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	314	DPM	CHB-C1B	-9.42	1.40	1.50
2	A	314	DPM	CHB-C4A	-6.40	1.44	1.50
2	A	314	DPM	C1B-C2B	5.62	1.46	1.37
2	A	314	DPM	C4B-NB	5.48	1.44	1.36
3	A	315	ACY	CH3-C	3.72	1.65	1.49
2	A	314	DPM	C1A-NA	3.43	1.43	1.38
2	A	314	DPM	C4B-C3B	3.26	1.43	1.38
2	A	314	DPM	C1B-NB	3.03	1.42	1.37
2	A	314	DPM	C8B-C9B	3.03	1.58	1.50
2	A	314	DPM	C8A-C9A	2.97	1.58	1.50
2	A	314	DPM	C4A-C3A	2.77	1.42	1.37
2	A	314	DPM	C5B-C6B	2.58	1.57	1.50
2	A	314	DPM	C1A-C2A	2.53	1.42	1.38

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	314	DPM	C7A-C3A-C4A	-10.25	122.39	127.22
2	A	314	DPM	C1B-CHB-C4A	7.35	124.90	113.65
2	A	314	DPM	CHB-C1B-C2B	-4.90	122.67	130.25
2	A	314	DPM	CHB-C4A-C3A	-4.73	122.94	130.25
2	A	314	DPM	C3A-C4A-NA	4.39	109.27	105.77
2	A	314	DPM	O4A-C9A-C8A	4.03	128.47	114.22
2	A	314	DPM	C2A-C1A-NA	3.83	108.82	105.77
2	A	314	DPM	O2B-C6B-O1B	-3.67	113.96	123.30
3	A	315	ACY	OXT-C-O	3.49	136.19	121.96
2	A	314	DPM	C8B-C7B-C3B	-3.38	104.28	112.24
2	A	314	DPM	CHB-C1B-NB	3.33	130.08	123.16
2	A	314	DPM	O2B-C6B-C5B	3.21	121.86	113.80
2	A	314	DPM	O4A-C9A-O3A	-2.95	115.80	123.30
2	A	314	DPM	O4B-C9B-C8B	2.65	123.58	114.22
3	A	315	ACY	O-C-CH3	-2.58	110.82	122.06
2	A	314	DPM	C7B-C3B-C2B	-2.53	123.08	129.38
2	A	314	DPM	O3B-C9B-C8B	-2.46	114.57	123.03
2	A	314	DPM	C7A-C8A-C9A	-2.44	105.63	113.47
2	A	314	DPM	O3A-C9A-C8A	-2.30	115.13	123.03
2	A	314	DPM	C1A-NA-C4A	-2.01	106.03	109.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	314	DPM	O2A-C6A-O1A	-2.00	118.21	123.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	314	DPM	C4A

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	296/313 (94%)	-0.15	11 (3%)	39 48	12, 21, 41, 72	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	VAL	7.2
1	A	58	GLY	5.1
1	A	3	ASP	4.5
1	A	306	VAL	3.9
1	A	59	LYS	3.7
1	A	307	TYR	3.6
1	A	48	ILE	3.2
1	A	42	VAL	3.1
1	A	243	GLN	2.7
1	A	304	ALA	2.2
1	A	305	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACY	A	315	4/4	0.21	43.37	25,39,47,57	0
2	DPM	A	314	30/30	0.12	1.32	15,18,25,29	0

6.5 Other polymers ⓘ

There are no such residues in this entry.