



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 02:27 PM GMT

PDB ID : 1PF9
Title : GroEL-GroES-ADP
Authors : Chaudhry, C.; Farr, G.W.; Todd, M.J.; Rye, H.S.; Brunger, A.T.; Adams, P.D.; Horwich, A.L.; Sigler, P.B.
Deposited on : 2003-05-24
Resolution : 2.99 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	524	
1	B	524	
1	C	524	
1	D	524	
1	E	524	
1	F	524	
1	G	524	
1	H	524	
1	I	524	
1	J	524	
1	K	524	
1	L	524	
1	M	524	
1	N	524	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	O	97	
2	P	97	
2	Q	97	
2	R	97	
2	S	97	
2	T	97	
2	U	97	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	MG	A	601	-	X
3	MG	D	604	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 59283 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called groEL protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	B	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	C	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	D	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	E	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	F	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	G	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	H	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	I	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	J	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	K	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	L	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	M	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	N	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			


- Molecule 2 is a protein called groES protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	P	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	Q	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	R	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	S	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	T	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	U	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	F	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	G	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	1	Total	O	0	0
			1	1		
5	C	1	Total	O	0	0
			1	1		
5	D	1	Total	O	0	0
			1	1		
5	E	1	Total	O	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	1	Total	O	0	0
			1	1		
5	G	1	Total	O	0	0
			1	1		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: groEL protein




- Molecule 1: groEL protein






- Molecule 1: groEL protein

Chain E:



- Molecule 1: groEL protein


Chain F:



- Molecule 1: groEL protein


Chain G:






• Molecule 1: groEL protein


Chain H:



• Molecule 1: groEL protein


Chain I:






• Molecule 1: groEL protein


Chain J:



• Molecule 1: groEL protein


Chain K:






• Molecule 1: groEL protein


Chain L:



• Molecule 1: groEL protein


Chain M:






• Molecule 1: groEL protein

Chain N:




• Molecule 2: groES protein


Chain O:



• Molecule 2: groES protein


Chain P:





- Molecule 2: groES protein

Chain Q:




- Molecule 2: groES protein

Chain R:




- Molecule 2: groES protein

Chain S:




- Molecule 2: groES protein


Chain T:



- Molecule 2: groES protein

Chain U:





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	255.26Å 265.25Å 184.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.16 – 2.99 40.07 – 2.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.16-2.99) 96.7 (40.07-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.269 , 0.287 0.274 , 0.291	Depositor DCC
R_{free} test set	12081 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	50.3	Xtriage
Anisotropy	0.614	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 23.5	EDS
Estimated twinning fraction	0.003 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 242645 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	59283	wwPDB-VP
Average B, all atoms (Å ²)	7.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/3884	0.90	18/5243 (0.3%)
1	B	0.60	0/3884	0.95	18/5243 (0.3%)
1	C	0.64	0/3884	0.91	20/5243 (0.4%)
1	D	0.66	1/3884 (0.0%)	0.90	17/5243 (0.3%)
1	E	0.57	0/3884	0.86	15/5243 (0.3%)
1	F	0.55	0/3884	0.86	17/5243 (0.3%)
1	G	0.64	0/3884	0.91	20/5243 (0.4%)
1	H	0.59	0/3884	0.85	14/5243 (0.3%)
1	I	0.64	0/3884	0.86	15/5243 (0.3%)
1	J	0.61	0/3884	0.87	18/5243 (0.3%)
1	K	0.54	0/3884	0.83	17/5243 (0.3%)
1	L	0.51	0/3884	0.87	20/5243 (0.4%)
1	M	0.59	0/3884	0.90	20/5243 (0.4%)
1	N	0.59	0/3884	0.85	17/5243 (0.3%)
2	O	0.37	0/732	0.73	3/983 (0.3%)
2	P	0.37	0/732	0.73	4/983 (0.4%)
2	Q	0.38	0/732	0.73	2/983 (0.2%)
2	R	0.41	0/732	0.74	4/983 (0.4%)
2	S	0.39	0/732	0.74	4/983 (0.4%)
2	T	0.39	0/732	0.74	3/983 (0.3%)
2	U	0.38	0/732	0.72	3/983 (0.3%)
All	All	0.59	1/59500 (0.0%)	0.87	269/80283 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	N	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	130	GLU	CD-OE1	5.35	1.31	1.25

The worst 5 of 269 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	231	ARG	NE-CZ-NH1	-19.37	110.61	120.30
1	B	231	ARG	NE-CZ-NH2	18.28	129.44	120.30
1	L	268	ARG	NE-CZ-NH2	14.85	127.72	120.30
1	M	268	ARG	NE-CZ-NH2	14.77	127.68	120.30
1	M	268	ARG	NE-CZ-NH1	-14.67	112.97	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	N	433	ASN	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3856	0	3976	105	0
1	B	3856	0	3976	99	0
1	C	3856	0	3976	109	0
1	D	3856	0	3976	115	0
1	E	3856	0	3976	104	0
1	F	3856	0	3976	99	0
1	G	3856	0	3976	113	0
1	H	3856	0	3976	107	0
1	I	3856	0	3976	104	0
1	J	3856	0	3976	106	0
1	K	3856	0	3976	110	0
1	L	3856	0	3976	112	0
1	M	3856	0	3976	115	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	N	3856	0	3976	111	0
2	O	728	0	762	20	0
2	P	728	0	762	21	0
2	Q	728	0	762	18	0
2	R	728	0	762	17	0
2	S	728	0	762	19	0
2	T	728	0	762	15	0
2	U	728	0	762	17	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
4	A	27	0	12	2	0
4	B	27	0	12	2	0
4	C	27	0	12	2	0
4	D	27	0	12	3	0
4	E	27	0	12	1	0
4	F	27	0	12	1	0
4	G	27	0	12	3	0
5	A	1	0	0	1	0
5	B	1	0	0	1	0
5	C	1	0	0	1	0
5	D	1	0	0	1	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	1	0
All	All	59283	0	61082	1567	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 1567 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:E:114:MET:CE	1:E:114:MET:SD	2.05	1.45
1:C:73:MET:SD	1:C:73:MET:CE	2.05	1.42
1:C:18:ARG:HG2	1:C:18:ARG:HH11	1.07	1.19
1:G:18:ARG:HH11	1:G:18:ARG:HG2	1.07	1.17
1:D:432:GLN:NE2	1:D:436:GLN:HE22	1.47	1.13

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/524 (100%)	485 (93%)	33 (6%)	4 (1%)	27	76
1	B	522/524 (100%)	484 (93%)	33 (6%)	5 (1%)	22	70
1	C	522/524 (100%)	484 (93%)	34 (6%)	4 (1%)	27	76
1	D	522/524 (100%)	485 (93%)	32 (6%)	5 (1%)	22	70
1	E	522/524 (100%)	489 (94%)	29 (6%)	4 (1%)	27	76
1	F	522/524 (100%)	490 (94%)	27 (5%)	5 (1%)	22	70
1	G	522/524 (100%)	481 (92%)	36 (7%)	5 (1%)	22	70
1	H	522/524 (100%)	481 (92%)	36 (7%)	5 (1%)	22	70
1	I	522/524 (100%)	483 (92%)	32 (6%)	7 (1%)	18	62
1	J	522/524 (100%)	480 (92%)	38 (7%)	4 (1%)	27	76
1	K	522/524 (100%)	479 (92%)	38 (7%)	5 (1%)	22	70
1	L	522/524 (100%)	483 (92%)	34 (6%)	5 (1%)	22	70
1	M	522/524 (100%)	480 (92%)	37 (7%)	5 (1%)	22	70
1	N	522/524 (100%)	483 (92%)	35 (7%)	4 (1%)	27	76
2	O	95/97 (98%)	77 (81%)	13 (14%)	5 (5%)	3	18
2	P	95/97 (98%)	77 (81%)	13 (14%)	5 (5%)	3	18
2	Q	95/97 (98%)	76 (80%)	13 (14%)	6 (6%)	2	12
2	R	95/97 (98%)	77 (81%)	13 (14%)	5 (5%)	3	18
2	S	95/97 (98%)	77 (81%)	13 (14%)	5 (5%)	3	18
2	T	95/97 (98%)	76 (80%)	13 (14%)	6 (6%)	2	12
2	U	95/97 (98%)	76 (80%)	14 (15%)	5 (5%)	3	18
All	All	7973/8015 (100%)	7303 (92%)	566 (7%)	104 (1%)	18	62

5 of 104 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PHE
1	B	44	PHE
1	C	44	PHE
1	D	44	PHE
1	E	44	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/404 (100%)	359 (89%)	45 (11%)	9	34
1	B	404/404 (100%)	358 (89%)	46 (11%)	8	33
1	C	404/404 (100%)	356 (88%)	48 (12%)	8	30
1	D	404/404 (100%)	357 (88%)	47 (12%)	8	31
1	E	404/404 (100%)	358 (89%)	46 (11%)	8	33
1	F	404/404 (100%)	357 (88%)	47 (12%)	8	31
1	G	404/404 (100%)	354 (88%)	50 (12%)	7	28
1	H	404/404 (100%)	347 (86%)	57 (14%)	5	23
1	I	404/404 (100%)	349 (86%)	55 (14%)	5	24
1	J	404/404 (100%)	349 (86%)	55 (14%)	5	24
1	K	404/404 (100%)	347 (86%)	57 (14%)	5	23
1	L	404/404 (100%)	348 (86%)	56 (14%)	5	23
1	M	404/404 (100%)	347 (86%)	57 (14%)	5	23
1	N	404/404 (100%)	349 (86%)	55 (14%)	5	24
2	O	80/80 (100%)	70 (88%)	10 (12%)	7	28
2	P	80/80 (100%)	70 (88%)	10 (12%)	7	28
2	Q	80/80 (100%)	70 (88%)	10 (12%)	7	28
2	R	80/80 (100%)	69 (86%)	11 (14%)	5	24
2	S	80/80 (100%)	69 (86%)	11 (14%)	5	24
2	T	80/80 (100%)	70 (88%)	10 (12%)	7	28
2	U	80/80 (100%)	69 (86%)	11 (14%)	5	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	6216/6216 (100%)	5422 (87%)	794 (13%)	6 27

5 of 794 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	300	VAL
1	J	37	ASN
2	P	6	LEU
1	H	385	THR
1	I	187	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 135 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	153	ASN
1	H	436	GLN
2	P	68	ASN
1	G	265	ASN
1	H	21	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 7 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	A	700	3	29,29,29	1.15	3 (10%)	45,45,45	2.64	18 (40%)
4	ADP	B	701	3	29,29,29	1.24	3 (10%)	45,45,45	2.53	14 (31%)
4	ADP	C	702	3	29,29,29	1.12	3 (10%)	45,45,45	2.63	17 (37%)
4	ADP	D	703	3	29,29,29	1.34	5 (17%)	45,45,45	2.67	15 (33%)
4	ADP	E	704	3	29,29,29	1.12	3 (10%)	45,45,45	2.52	14 (31%)
4	ADP	F	705	3	29,29,29	0.99	2 (6%)	45,45,45	2.38	12 (26%)
4	ADP	G	706	3	29,29,29	1.09	1 (3%)	45,45,45	2.51	17 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	700	3	-	0/16/32/32	0/1/3/3
4	ADP	B	701	3	-	0/16/32/32	0/1/3/3
4	ADP	C	702	3	-	0/16/32/32	0/1/3/3
4	ADP	D	703	3	-	0/16/32/32	0/1/3/3
4	ADP	E	704	3	-	0/16/32/32	0/1/3/3
4	ADP	F	705	3	-	0/16/32/32	0/1/3/3
4	ADP	G	706	3	-	0/16/32/32	0/1/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	706	ADP	C2-N3	3.57	1.39	1.32
4	B	701	ADP	C2-N3	3.36	1.38	1.32
4	D	703	ADP	C2-N3	3.12	1.38	1.32
4	A	700	ADP	C2-N3	3.03	1.38	1.32
4	E	704	ADP	C2-N3	3.01	1.38	1.32

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	703	ADP	N3-C2-N1	-12.03	118.65	128.71
4	A	700	ADP	N3-C2-N1	-11.33	119.24	128.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	705	ADP	N3-C2-N1	-10.79	119.69	128.71
4	C	702	ADP	N3-C2-N1	-10.43	119.99	128.71
4	E	704	ADP	N3-C2-N1	-10.40	120.02	128.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/524 (100%)	0.62	59 (11%) 6 2	5, 7, 9, 10	0
1	B	524/524 (100%)	0.70	69 (13%) 4 1	5, 7, 9, 10	0
1	C	524/524 (100%)	0.48	50 (9%) 8 2	5, 7, 9, 10	0
1	D	524/524 (100%)	0.55	53 (10%) 7 2	5, 7, 9, 10	0
1	E	524/524 (100%)	0.83	84 (16%) 3 1	5, 7, 9, 10	0
1	F	524/524 (100%)	0.89	78 (14%) 3 1	5, 7, 9, 10	0
1	G	524/524 (100%)	0.70	67 (12%) 4 1	5, 7, 9, 10	0
1	H	524/524 (100%)	0.40	22 (4%) 35 7	6, 7, 9, 10	0
1	I	524/524 (100%)	0.32	15 (2%) 49 9	5, 7, 9, 10	0
1	J	524/524 (100%)	0.45	33 (6%) 19 5	5, 7, 9, 10	0
1	K	524/524 (100%)	0.63	49 (9%) 9 2	5, 7, 9, 10	0
1	L	524/524 (100%)	0.63	43 (8%) 12 3	5, 7, 9, 10	0
1	M	524/524 (100%)	0.50	39 (7%) 14 3	5, 7, 9, 10	0
1	N	524/524 (100%)	0.55	45 (8%) 11 3	5, 7, 9, 10	0
2	O	97/97 (100%)	1.83	39 (40%) 1 0	7, 7, 8, 8	0
2	P	97/97 (100%)	1.74	32 (32%) 1 0	7, 7, 8, 8	0
2	Q	97/97 (100%)	1.47	30 (30%) 1 0	7, 7, 8, 8	0
2	R	97/97 (100%)	1.52	28 (28%) 1 0	7, 7, 8, 8	0
2	S	97/97 (100%)	1.87	37 (38%) 1 0	7, 7, 8, 8	0
2	T	97/97 (100%)	1.74	30 (30%) 1 0	7, 7, 8, 8	0
2	U	97/97 (100%)	3.40	78 (80%) 0 0	7, 7, 8, 8	0
All	All	8015/8015 (100%)	0.70	980 (12%) 5 1	5, 7, 9, 10	0

The worst 5 of 980 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	361	ASP	15.0
1	F	361	ASP	10.3
1	A	361	ASP	9.8
1	F	210	THR	9.2
1	G	208	PRO	8.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	D	604	1/1	0.32	14.84	2,2,2,2	0
3	MG	A	601	1/1	0.29	12.83	2,2,2,2	0
3	MG	G	607	1/1	0.21	1.28	2,2,2,2	0
3	MG	B	602	1/1	0.23	0.95	2,2,2,2	0
3	MG	C	603	1/1	0.20	0.74	2,2,2,2	0
4	ADP	B	701	27/27	0.21	0.13	6,8,9,10	0
4	ADP	A	700	27/27	0.18	0.06	6,8,9,11	0
3	MG	F	606	1/1	0.25	-0.12	2,2,2,2	0
4	ADP	F	705	27/27	0.21	-0.17	6,8,9,10	0
4	ADP	E	704	27/27	0.20	-0.26	6,8,9,10	0
4	ADP	D	703	27/27	0.16	-0.71	6,8,9,10	0
4	ADP	G	706	27/27	0.16	-0.78	6,8,9,10	0
3	MG	E	605	1/1	0.20	-1.24	2,2,2,2	0
4	ADP	C	702	27/27	0.14	-1.63	6,8,9,10	0

6.5 Other polymers ⓘ

There are no such residues in this entry.