



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 03:24 PM GMT

PDB ID : 2PFH
Title : Complex of Aldose Reductase with NADP+ and simultaneously bound competitive inhibitors Fidarestat and IDD594. Concentration of Fidarestat in soaking solution is less than concentration of IDD594.
Authors : Petrova, T.; Hazemann, I.; Cousido, A.; Mitschler, A.; Ginell, S.; Joachimiak, A.; Podjarny, A.
Deposited on : 2007-04-05
Resolution : 0.85 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 0.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1084 (1.12-0.62)
Clashscore	79885	1127 (1.10-0.62)
Ramachandran outliers	78287	1051 (1.10-0.62)
Sidechain outliers	78261	1050 (1.10-0.62)
RSRZ outliers	66119	1084 (1.12-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	316	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
6	CIT	A	450[A]	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 3919 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	3098	1992	518	571	17	7	105	1


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ILE	LEU	SEE REMARK 999	UNP P15121

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	1
			1	1		

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is IDD594 (three-letter code: LDT) (formula: $C_{16}H_{12}BrF_2NO_3S$).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
4	A	1	Total	Br	C	F	N	O	S	0	1
			24	1	16	2	1	3	1		

- Molecule 5 is (2S,4S)-2-AMINOFORMYL-6-FLUORO-SPIRO[CHROMAN-4,4'-IMIDAZO LIDINE]-2',5'-DIONE (three-letter code: FID) (formula: $C_{12}H_{10}FN_3O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	1
			20	12	1	3	4		

- Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			13	6	7		
6	A	1	Total	C	O	0	1
			13	6	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	656	Total	O	0	305
			702	702		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

● Molecule 1: Aldose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.15Å 66.55Å 47.24Å 90.00° 92.31° 90.00°	Depositor
Resolution (Å)	50.00 – 0.85 34.74 – 0.85	Depositor EDS
% Data completeness (in resolution range)	95.4 (50.00-0.85) 95.4 (34.74-0.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.02 (at 0.85Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.082 , 0.095 0.095 , 0.096	Depositor DCC
R_{free} test set	12783 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	5.3	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 84.6	EDS
Estimated twinning fraction	0.002 for -l,k,h 0.018 for -h,-k,l 0.015 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 253543 reflections	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3919	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, CIT, LDT, FID, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	6.18	17/3368 (0.5%)	2.58	60/4566 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	312[A]	HIS	CG-CD2	178.57	4.39	1.35
1	A	312[B]	HIS	CG-CD2	178.57	4.39	1.35
1	A	312[A]	HIS	CG-ND1	124.21	4.12	1.38
1	A	312[B]	HIS	CG-ND1	124.21	4.12	1.38
1	A	26[A]	GLN	CD-NE2	90.05	3.58	1.32
1	A	26[B]	GLN	CD-NE2	90.05	3.58	1.32
1	A	26[A]	GLN	CD-OE1	77.60	2.94	1.24
1	A	26[B]	GLN	CD-OE1	77.60	2.94	1.24
1	A	293[A]	ARG	CB-CG	-7.41	1.32	1.52
1	A	293[B]	ARG	CB-CG	-7.41	1.32	1.52
1	A	29[A]	GLU	CD-OE1	-6.96	1.18	1.25
1	A	29[B]	GLU	CD-OE1	-6.96	1.18	1.25
1	A	64[A]	GLU	CD-OE1	-5.65	1.19	1.25
1	A	64[B]	GLU	CD-OE1	-5.65	1.19	1.25
1	A	150	GLU	CD-OE2	-5.45	1.19	1.25
1	A	40[A]	ARG	CZ-NH2	5.21	1.39	1.33
1	A	40[B]	ARG	CZ-NH2	5.21	1.39	1.33

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312[A]	HIS	ND1-CG-CD2	-54.82	29.25	106.00
1	A	312[B]	HIS	ND1-CG-CD2	-54.82	29.25	106.00
1	A	26[A]	GLN	CG-CD-OE1	-40.72	40.15	121.60
1	A	26[B]	GLN	CG-CD-OE1	-40.72	40.15	121.60
1	A	26[A]	GLN	OE1-CD-NE2	-36.71	37.48	121.90
1	A	26[B]	GLN	OE1-CD-NE2	-36.71	37.48	121.90
1	A	26[A]	GLN	CG-CD-NE2	-34.63	33.59	116.70
1	A	26[B]	GLN	CG-CD-NE2	-34.63	33.59	116.70
1	A	312[A]	HIS	CG-ND1-CE1	32.17	153.24	108.20
1	A	312[B]	HIS	CG-ND1-CE1	32.17	153.24	108.20
1	A	312[A]	HIS	CB-CG-ND1	-20.63	71.63	123.20
1	A	312[B]	HIS	CB-CG-ND1	-20.63	71.63	123.20
1	A	3	ARG	NE-CZ-NH2	-15.92	112.34	120.30
1	A	312[A]	HIS	CG-CD2-NE2	14.98	137.65	109.20
1	A	312[B]	HIS	CG-CD2-NE2	14.98	137.65	109.20
1	A	232	ARG	NE-CZ-NH1	11.01	125.80	120.30
1	A	293[A]	ARG	NE-CZ-NH1	-10.57	115.01	120.30
1	A	293[B]	ARG	NE-CZ-NH1	-10.57	115.01	120.30
1	A	276[A]	PHE	CB-CG-CD1	10.03	127.82	120.80
1	A	276[B]	PHE	CB-CG-CD1	10.03	127.82	120.80
1	A	312[A]	HIS	CB-CG-CD2	-9.74	100.59	130.80
1	A	312[B]	HIS	CB-CG-CD2	-9.74	100.59	130.80
1	A	217[A]	ARG	NE-CZ-NH2	-9.21	115.69	120.30
1	A	217[B]	ARG	NE-CZ-NH2	-9.21	115.69	120.30
1	A	217[A]	ARG	NE-CZ-NH1	8.29	124.44	120.30
1	A	217[B]	ARG	NE-CZ-NH1	8.29	124.44	120.30
1	A	268	ARG	NE-CZ-NH1	7.89	124.25	120.30
1	A	224	ASP	CB-CG-OD1	7.62	125.16	118.30
1	A	63[A]	ARG	NE-CZ-NH2	-7.59	116.51	120.30
1	A	216	ASP	CB-CG-OD2	-7.04	111.96	118.30
1	A	276[A]	PHE	CB-CG-CD2	-6.66	116.14	120.80
1	A	276[B]	PHE	CB-CG-CD2	-6.66	116.14	120.80
1	A	3	ARG	NH1-CZ-NH2	6.59	126.66	119.40
1	A	134[A]	ASP	CB-CG-OD2	-6.59	112.36	118.30
1	A	134[B]	ASP	CB-CG-OD2	-6.59	112.36	118.30
1	A	82	TYR	CB-CG-CD1	6.43	124.86	121.00
1	A	296[A]	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	296[B]	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	277[A]	ASP	CB-CG-OD2	-5.87	113.01	118.30
1	A	277[B]	ASP	CB-CG-OD2	-5.87	113.01	118.30
1	A	278[A]	PHE	CB-CG-CD1	5.82	124.87	120.80
1	A	278[B]	PHE	CB-CG-CD1	5.82	124.87	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	TYR	CB-CG-CD1	5.73	124.44	121.00
1	A	193[A]	GLU	CB-CA-C	-5.69	99.01	110.40
1	A	193[B]	GLU	CB-CA-C	-5.69	99.01	110.40
1	A	135[A]	THR	O-C-N	-5.62	113.72	122.70
1	A	135[B]	THR	O-C-N	-5.62	113.72	122.70
1	A	135[C]	THR	O-C-N	-5.62	113.72	122.70
1	A	291	TYR	CD1-CE1-CZ	5.59	124.83	119.80
1	A	73	PHE	CB-CG-CD1	5.48	124.64	120.80
1	A	67[A]	VAL	CB-CA-C	-5.41	101.12	111.40
1	A	67[B]	VAL	CB-CA-C	-5.41	101.12	111.40
1	A	232	ARG	NH1-CZ-NH2	-5.32	113.54	119.40
1	A	291	TYR	CB-CG-CD2	5.19	124.12	121.00
1	A	209	TYR	CA-CB-CG	5.16	123.21	113.40
1	A	70[A]	GLU	CB-CG-CD	5.06	127.85	114.20
1	A	70[B]	GLU	CB-CG-CD	5.06	127.85	114.20
1	A	115	PHE	CB-CG-CD2	-5.05	117.27	120.80
1	A	127[A]	SER	O-C-N	-5.02	114.67	123.20
1	A	127[B]	SER	O-C-N	-5.02	114.67	123.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	0	11	7
2	A	1	0	0	1	0
3	A	48	0	0	0	0
4	A	24	0	0	1	0
5	A	20	0	0	3	0
6	A	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	702	0	0	6	9
All	All	3919	0	0	11	9

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (11) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:295[A]:TRP:CZ3	7:A:3675[A]:HOH:O	2.29	0.84
1:A:136[C]:ASN:N	7:A:3302:HOH:O	2.43	0.52
1:A:110:HIS:ND1	2:A:2000[A]:CL:CL	2.80	0.52
1:A:226:SER:N	7:A:3634[B]:HOH:O	2.45	0.49
1:A:113:THR:OG1	4:A:320[A]:LDT:BR8	2.85	0.49
1:A:111:TRP:NE1	5:A:321[B]:FID:O6I	2.47	0.48
1:A:116[A]:LYS:NZ	7:A:3267:HOH:O	2.48	0.47
1:A:194[A]:LYS:NZ	7:A:3504[A]:HOH:O	2.48	0.46
1:A:298[B]:CYS:SG	5:A:321[B]:FID:C8I	3.06	0.44
1:A:48:TYR:OH	5:A:321[B]:FID:O3I	2.35	0.44
1:A:242[B]:LYS:NZ	7:A:3471[B]:HOH:O	2.53	0.42

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:202[A]:LYS:CB	7:A:3360:HOH:O[2_546]	1.25	0.95
1:A:202[A]:LYS:CA	7:A:3360:HOH:O[2_546]	1.30	0.90
1:A:202[B]:LYS:CB	7:A:3360:HOH:O[2_546]	1.35	0.85
1:A:202[B]:LYS:CA	7:A:3360:HOH:O[2_546]	1.68	0.52
1:A:202[A]:LYS:N	7:A:3360:HOH:O[2_546]	1.71	0.49
1:A:202[B]:LYS:N	7:A:3360:HOH:O[2_546]	1.78	0.42
7:A:3492[A]:HOH:O	7:A:3550:HOH:O[1_554]	1.86	0.34
7:A:3402:HOH:O	7:A:3479[A]:HOH:O[1_655]	1.97	0.23
1:A:202[A]:LYS:C	7:A:3360:HOH:O[2_546]	2.03	0.17

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/316 (129%)	404 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/281 (131%)	368 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	A	318	-	52,52,52	1.85	10 (19%)	80,80,80	1.45	13 (16%)
4	LDT	A	320[A]	-	25,25,25	1.36	2 (8%)	34,34,34	1.53	6 (17%)
5	FID	A	321[B]	-	22,22,22	1.65	1 (4%)	34,34,34	2.86	13 (38%)
6	CIT	A	400[A]	-	12,12,12	1.85	5 (41%)	17,17,17	3.16	4 (23%)
6	CIT	A	450[A]	-	12,12,12	0.91	0	17,17,17	2.18	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	A	318	-	-	0/35/77/77	0/3/5/5
4	LDT	A	320[A]	-	-	0/14/14/14	0/2/2/2
5	FID	A	321[B]	-	-	0/4/34/34	0/1/3/3
6	CIT	A	400[A]	-	-	0/16/16/16	0/0/0/0
6	CIT	A	450[A]	-	-	0/16/16/16	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	321[B]	FID	C8I-C7I	-6.71	1.47	1.54
3	A	318	NDP	C4N-C3N	-6.52	1.37	1.50
3	A	318	NDP	C4N-C5N	-4.51	1.39	1.49
3	A	318	NDP	C1D-N1N	3.81	1.54	1.46
3	A	318	NDP	C6N-C5N	3.77	1.41	1.33
3	A	318	NDP	C2N-C3N	3.73	1.42	1.34
3	A	318	NDP	PA-O3	3.35	1.65	1.59
4	A	320[A]	LDT	C6-C5	3.23	1.43	1.36
6	A	400[A]	CIT	O1-C1	2.86	1.32	1.22
6	A	400[A]	CIT	O3-C5	2.74	1.32	1.22
6	A	400[A]	CIT	C3-C6	2.72	1.56	1.53
3	A	318	NDP	O4B-C1B	2.66	1.45	1.41
6	A	400[A]	CIT	O5-C6	2.60	1.31	1.22
6	A	400[A]	CIT	O4-C5	-2.46	1.21	1.30
3	A	318	NDP	C4A-N3A	-2.25	1.32	1.35
3	A	318	NDP	PN-O3	-2.21	1.55	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	318	NDP	C3B-C2B	2.10	1.58	1.53
4	A	320[A]	LDT	O34-C32	2.05	1.29	1.22

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	400[A]	CIT	O5-C6-C3	-9.86	108.60	122.20
5	A	321[B]	FID	C7I-C8I-C9	7.29	122.33	110.27
3	A	318	NDP	C8A-N9A-C4A	-6.60	101.86	106.90
6	A	450[A]	CIT	O6-C6-C3	6.30	122.06	112.89
5	A	321[B]	FID	O10-C11-C12	6.19	128.91	122.85
6	A	400[A]	CIT	O6-C6-C3	5.82	121.35	112.89
5	A	321[B]	FID	C7I-C5-N4	-5.66	103.38	107.59
5	A	321[B]	FID	O6I-C5-N4	5.21	133.79	126.30
6	A	450[A]	CIT	O5-C6-C3	-4.65	115.79	122.20
5	A	321[B]	FID	C8I-C7I-N1I	4.50	119.98	112.27
4	A	320[A]	LDT	C13-N17-C11	-4.07	121.59	124.85
3	A	318	NDP	N3A-C2A-N1A	4.01	132.06	128.71
6	A	400[A]	CIT	O7-C3-C6	3.97	114.68	108.95
5	A	321[B]	FID	C5-N4-C2I	3.91	115.40	111.67
5	A	321[B]	FID	O6I-C5-C7I	-3.91	122.59	126.05
5	A	321[B]	FID	C8I-C7I-C12	-3.74	102.99	111.39
5	A	321[B]	FID	N4-C2I-N1I	-3.69	103.55	107.64
3	A	318	NDP	PN-O3-PA	2.86	140.07	131.68
4	A	320[A]	LDT	F9-C5-C2	-2.79	113.90	118.22
6	A	400[A]	CIT	C4-C3-C6	-2.62	104.05	110.12
5	A	321[B]	FID	C13-C12-C11	2.60	120.70	116.67
5	A	321[B]	FID	O10-C9-C8I	-2.51	106.09	111.32
6	A	450[A]	CIT	O7-C3-C6	-2.51	105.34	108.95
5	A	321[B]	FID	O10-C11-C16	-2.46	111.91	116.72
3	A	318	NDP	C5A-C6A-N6A	2.42	126.19	120.72
3	A	318	NDP	O2D-C2D-C3D	-2.40	104.01	111.83
5	A	321[B]	FID	C8I-C9-C19	2.40	115.64	112.20
3	A	318	NDP	C6A-C5A-C4A	2.40	121.65	117.25
3	A	318	NDP	C2N-C3N-C7N	2.34	124.47	118.49
4	A	320[A]	LDT	O33-C32-O34	2.26	129.04	123.30
3	A	318	NDP	C6N-N1N-C2N	-2.21	116.55	119.44
3	A	318	NDP	O4B-C1B-N9A	2.19	110.48	108.44
3	A	318	NDP	P2B-O2B-C2B	2.15	126.49	121.96
3	A	318	NDP	C1B-N9A-C4A	2.13	130.31	126.64
3	A	318	NDP	C2A-N3A-C4A	-2.12	107.99	114.01
6	A	450[A]	CIT	O7-C3-C4	2.12	113.49	109.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	320[A]	LDT	C28-C27-C24	2.06	127.09	124.16
4	A	320[A]	LDT	C3-C7-C4	2.04	120.95	118.20
3	A	318	NDP	O2X-P2B-O2B	-2.03	101.25	107.09
4	A	320[A]	LDT	O15-C4-C2	-2.02	118.68	123.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/316 (99%)	-0.73	2 (0%) 86 82	3, 6, 13, 27	26 (8%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0[A]	MET	5.1
1	A	1	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	CIT	A	450[A]	13/13	0.22	40.70	1,6,7,8	13
6	CIT	A	400[A]	13/13	0.10	1.35	7,10,12,16	13
5	FID	A	321[B]	20/20	0.05	0.22	3,4,7,9	20

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NDP	A	318	48/48	0.03	-0.96	2,3,5,6	0
4	LDT	A	320[A]	24/24	0.03	-1.42	3,4,5,5	24
2	CL	A	2000[A]	1/1	0.03	-2.14	5,5,5,5	1

6.5 Other polymers ⓘ

There are no such residues in this entry.