



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 08:59 AM GMT

PDB ID : 2PI8
Title : Crystal structure of E. coli MltA with bound chitohexaose
Authors : van Straaten, K.E.; Barends, T.R.M.; Dijkstra, B.W.; Thunnissen, A.M.W.H.
Deposited on : 2007-04-13
Resolution : 2.25 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance





The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	345	
1	B	345	
1	C	345	
1	D	345	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	PO4	A	699	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11043 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-bound lytic murein transglycosylase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	Se	0	0	0
			2599	1645	466	480	8			
1	B	335	Total	C	N	O	Se	0	0	0
			2634	1666	472	488	8			
1	C	332	Total	C	N	O	Se	0	0	0
			2608	1651	468	481	8			
1	D	335	Total	C	N	O	Se	0	0	0
			2634	1666	472	488	8			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
A	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	131	SER	PRO	CONFLICT	UNP P0A935
A	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	272	ILE	LEU	CONFLICT	UNP P0A935
A	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	308	ALA	ASP	ENGINEERED	UNP P0A935
B	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
B	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	131	SER	PRO	CONFLICT	UNP P0A935
B	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	272	ILE	LEU	CONFLICT	UNP P0A935
B	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	308	ALA	ASP	ENGINEERED	UNP P0A935
C	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
C	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	131	SER	PRO	CONFLICT	UNP P0A935
C	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	272	ILE	LEU	CONFLICT	UNP P0A935
C	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	308	ALA	ASP	ENGINEERED	UNP P0A935
D	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
D	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	131	SER	PRO	CONFLICT	UNP P0A935
D	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	272	ILE	LEU	CONFLICT	UNP P0A935
D	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	308	ALA	ASP	ENGINEERED	UNP P0A935

- Molecule 2 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	6	Total	C	N	O	0	0
			85	48	6	31		
2	B	6	Total	C	N	O	0	0
			85	48	6	31		
2	C	6	Total	C	N	O	0	0
			85	48	6	31		
2	D	6	Total	C	N	O	0	0
			85	48	6	31		

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
A	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	131	SER	PRO	CONFLICT	UNP P0A935
A	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	272	ILE	LEU	CONFLICT	UNP P0A935
A	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
A	308	ALA	ASP	ENGINEERED	UNP P0A935
B	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
B	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	131	SER	PRO	CONFLICT	UNP P0A935
B	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	272	ILE	LEU	CONFLICT	UNP P0A935
B	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
B	308	ALA	ASP	ENGINEERED	UNP P0A935
C	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
C	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	131	SER	PRO	CONFLICT	UNP P0A935
C	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	272	ILE	LEU	CONFLICT	UNP P0A935
C	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
C	308	ALA	ASP	ENGINEERED	UNP P0A935
D	1	MSE	-	MODIFIED RESIDUE	UNP P0A935
D	77	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	87	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	123	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	131	SER	PRO	CONFLICT	UNP P0A935
D	154	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	159	MSE	MET	MODIFIED RESIDUE	UNP P0A935

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	206	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	208	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	272	ILE	LEU	CONFLICT	UNP P0A935
D	293	MSE	MET	MODIFIED RESIDUE	UNP P0A935
D	308	ALA	ASP	ENGINEERED	UNP P0A935

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total	O	0	0
			46	46		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	61	Total 61	O 61	0	0
4	C	46	Total 46	O 46	0	0
4	D	45	Total 45	O 45	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Membrane-bound lytic murein transglycosylase A

Chain A:




- Molecule 1: Membrane-bound lytic murein transglycosylase A

Chain B:



- Molecule 1: Membrane-bound lytic murein transglycosylase A

Chain C:



- Molecule 1: Membrane-bound lytic murein transglycosylase A

Chain D:





4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	91.00Å 91.00Å 187.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.00 – 2.25 39.40 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.00-2.25) 97.4 (39.40-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.16Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.182 , 0.226 0.177 , 0.218	Depositor DCC
R_{free} test set	6610 reflections (7.77%)	DCC
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 18.8	EDS
Estimated twinning fraction	0.499 for -h,-k,l 0.499 for h,-h-k,-l 0.499 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 91662 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11043	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2661	0.48	0/3592
1	B	0.26	0/2696	0.47	0/3637
1	C	0.34	0/2670	0.49	0/3603
1	D	0.29	0/2696	0.48	0/3637
All	All	0.30	0/10723	0.48	0/14469

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2599	0	2497	49	0
1	B	2634	0	2534	46	0
1	C	2608	0	2510	64	0
1	D	2634	0	2534	54	0
2	A	85	0	75	2	0
2	B	85	0	75	3	0
2	C	85	0	75	7	0
2	D	85	0	75	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	10	0	0	0	0
3	D	10	0	0	0	0
4	A	46	0	0	0	0
4	B	61	0	0	1	0
4	C	46	0	0	0	0
4	D	45	0	0	0	0
All	All	11043	0	10375	197	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

The worst 5 of 197 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:200:GLU:HG2	1:A:214:TRP:HE1	1.27	0.98
1:D:190:SER:HB3	2:D:936:NAG:H81	1.47	0.95
1:A:259:ALA:HB3	1:A:308:ALA:HB3	1.62	0.80
1:D:259:ALA:HB3	1:D:308:ALA:HB3	1.62	0.79
1:A:27:PRO:HG2	1:A:74:THR:HB	1.65	0.79

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/345 (95%)	302 (92%)	22 (7%)	3 (1%)	25	20
1	B	333/345 (96%)	315 (95%)	16 (5%)	2 (1%)	33	33
1	C	328/345 (95%)	311 (95%)	15 (5%)	2 (1%)	33	33
1	D	333/345 (96%)	310 (93%)	20 (6%)	3 (1%)	25	20
All	All	1321/1380 (96%)	1238 (94%)	73 (6%)	10 (1%)	27	24

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	GLY
1	D	71	GLY
1	B	9	GLY
1	C	13	LYS
1	C	92	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/267 (100%)	260 (98%)	6 (2%)	63	72
1	B	270/267 (101%)	267 (99%)	3 (1%)	84	90
1	C	267/267 (100%)	261 (98%)	6 (2%)	64	74
1	D	270/267 (101%)	266 (98%)	4 (2%)	76	85
All	All	1073/1068 (100%)	1054 (98%)	19 (2%)	71	80

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	329	ARG
1	C	52	ARG
1	D	42	GLN
1	B	326	HIS
1	D	198	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

24 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	901	2	12,14,15	0.42	0	15,19,21	0.68	0
2	NAG	A	902	2	12,14,15	0.49	0	15,19,21	0.66	0
2	NAG	A	903	2	12,14,15	0.45	0	15,19,21	0.71	0
2	NAG	A	904	2	12,14,15	0.42	0	15,19,21	0.66	0
2	NAG	A	905	2	12,14,15	0.49	0	15,19,21	0.66	0
2	NAG	A	906	2	15,15,15	0.36	0	21,21,21	0.53	0
2	NAG	B	911	2	12,14,15	0.40	0	15,19,21	0.69	0
2	NAG	B	912	2	12,14,15	0.51	0	15,19,21	0.68	0
2	NAG	B	913	2	12,14,15	0.45	0	15,19,21	0.72	0
2	NAG	B	914	2	12,14,15	0.42	0	15,19,21	0.69	0
2	NAG	B	915	2	12,14,15	0.50	0	15,19,21	0.65	0
2	NAG	B	916	2	15,15,15	0.39	0	21,21,21	0.59	0
2	NAG	C	921	2	12,14,15	0.39	0	15,19,21	0.69	0
2	NAG	C	922	2	12,14,15	0.47	0	15,19,21	0.65	0
2	NAG	C	923	2	12,14,15	0.45	0	15,19,21	0.69	0
2	NAG	C	924	2	12,14,15	0.43	0	15,19,21	0.68	0
2	NAG	C	925	2	12,14,15	0.47	0	15,19,21	0.66	0
2	NAG	C	926	2	15,15,15	0.37	0	21,21,21	0.55	0
2	NAG	D	931	2	12,14,15	0.39	0	15,19,21	0.69	0
2	NAG	D	932	2	12,14,15	0.45	0	15,19,21	0.66	0
2	NAG	D	933	2	12,14,15	0.43	0	15,19,21	0.69	0
2	NAG	D	934	2	12,14,15	0.41	0	15,19,21	0.68	0
2	NAG	D	935	2	12,14,15	0.47	0	15,19,21	0.65	0
2	NAG	D	936	2	15,15,15	0.37	0	21,21,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	901	2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	902	2	-	0/6/23/26	0/1/1/1
2	NAG	A	903	2	-	0/6/23/26	0/1/1/1
2	NAG	A	904	2	-	0/6/23/26	0/1/1/1
2	NAG	A	905	2	-	0/6/23/26	0/1/1/1
2	NAG	A	906	2	-	0/6/26/26	0/1/1/1
2	NAG	B	911	2	-	0/6/23/26	0/1/1/1
2	NAG	B	912	2	-	0/6/23/26	0/1/1/1
2	NAG	B	913	2	-	0/6/23/26	0/1/1/1
2	NAG	B	914	2	-	0/6/23/26	0/1/1/1
2	NAG	B	915	2	-	0/6/23/26	0/1/1/1
2	NAG	B	916	2	-	0/6/26/26	0/1/1/1
2	NAG	C	921	2	-	0/6/23/26	0/1/1/1
2	NAG	C	922	2	-	0/6/23/26	0/1/1/1
2	NAG	C	923	2	-	0/6/23/26	0/1/1/1
2	NAG	C	924	2	-	0/6/23/26	0/1/1/1
2	NAG	C	925	2	-	0/6/23/26	0/1/1/1
2	NAG	C	926	2	-	0/6/26/26	0/1/1/1
2	NAG	D	931	2	-	0/6/23/26	0/1/1/1
2	NAG	D	932	2	-	0/6/23/26	0/1/1/1
2	NAG	D	933	2	-	0/6/23/26	0/1/1/1
2	NAG	D	934	2	-	0/6/23/26	0/1/1/1
2	NAG	D	935	2	-	0/6/23/26	0/1/1/1
2	NAG	D	936	2	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	A	699	-	4,4,4	0.92	0	6,6,6	0.31	0
3	PO4	B	696	-	4,4,4	0.89	0	6,6,6	0.31	0
3	PO4	C	697	-	4,4,4	0.89	0	6,6,6	0.31	0
3	PO4	C	701	-	4,4,4	0.88	0	6,6,6	0.31	0
3	PO4	D	698	-	4,4,4	0.90	0	6,6,6	0.31	0
3	PO4	D	700	-	4,4,4	0.89	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	699	-	-	0/0/0/0	0/0/0/0
3	PO4	B	696	-	-	0/0/0/0	0/0/0/0
3	PO4	C	697	-	-	0/0/0/0	0/0/0/0
3	PO4	C	701	-	-	0/0/0/0	0/0/0/0
3	PO4	D	698	-	-	0/0/0/0	0/0/0/0
3	PO4	D	700	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/345 (95%)	-0.43	0 100 100	20, 37, 54, 72	0
1	B	335/345 (97%)	-0.45	0 100 100	16, 36, 52, 66	0
1	C	332/345 (96%)	-0.46	1 (0%) 91 95	17, 36, 53, 69	0
1	D	335/345 (97%)	-0.40	2 (0%) 86 91	17, 38, 57, 65	0
All	All	1333/1380 (96%)	-0.44	3 (0%) 93 96	16, 37, 54, 72	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	17	PHE	2.7
1	D	18	THR	2.3
1	C	90	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	906	15/15	0.17	3.58	54,61,62,62	0
2	NAG	C	926	15/15	0.14	3.46	50,55,57,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	D	936	15/15	0.15	2.40	38,44,47,48	0
2	NAG	B	916	15/15	0.11	1.85	42,49,51,53	0
2	NAG	B	914	14/15	0.11	0.74	23,26,29,32	0
2	NAG	D	933	14/15	0.11	0.63	19,22,24,26	0
2	NAG	A	905	14/15	0.10	0.55	39,45,48,49	0
2	NAG	D	934	14/15	0.11	0.23	24,28,31,33	0
2	NAG	C	923	14/15	0.10	0.15	24,29,32,34	0
2	NAG	B	912	14/15	0.10	0.15	19,21,25,26	0
2	NAG	C	924	14/15	0.11	0.05	31,37,39,40	0
2	NAG	B	913	14/15	0.10	0.04	16,22,22,24	0
2	NAG	C	922	14/15	0.09	-0.03	22,26,31,31	0
2	NAG	A	903	14/15	0.10	-0.09	24,26,27,27	0
2	NAG	A	904	14/15	0.10	-0.21	27,30,35,38	0
2	NAG	C	925	14/15	0.10	-0.22	37,41,44,46	0
2	NAG	A	902	14/15	0.09	-0.29	22,25,27,29	0
2	NAG	D	932	14/15	0.10	-0.35	23,25,27,28	0
2	NAG	D	935	14/15	0.08	-1.02	32,35,38,40	0
2	NAG	B	911	14/15	0.08	-1.44	26,28,30,32	0
2	NAG	B	915	14/15	0.08	-1.49	30,36,39,39	0
2	NAG	A	901	14/15	0.07	-2.04	31,33,35,36	0
2	NAG	D	931	14/15	0.08	-2.88	24,31,35,37	0
2	NAG	C	921	14/15	0.07	-4.96	23,29,32,36	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PO4	A	699	5/5	0.12	3.14	20,21,22,25	0
3	PO4	D	698	5/5	0.10	-0.06	20,22,24,25	0
3	PO4	B	696	5/5	0.10	-0.12	27,28,28,29	0
3	PO4	C	697	5/5	0.10	-0.25	23,24,25,25	0
3	PO4	C	701	5/5	0.10	-0.37	44,44,46,46	0
3	PO4	D	700	5/5	0.09	-2.04	42,42,42,43	0

6.5 Other polymers ⓘ

There are no such residues in this entry.