



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:07 AM GMT

PDB ID : 3PIC
Title : Glucuronoyl Esterase catalytic domain (Cip2_GE) from Hypocrea jecorina
Authors : Pokkuluri, P.R.
Deposited on : 2010-11-05
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	375	
1	B	375	
1	C	375	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NAG	A	500	-	X
2	NAG	B	500	-	X
2	NAG	C	500	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9221 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cip2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	0	0
			2692	1696	461	520	15			
1	B	366	Total	C	N	O	S	0	0	0
			2730	1718	469	527	16			
1	C	362	Total	C	N	O	S	0	0	0
			2644	1667	450	512	15			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
A	87	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
A	88	THR	-	CLONING ARTIFACT	UNP Q7Z9N1
A	89	SER	-	CLONING ARTIFACT	UNP Q7Z9N1
B	86	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
B	87	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
B	88	THR	-	CLONING ARTIFACT	UNP Q7Z9N1
B	89	SER	-	CLONING ARTIFACT	UNP Q7Z9N1
C	86	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
C	87	GLN	-	CLONING ARTIFACT	UNP Q7Z9N1
C	88	THR	-	CLONING ARTIFACT	UNP Q7Z9N1
C	89	SER	-	CLONING ARTIFACT	UNP Q7Z9N1

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	375	Total	O	0	0
			375	375		
3	B	469	Total	O	0	0
			469	469		
3	C	269	Total	O	0	0
			269	269		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cip2

Chain A:




• Molecule 1: Cip2

Chain B:



• Molecule 1: Cip2

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.26Å 82.01Å 185.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.20 – 1.90 31.20 – 1.88	Depositor EDS
% Data completeness (in resolution range)	90.1 (31.20-1.90) 88.9 (31.20-1.88)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.88Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.195 , 0.234 0.196 , 0.234	Depositor DCC
R_{free} test set	8748 reflections (10.89%)	DCC
Wilson B-factor (Å ²)	12.1	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.1	EDS
Estimated twinning fraction	0.044 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 91298 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9221	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2758	0.58	0/3759
1	B	0.33	0/2796	0.58	0/3805
1	C	0.30	0/2709	0.56	0/3696
All	All	0.32	0/8263	0.57	0/11260

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2692	0	2565	29	0
1	B	2730	0	2619	23	0
1	C	2644	0	2473	35	0
2	A	14	0	13	4	0
2	B	14	0	13	0	0
2	C	14	0	13	0	0
3	A	375	0	0	3	0
3	B	469	0	0	5	0
3	C	269	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9221	0	7696	87	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (87) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:447:ASN:HD21	2:A:500:NAG:C1	1.72	1.01
1:A:447:ASN:ND2	2:A:500:NAG:C1	2.37	0.88
1:C:182:THR:HG22	1:C:210:MET:HB2	1.67	0.77
1:C:330:ILE:HD13	1:C:344:VAL:HG11	1.69	0.73
1:C:346:GLN:O	1:C:349:VAL:HG22	1.91	0.71
1:A:297:THR:HG22	1:A:299:PRO:HD3	1.75	0.69
1:A:327:SER:HA	1:A:344:VAL:CG1	2.24	0.67
1:B:346:GLN:O	1:B:349:VAL:HG22	1.95	0.67
1:C:182:THR:CG2	1:C:210:MET:HB2	2.26	0.66
1:A:279:ARG:HA	1:A:282:LYS:HE2	1.78	0.65
1:C:268:ASP:OD1	1:C:270:THR:HG22	1.97	0.65
1:A:327:SER:HA	1:A:344:VAL:HG13	1.79	0.64
1:A:99:LEU:HD12	1:A:100:PRO:HD2	1.80	0.63
1:A:427:GLN:OE1	1:A:435:THR:HG21	2.02	0.60
1:C:256:ILE:O	1:C:260:GLU:HG3	2.02	0.59
1:C:460:SER:HB3	3:C:935:HOH:O	2.03	0.59
1:C:166:ILE:N	1:C:166:ILE:HD12	2.17	0.59
1:B:99:LEU:HB2	3:B:738:HOH:O	2.02	0.59
1:A:108:ASN:HD22	1:A:108:ASN:C	2.05	0.58
1:B:421:GLN:HG3	3:B:613:HOH:O	2.04	0.57
1:C:122:LYS:HE3	3:C:882:HOH:O	2.03	0.57
1:A:346:GLN:O	1:A:349:VAL:HG22	2.05	0.57
1:C:426:VAL:HG13	1:C:430:LEU:HD12	1.87	0.56
1:C:279:ARG:HG2	1:C:280:ASN:N	2.19	0.56
1:C:347:VAL:N	1:C:348:PRO:HD2	2.21	0.55
1:A:379:GLN:CD	2:A:500:NAG:H82	2.27	0.55
1:C:423:THR:O	1:C:427:GLN:HG3	2.07	0.55
1:B:245:MET:HE3	1:B:335:TRP:O	2.07	0.54
1:B:100:PRO:HG2	1:B:103:ILE:CG2	2.37	0.54
1:B:330:ILE:HD13	1:B:344:VAL:HG11	1.90	0.54
1:C:268:ASP:CG	1:C:270:THR:HG22	2.29	0.53
1:B:244:ALA:HB3	1:B:335:TRP:HA	1.90	0.53
1:A:324:GLN:HB2	1:A:375:TRP:CE2	2.44	0.53
1:B:250:TRP:O	1:B:254:ARG:HG2	2.09	0.53
1:B:297:THR:HG22	1:B:299:PRO:HD3	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:324:GLN:HB2	1:C:375:TRP:CE2	2.45	0.52
1:C:250:TRP:O	1:C:254:ARG:HG2	2.09	0.51
1:A:295:VAL:HG12	1:A:429:PHE:HB3	1.92	0.51
1:A:295:VAL:HG13	3:A:480:HOH:O	2.10	0.51
1:C:426:VAL:O	1:C:430:LEU:HB2	2.11	0.50
1:B:428:LYS:HG2	1:B:429:PHE:CE1	2.48	0.49
1:A:326:ALA:C	1:A:344:VAL:HG22	2.32	0.49
1:C:214:ASN:ND2	1:C:217:ASN:ND2	2.60	0.49
1:C:377:GLY:N	1:C:378:PRO:HD3	2.27	0.49
1:A:327:SER:CA	1:A:344:VAL:HG13	2.43	0.48
1:C:239:SER:HB2	3:C:542:HOH:O	2.13	0.48
1:A:221:GLN:OE1	1:A:279:ARG:HD2	2.13	0.48
1:C:158:SER:O	1:C:164:LEU:HD12	2.14	0.47
1:B:189:ALA:HA	1:B:190:PRO:C	2.35	0.47
1:C:312:ILE:HG21	1:C:380:SER:HA	1.96	0.47
1:C:297:THR:HG22	1:C:299:PRO:HD3	1.97	0.47
1:B:107:SER:HA	1:B:349:VAL:HA	1.98	0.46
1:A:327:SER:HA	1:A:344:VAL:HG11	1.97	0.46
1:C:268:ASP:OD2	1:C:270:THR:HG22	2.16	0.46
1:A:421:GLN:HG3	3:A:612:HOH:O	2.16	0.45
1:B:292:LYS:HD2	1:B:292:LYS:N	2.32	0.45
1:C:155:LEU:HD23	1:C:155:LEU:C	2.37	0.45
1:B:436:ASN:HB2	3:B:886:HOH:O	2.17	0.45
1:B:189:ALA:HB2	1:B:191:TYR:CE2	2.53	0.44
1:B:256:ILE:O	1:B:260:GLU:HG3	2.17	0.44
1:A:256:ILE:O	1:A:260:GLU:HG3	2.18	0.44
1:B:245:MET:HE1	1:B:336:PHE:CE1	2.53	0.44
1:A:291:GLU:HG3	1:A:294:ILE:HG13	1.99	0.43
1:A:305:GLY:O	1:A:354:HIS:HE1	2.02	0.43
1:B:310:TRP:CE3	1:B:326:ALA:HB2	2.53	0.43
1:A:250:TRP:O	1:A:254:ARG:HG2	2.19	0.43
1:C:366:LEU:HD23	1:C:401:MET:HG3	2.01	0.43
1:B:366:LEU:O	1:B:401:MET:HA	2.19	0.42
1:A:136:MET:O	1:A:140:ILE:HG13	2.20	0.42
1:C:155:LEU:HA	1:C:167:ASN:O	2.19	0.42
1:C:421:GLN:HB3	1:C:439:ILE:HD13	2.01	0.42
1:C:184:PRO:HD3	1:C:209:ALA:HA	2.01	0.42
1:A:203:PRO:HG2	1:A:419:GLN:HE22	1.85	0.42
1:C:417:ASN:HB2	3:C:1027:HOH:O	2.20	0.42
1:C:231:LYS:HA	1:C:234:ASP:OD2	2.20	0.41
1:A:447:ASN:CG	2:A:500:NAG:C1	2.89	0.41
1:B:224:THR:HG22	3:B:754:HOH:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:105:LEU:HD22	1:B:351:PRO:HD3	2.02	0.41
1:A:295:VAL:HG11	3:A:61:HOH:O	2.20	0.41
1:B:282:LYS:HG2	3:B:581:HOH:O	2.20	0.41
1:C:132:ARG:O	1:C:136:MET:HG3	2.21	0.41
1:A:244:ALA:HB3	1:A:335:TRP:HA	2.02	0.40
1:A:330:ILE:C	1:A:330:ILE:HD12	2.41	0.40
1:C:166:ILE:HG13	1:C:258:ALA:CB	2.52	0.40
1:C:254:ARG:HA	1:C:254:ARG:NE	2.37	0.40
1:C:366:LEU:O	1:C:401:MET:HA	2.21	0.40
1:B:136:MET:O	1:B:140:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/375 (96%)	348 (97%)	12 (3%)	0	100	100
1	B	364/375 (97%)	353 (97%)	11 (3%)	0	100	100
1	C	360/375 (96%)	344 (96%)	16 (4%)	0	100	100
All	All	1084/1125 (96%)	1045 (96%)	39 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/295 (95%)	278 (99%)	3 (1%)	84	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	288/295 (98%)	283 (98%)	5 (2%)	73	68
1	C	267/295 (90%)	262 (98%)	5 (2%)	69	63
All	All	836/885 (94%)	823 (98%)	13 (2%)	75	70

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	344	VAL
1	A	440	PHE
1	B	216	ASP
1	B	279	ARG
1	B	401	MET
1	B	428	LYS
1	B	440	PHE
1	C	188	THR
1	C	279	ARG
1	C	401	MET
1	C	417	ASN
1	C	445	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	419	GLN
1	A	447	ASN
1	B	322	ASN
1	B	419	GLN
1	B	448	GLN
1	C	217	ASN
1	C	240	HIS
1	C	341	ASN
1	C	417	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	500	-	12,14,15	0.45	0	15,19,21	0.60	0
2	NAG	B	500	1	12,14,15	0.40	0	15,19,21	0.72	0
2	NAG	C	500	1	12,14,15	0.40	0	15,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	500	-	-	0/6/23/26	0/1/1/1
2	NAG	B	500	1	-	0/6/23/26	0/1/1/1
2	NAG	C	500	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/375 (96%)	-0.27	1 (0%) 91 93	4, 11, 23, 33	0
1	B	366/375 (97%)	-0.34	1 (0%) 91 93	3, 9, 18, 32	0
1	C	362/375 (96%)	0.16	10 (2%) 50 51	7, 19, 32, 37	0
All	All	1090/1125 (96%)	-0.15	12 (1%) 77 79	3, 12, 28, 37	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	431	LEU	3.4
1	A	188	THR	3.1
1	C	432	GLY	2.9
1	C	435	THR	2.7
1	B	95	THR	2.6
1	C	261	LEU	2.6
1	C	157	ALA	2.5
1	C	172	GLY	2.3
1	C	171	ALA	2.3
1	C	184	PRO	2.1
1	C	265	ALA	2.0
1	C	460	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	B	500	14/15	0.14	3.20	12,17,19,20	0
2	NAG	C	500	14/15	0.17	2.29	25,30,33,35	0
2	NAG	A	500	14/15	0.13	2.11	14,17,21,24	0

6.5 Other polymers

There are no such residues in this entry.