



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 03:50 PM GMT

PDB ID : 3PMZ
Title : Crystal Structure of the Complex of Acetylcholine Binding Protein and d-tubocurarine
Authors : Talley, T.T.; Harel, M.; Yamauchi, J.G.; Radic, Z.; Hansen, S.; Huxford, T.; Taylor, P.W.
Deposited on : 2010-11-18
Resolution : 2.44 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2989 (2.48-2.40)
Clashscore	79885	3698 (2.48-2.40)
Ramachandran outliers	78287	3639 (2.48-2.40)
Sidechain outliers	78261	3640 (2.48-2.40)
RSRZ outliers	66119	2993 (2.48-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	227	
1	B	227	
1	C	227	
1	D	227	
1	E	227	
1	F	227	
1	G	227	
1	H	227	
1	I	227	
1	J	227	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	MG	A	221	-	X
2	MG	B	220	-	X
2	MG	B	221	-	X
2	MG	D	220	-	X
2	MG	E	221	-	X
2	MG	E	222	-	X
2	MG	F	221	-	X
2	MG	I	220	-	X
2	MG	I	221	-	X
2	MG	J	220	-	X
3	TUB	E	220	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17405 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Soluble acetylcholine receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1700	1071	280	340	9			
1	B	211	Total	C	N	O	S	0	0	0
			1685	1061	278	338	8			
1	C	215	Total	C	N	O	S	0	3	0
			1744	1099	290	346	9			
1	D	214	Total	C	N	O	S	0	2	0
			1725	1087	288	341	9			
1	E	211	Total	C	N	O	S	0	0	0
			1686	1064	278	335	9			
1	F	207	Total	C	N	O	S	0	0	0
			1648	1041	270	328	9			
1	G	211	Total	C	N	O	S	0	0	0
			1683	1061	276	338	8			
1	H	212	Total	C	N	O	S	0	0	0
			1693	1066	279	339	9			
1	I	212	Total	C	N	O	S	0	0	0
			1692	1067	279	337	9			
1	J	213	Total	C	N	O	S	0	0	0
			1697	1070	278	340	9			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
A	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
A	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
A	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
A	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
A	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
A	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
A	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
B	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
B	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
B	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
B	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
B	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
B	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
B	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
C	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
C	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
C	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
C	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
C	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
C	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
C	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
C	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
D	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
D	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
D	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
D	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
D	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
D	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
D	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
D	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
E	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
E	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
E	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
E	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
E	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
E	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
E	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
E	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
F	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
F	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
F	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
F	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
F	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
F	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
F	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
F	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
G	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
G	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
G	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
G	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
G	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
G	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
G	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
H	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
H	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
H	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
H	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
H	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
H	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
H	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
H	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
I	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
I	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
I	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
I	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
I	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
I	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
I	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
I	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8
J	-7	ASP	-	EXPRESSION TAG	UNP Q8WSF8
J	-6	TYR	-	EXPRESSION TAG	UNP Q8WSF8
J	-5	LYS	-	EXPRESSION TAG	UNP Q8WSF8
J	-4	ASP	-	EXPRESSION TAG	UNP Q8WSF8
J	-3	ASP	-	EXPRESSION TAG	UNP Q8WSF8
J	-2	ASP	-	EXPRESSION TAG	UNP Q8WSF8
J	-1	LYS	-	EXPRESSION TAG	UNP Q8WSF8
J	0	LEU	-	EXPRESSION TAG	UNP Q8WSF8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Mg 1 1	0	0
2	J	1	Total Mg 1 1	0	0
2	D	3	Total Mg 3 3	0	0
2	E	2	Total Mg 2 2	0	0
2	B	2	Total Mg 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	I	2	Total	Mg	0	0
			2	2		
2	C	1	Total	Mg	0	0
			1	1		
2	A	2	Total	Mg	0	0
			2	2		
2	F	2	Total	Mg	0	0
			2	2		

- Molecule 3 is (1BETA,1'ALPHA)-7',12'-DIHYDROXY-6,6'-DIMETHOXY-2,2',2'-TRIMETHYLTUBOCURARAN-2'-IUM (three-letter code: TUB) (formula: C₃₇H₄₁N₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	E	1	Total	C	N	O	0	0
			45	37	2	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	58	Total	O	0	0
			58	58		
4	B	41	Total	O	0	0
			41	41		
4	C	57	Total	O	0	0
			57	57		
4	D	56	Total	O	0	0
			56	56		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	71	Total 71	O 71	0	0
4	F	22	Total 22	O 22	0	0
4	G	14	Total 14	O 14	0	0
4	H	11	Total 11	O 11	0	0
4	I	22	Total 22	O 22	0	0
4	J	39	Total 39	O 39	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Soluble acetylcholine receptor

Chain A: 




- Molecule 1: Soluble acetylcholine receptor

Chain B: 




- Molecule 1: Soluble acetylcholine receptor

Chain C: 




- Molecule 1: Soluble acetylcholine receptor

Chain D: 




- Molecule 1: Soluble acetylcholine receptor

Chain E:




- Molecule 1: Soluble acetylcholine receptor

Chain F:




- Molecule 1: Soluble acetylcholine receptor

Chain G:




- Molecule 1: Soluble acetylcholine receptor

Chain H:




- Molecule 1: Soluble acetylcholine receptor

Chain I: 



● Molecule 1: Soluble acetylcholine receptor

Chain J: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	142.74Å 194.04Å 101.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	114.71 – 2.44 48.51 – 2.44	Depositor EDS
% Data completeness (in resolution range)	96.8 (114.71-2.44) 96.9 (48.51-2.44)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.228 , 0.294 0.226 , 0.289	Depositor DCC
R_{free} test set	5105 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	40.7	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 22.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 101833 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17405	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TUB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	1/1741 (0.1%)	0.87	2/2373 (0.1%)
1	B	0.84	0/1724	0.84	0/2348
1	C	0.94	0/1793	0.93	3/2439 (0.1%)
1	D	0.96	0/1772	0.93	1/2413 (0.0%)
1	E	0.93	0/1726	0.92	1/2350 (0.0%)
1	F	0.77	0/1689	0.86	0/2304
1	G	0.77	0/1722	0.85	2/2345 (0.1%)
1	H	0.76	0/1732	0.84	2/2358 (0.1%)
1	I	0.80	0/1733	0.79	0/2362
1	J	0.85	0/1736	0.89	5/2366 (0.2%)
All	All	0.85	1/17368 (0.0%)	0.87	16/23658 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127	CYS	CB-SG	5.07	1.90	1.82

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	0	LEU	CA-CB-CG	6.13	129.40	115.30
1	C	122	ARG	NE-CZ-NH1	5.74	123.17	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	195	TYR	N-CA-C	-5.67	95.70	111.00
1	A	97	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	G	51	ASP	CB-CG-OD1	5.51	123.26	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	17	SER	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1700	0	1625	20	0
1	B	1685	0	1608	24	0
1	C	1744	0	1677	41	0
1	D	1725	0	1664	32	0
1	E	1686	0	1616	22	0
1	F	1648	0	1576	60	0
1	G	1683	0	1608	75	0
1	H	1693	0	1617	69	0
1	I	1692	0	1621	49	0
1	J	1697	0	1616	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	3	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	1	0	0	0	0
2	I	2	0	0	0	0
2	J	1	0	0	0	0
3	E	45	0	40	6	0
4	A	58	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	41	0	0	0	0
4	C	57	0	0	0	0
4	D	56	0	0	1	0
4	E	71	0	0	0	0
4	F	22	0	0	0	0
4	G	14	0	0	2	0
4	H	11	0	0	0	0
4	I	22	0	0	2	0
4	J	39	0	0	0	0
All	All	17405	0	16268	374	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

The worst 5 of 374 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:F:184:GLN:NE2	1:F:199:ASN:HB2	1.56	1.19
1:I:59:ARG:HD3	1:I:116:MET:CE	1.85	1.06
1:F:173:LYS:HE3	1:G:45:SER:O	1.60	1.00
1:G:105:GLN:HE21	1:G:105:GLN:HA	1.30	0.97
1:F:38:GLN:HB3	1:G:126:MET:HE1	1.45	0.95

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/227 (93%)	202 (96%)	6 (3%)	3 (1%)	16	21
1	B	207/227 (91%)	201 (97%)	6 (3%)	0	100	100
1	C	214/227 (94%)	204 (95%)	8 (4%)	2 (1%)	25	33
1	D	214/227 (94%)	206 (96%)	6 (3%)	2 (1%)	25	33
1	E	207/227 (91%)	201 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	205/227 (90%)	178 (87%)	18 (9%)	9 (4%)	4	2
1	G	207/227 (91%)	174 (84%)	25 (12%)	8 (4%)	5	3
1	H	208/227 (92%)	183 (88%)	22 (11%)	3 (1%)	16	21
1	I	210/227 (92%)	196 (93%)	11 (5%)	3 (1%)	16	21
1	J	211/227 (93%)	194 (92%)	15 (7%)	2 (1%)	25	33
All	All	2094/2270 (92%)	1939 (93%)	123 (6%)	32 (2%)	15	19

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	18	PRO
1	F	3	GLN
1	F	134	SER
1	F	166	SER
1	G	150	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/205 (95%)	184 (95%)	10 (5%)	32	49
1	B	192/205 (94%)	180 (94%)	12 (6%)	25	37
1	C	199/205 (97%)	183 (92%)	16 (8%)	17	25
1	D	197/205 (96%)	182 (92%)	15 (8%)	19	28
1	E	192/205 (94%)	183 (95%)	9 (5%)	36	53
1	F	188/205 (92%)	173 (92%)	15 (8%)	17	25
1	G	192/205 (94%)	166 (86%)	26 (14%)	6	6
1	H	193/205 (94%)	177 (92%)	16 (8%)	16	23
1	I	193/205 (94%)	174 (90%)	19 (10%)	12	16
1	J	193/205 (94%)	182 (94%)	11 (6%)	29	43
All	All	1933/2050 (94%)	1784 (92%)	149 (8%)	19	27

5 of 149 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	77	ASP
1	G	45	SER
1	I	205	ARG
1	F	104	PRO
1	F	203	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	38	GLN
1	G	15	ASN
1	J	74	ASN
1	F	184	GLN
1	G	57	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 16 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TUB	E	220	-	51,51,51	1.73	8 (15%)	77,77,77	4.24	23 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TUB	E	220	-	-	0/20/49/49	0/0/7/7

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	220	TUB	O21-C22	6.44	1.54	1.39
3	E	220	TUB	O37-C1	4.73	1.44	1.37
3	E	220	TUB	C4-C7	-3.98	1.47	1.52
3	E	220	TUB	C12-C11	3.41	1.45	1.40
3	E	220	TUB	O38-C29	3.33	1.43	1.36

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	220	TUB	O21-C22-C29	17.93	152.85	116.33
3	E	220	TUB	C35-C26-C25	-15.18	93.50	120.41
3	E	220	TUB	O21-C22-C25	-14.64	81.68	121.61
3	E	220	TUB	C35-C26-C27	12.91	146.23	120.90
3	E	220	TUB	C5-C4-C7	8.67	129.81	121.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/227 (93%)	-0.14	5 (2%) 57 55	17, 31, 56, 71	0
1	B	211/227 (92%)	-0.12	5 (2%) 56 54	19, 33, 61, 72	0
1	C	215/227 (94%)	-0.20	3 (1%) 72 71	17, 30, 61, 84	0
1	D	214/227 (94%)	-0.11	3 (1%) 72 71	14, 25, 60, 73	0
1	E	211/227 (92%)	-0.25	2 (0%) 81 81	13, 25, 49, 76	0
1	F	207/227 (91%)	0.28	11 (5%) 25 23	31, 49, 66, 75	0
1	G	211/227 (92%)	0.56	16 (7%) 14 12	32, 57, 79, 84	0
1	H	212/227 (93%)	0.43	13 (6%) 21 18	30, 56, 77, 87	0
1	I	212/227 (93%)	-0.10	3 (1%) 72 71	23, 40, 64, 77	0
1	J	213/227 (93%)	-0.11	6 (2%) 50 48	22, 36, 61, 79	0
All	All	2119/2270 (93%)	0.02	67 (3%) 45 43	13, 37, 69, 87	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	19	MET	5.6
1	G	191	CYS	4.9
1	G	137	GLY	4.8
1	C	-6	TYR	4.5
1	B	189	SER	4.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MG	E	222	1/1	0.62	25.88	36,36,36,36	0
2	MG	B	220	1/1	0.31	13.43	44,44,44,44	0
2	MG	I	220	1/1	0.29	10.20	43,43,43,43	0
2	MG	A	221	1/1	0.30	9.02	43,43,43,43	0
2	MG	I	221	1/1	0.38	6.04	52,52,52,52	0
2	MG	E	221	1/1	0.31	5.49	45,45,45,45	0
2	MG	J	220	1/1	0.33	5.37	46,46,46,46	0
2	MG	F	221	1/1	0.29	4.24	52,52,52,52	0
2	MG	B	221	1/1	0.27	3.01	53,53,53,53	0
3	TUB	E	220	45/45	0.19	2.21	15,20,26,28	0
2	MG	D	220	1/1	0.16	2.01	49,49,49,49	0
2	MG	D	222	1/1	0.21	1.98	48,48,48,48	0
2	MG	A	220	1/1	0.23	1.74	43,43,43,43	0
2	MG	C	220	1/1	0.26	1.36	40,40,40,40	0
2	MG	D	221	1/1	0.15	0.92	57,57,57,57	0
2	MG	F	220	1/1	0.20	0.29	54,54,54,54	0
2	MG	G	220	1/1	0.12	-1.85	55,55,55,55	0

6.5 Other polymers ⓘ

There are no such residues in this entry.