



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:50 PM GMT

PDB ID : 2PQR
Title : Crystal structure of yeast Fis1 complexed with a fragment of yeast Caf4
Authors : Zhang, Y.; Chan, D.C.
Deposited on : 2007-05-02
Resolution : 1.88 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	5260 (1.90-1.86)
Clashscore	79885	6268 (1.90-1.86)
Ramachandran outliers	78287	6195 (1.90-1.86)
Sidechain outliers	78261	6196 (1.90-1.86)
RSRZ outliers	66119	5262 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	129	
1	B	129	
2	C	64	
2	D	64	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	AU	B	301[A]	-	X
3	AU	B	301[B]	-	X
3	AU	D	302	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2666 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondria fission 1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	98	Total	C	N	O	S	4	0	0
			809	518	134	154	3			
1	B	124	Total	C	N	O	S	0	3	0
			1037	664	173	196	4			

- Molecule 2 is a protein called WD repeat protein YKR036C.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	40	Total	C	N	O	0	0	0
			323	209	56	58			
2	D	45	Total	C	N	O	0	0	0
			353	227	59	67			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	77	GLY	-	EXPRESSION TAG	UNP P36130
C	78	SER	-	EXPRESSION TAG	UNP P36130
C	79	HIS	-	EXPRESSION TAG	UNP P36130
C	80	MET	-	EXPRESSION TAG	UNP P36130
C	110	GLN	HIS	VARIANT	UNP P36130
C	111	ARG	GLY	VARIANT	UNP P36130
D	77	GLY	-	EXPRESSION TAG	UNP P36130
D	78	SER	-	EXPRESSION TAG	UNP P36130
D	79	HIS	-	EXPRESSION TAG	UNP P36130
D	80	MET	-	EXPRESSION TAG	UNP P36130
D	110	GLN	HIS	VARIANT	UNP P36130
D	111	ARG	GLY	VARIANT	UNP P36130

- Molecule 3 is GOLD ION (three-letter code: AU) (formula: Au).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 2	Au 2	0	1
3	D	1	Total 1	Au 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	15	Total 15	O 15	0	0
4	B	85	Total 85	O 85	0	0
4	C	14	Total 14	O 14	0	0
4	D	27	Total 27	O 27	0	0

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitochondria fission 1 protein

Chain A: 




- Molecule 1: Mitochondria fission 1 protein

Chain B: 




- Molecule 2: WD repeat protein YKR036C

Chain C: 



- Molecule 2: WD repeat protein YKR036C

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.41Å 46.95Å 64.47Å 90.00° 99.21° 90.00°	Depositor
Resolution (Å)	40.40 – 1.88 40.40 – 1.88	Depositor EDS
% Data completeness (in resolution range)	92.9 (40.40-1.88) 93.0 (40.40-1.88)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.3.0026	Depositor
R, R_{free}	0.208 , 0.241 0.208 , 0.241	Depositor DCC
R_{free} test set	1333 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.274	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 70.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 28916 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2666	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	1/827 (0.1%)	0.64	1/1120 (0.1%)
1	B	0.85	1/1066 (0.1%)	0.81	0/1439
2	C	1.45	3/329 (0.9%)	1.33	3/443 (0.7%)
2	D	0.64	0/361	0.86	1/490 (0.2%)
All	All	0.87	5/2583 (0.2%)	0.86	5/3492 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	111	ARG	CZ-NH1	19.52	1.58	1.33
1	A	11	LYS	CB-CG	-12.59	1.18	1.52
2	C	111	ARG	NE-CZ	10.05	1.46	1.33
1	B	71	GLU	CB-CG	-6.04	1.40	1.52
2	C	121	LYS	C-O	5.10	1.33	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	111	ARG	NE-CZ-NH2	-20.57	110.01	120.30
2	C	111	ARG	NE-CZ-NH1	12.53	126.56	120.30
2	D	96	ASP	N-CA-C	-9.12	86.37	111.00
2	C	111	ARG	CD-NE-CZ	-6.13	115.02	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	LYS	CA-CB-CG	6.01	126.63	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	95	ALA	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	809	0	796	11	0
1	B	1037	0	1044	10	0
2	C	323	0	326	10	0
2	D	353	0	347	11	0
3	B	2	0	0	0	0
3	D	1	0	0	0	0
4	A	15	0	0	0	0
4	B	85	0	0	0	0
4	C	14	0	0	1	0
4	D	27	0	0	0	0
All	All	2666	0	2513	32	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (32) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:39[B]:ILE:HG23	1:B:68[B]:ILE:HD12	1.70	0.74
2:C:133:THR:HG23	2:D:136:ILE:HD13	1.69	0.73
1:A:12:ASP:OD2	2:C:131:LYS:NZ	2.27	0.65
2:C:136:ILE:HD11	2:D:136:ILE:CG2	2.27	0.65
2:C:136:ILE:HD11	2:D:136:ILE:HG22	1.81	0.63
2:D:118:ALA:HB3	2:D:119:ALA:HA	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:67:ASP:O	1:B:71:GLU:HG2	2.00	0.60
1:A:40:GLN:HE21	1:A:44:ASN:HD21	1.51	0.58
1:B:50:ILE:HD11	1:B:65:LEU:HD22	1.84	0.58
1:A:21:GLN:NE2	2:C:138:GLN:HE22	2.06	0.54
1:A:66:THR:HG23	2:C:107:LEU:HD12	1.90	0.53
1:B:50:ILE:CD1	1:B:65:LEU:HD22	2.40	0.51
2:D:117:GLY:O	2:D:118:ALA:HB3	2.10	0.50
1:B:5:ASP:HB3	1:B:7:TRP:CE2	2.46	0.50
1:B:123:LYS:O	1:B:127:GLU:HG2	2.12	0.50
1:A:73:GLU:HA	1:A:76:ARG:HG3	1.94	0.49
2:C:108:ASP:HB3	2:C:111:ARG:HB3	1.95	0.49
1:B:110:ASN:HD22	1:B:110:ASN:C	2.16	0.48
4:C:148:HOH:O	2:D:136:ILE:HD12	2.14	0.48
2:D:117:GLY:O	2:D:118:ALA:CB	2.62	0.47
2:C:97:SER:OG	2:C:98:ALA:N	2.48	0.46
2:D:117:GLY:C	2:D:119:ALA:HB2	2.36	0.45
1:A:40:GLN:HE21	1:A:44:ASN:ND2	2.12	0.45
1:A:49:LEU:O	1:A:52:SER:HB3	2.18	0.44
1:A:50:ILE:HD11	1:A:65:LEU:HD22	2.00	0.44
2:C:133:THR:CG2	2:D:136:ILE:HD13	2.45	0.43
1:B:10:LEU:HD23	1:B:85:ILE:HG23	2.01	0.42
1:B:77:ARG:HA	2:D:114:LEU:HD11	2.02	0.42
1:A:69:TYR:HE1	1:A:80:LEU:HG	1.84	0.42
1:A:7:TRP:HB3	1:A:81:TYR:CE2	2.55	0.41
1:A:69:TYR:CD2	2:C:107:LEU:HD11	2.56	0.41
1:B:73:GLU:HG2	2:D:115:PRO:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/129 (74%)	94 (98%)	2 (2%)	0	100	100
1	B	125/129 (97%)	124 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	36/64 (56%)	33 (92%)	3 (8%)	0	100	100
2	D	43/64 (67%)	40 (93%)	1 (2%)	2 (5%)	4	0
All	All	300/386 (78%)	291 (97%)	7 (2%)	2 (1%)	30	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	118	ALA
2	D	95	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/116 (75%)	83 (95%)	4 (5%)	37	21
1	B	114/116 (98%)	112 (98%)	2 (2%)	71	63
2	C	35/53 (66%)	29 (83%)	6 (17%)	3	0
2	D	37/53 (70%)	37 (100%)	0	100	100
All	All	273/338 (81%)	261 (96%)	12 (4%)	39	23

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	31	SER
1	A	65	LEU
1	A	97	LYS
1	B	65	LEU
1	B	110	ASN
2	C	100	THR
2	C	110	GLN
2	C	114	LEU
2	C	131	LYS
2	C	136	ILE
2	C	139	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	21	GLN
1	A	44	ASN
1	B	27	GLN
1	B	106	HIS
1	B	110	ASN
1	B	112	GLN
2	C	110	GLN
2	D	138	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	98/129 (75%)	0.38	4 (4%) 35 35	36, 43, 57, 63	1 (1%)
1	B	124/129 (96%)	-0.11	0 100 100	19, 24, 31, 36	0
2	C	40/64 (62%)	0.87	6 (15%) 3 3	29, 44, 61, 61	0
2	D	45/64 (70%)	0.41	5 (11%) 6 5	25, 29, 44, 54	0
All	All	307/386 (79%)	0.25	15 (4%) 28 28	19, 32, 57, 63	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	115	PRO	4.2
1	A	99	TYR	3.9
2	D	121	LYS	3.9
2	C	122	ASN	3.9
2	C	108	ASP	2.8
2	C	121	LYS	2.7
2	D	119	ALA	2.7
1	A	19	PRO	2.7
2	D	122	ASN	2.2
1	A	18	TYR	2.2
2	C	113	PRO	2.2
2	D	118	ALA	2.1
2	D	117	GLY	2.1
2	C	110	GLN	2.0
1	A	102	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	AU	B	301[B]	1/1	0.13	2.60	44,44,44,44	1
3	AU	B	301[A]	1/1	0.13	2.58	41,41,41,41	1
3	AU	D	302	1/1	0.15	2.54	50,50,50,50	1

6.5 Other polymers

There are no such residues in this entry.