



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 04:14 PM GMT

PDB ID : 3PSI
Title : Crystal Structure of the Spt6 core domain from *Saccharomyces cerevisiae*,
Form Spt6(239-1451)
Authors : Close, D.; Hill, C.P.; Johnson, S.J.
Deposited on : 2010-12-01
Resolution : 3.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1341 (3.40-3.20)
Clashscore	79885	1696 (3.40-3.20)
Ramachandran outliers	78287	1664 (3.40-3.20)
Sidechain outliers	78261	1662 (3.40-3.20)
RSRZ outliers	66119	1342 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	1219	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 13677 atoms, of which 6800 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription elongation factor SPT6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	843	Total	C	H	N	O	S	0	0	0
			13677	4370	6800	1164	1326	17			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	233	GLY	-	EXPRESSION TAG	UNP P23615
A	234	ILE	-	EXPRESSION TAG	UNP P23615
A	235	ASP	-	EXPRESSION TAG	UNP P23615
A	236	PRO	-	EXPRESSION TAG	UNP P23615
A	237	PHE	-	EXPRESSION TAG	UNP P23615
A	238	THR	-	EXPRESSION TAG	UNP P23615

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcription elongation factor SPT6

Chain A: 



PHE GLN HIS ILE ASP ILE GLN GLU LEU LEU LYS GLY ASN PRO LEU ALA LEU LEU VAL ASP ASN GLN LYS TYR ASN ASP LEU ASP GLN ILE ILE VAL VAL GLU TYR LEU LEU GLN ASN LYS VAL ARG LEU LEU LEU ASN LYS THR PHE LYS SER SER SER LYS THR XYS

LYS ASP VAL VAL VAL PHE ILE GLU ASP TYR ARG SER VAL ASN PRO ASN LYS SER VAL TYR PHE SER LEU ASN HIS ASP ASN PRO GLY TRP PHE TYR MET PHE PHE LYS ILE ASN ALA ASN SER LYS TYR THR TRP ASN VAL LYS LEU THR THR ASN THR GLY TYR PHE LEU VAL ASN

TYR ASN TYR PRO SER VAL ILE GLN LEU CYS ASN GLY PHE LYS THR LEU LEU LYS SER ASN SER SER LYS ASN ARG MET ASN ASN TYR ARG

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.75Å 118.75Å 214.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.67 – 3.30 45.67 – 3.29	Depositor EDS
% Data completeness (in resolution range)	99.4 (45.67-3.30) 99.4 (45.67-3.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 3.32Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.265 , 0.308 0.250 , 0.339	Depositor DCC
R_{free} test set	2835 reflections (10.50%)	DCC
Wilson B-factor (Å ²)	111.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 110.0	EDS
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 27006 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13677	wwPDB-VP
Average B, all atoms (Å ²)	170.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/7015	0.77	2/9487 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1020	ARG	NE-CZ-NH1	-7.92	116.34	120.30
1	A	1101	LEU	CB-CG-CD1	-5.88	101.00	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6877	6800	1	222	0
All	All	6877	6800	1	222	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 222 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:980:ILE:HD12	1:A:981:THR:N	1.84	0.91

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:601:LEU:HD11	1:A:658:HIS:HB2	1.61	0.82
1:A:980:ILE:HD12	1:A:981:THR:H	1.41	0.81
1:A:641:PHE:CE1	1:A:713:LEU:HD23	2.22	0.74
1:A:653:LEU:HD12	1:A:653:LEU:C	2.07	0.74

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	831/1219 (68%)	760 (92%)	58 (7%)	13 (2%)	14 64

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1117	GLY
1	A	742	VAL
1	A	551	SER
1	A	628	ASN
1	A	762	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	762/1108 (69%)	759 (100%)	3 (0%)	95 99

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	689	PHE
1	A	726	ARG
1	A	1204	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	828	GLN
1	A	907	HIS
1	A	982	HIS
1	A	615	GLN
1	A	909	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	843/1219 (69%)	0.15	19 (2%) 57 15	74, 151, 300, 453	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	534	ILE	5.7
1	A	519	ILE	4.4
1	A	679	THR	3.5
1	A	623	ILE	3.3
1	A	532	HIS	3.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.