



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:51 PM GMT

PDB ID : 3PWF
Title : High resolution structure of the fully reduced form of rubrerythrin from *P. furiosus*
Authors : Dillard, B.D.; Demick, J.M.; Adams, M.W.; Lanzilotta, W.N.
Deposited on : 2010-12-08
Resolution : 1.64 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1326 (1.66-1.62)
Clashscore	79885	1525 (1.66-1.62)
Ramachandran outliers	78287	1490 (1.66-1.62)
Sidechain outliers	78261	1490 (1.66-1.62)
RSRZ outliers	66119	1326 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	170	
1	B	170	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3019 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rubrerythrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	0	1	0
			1371	886	223	253	9			
1	B	170	Total	C	N	O	S	0	2	0
			1380	890	223	258	9			

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Fe	0	0
			3	3		
2	A	3	Total	Fe	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total	O	0	0
			133	133		
3	B	129	Total	O	0	0
			129	129		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rubrerythrin



• Molecule 1: Rubrerythrin



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	104.87Å 104.87Å 79.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.63 – 1.64 33.63 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.9 (33.63-1.64) 99.9 (33.63-1.64)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.13 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.166 , 0.188 0.165 , 0.189	Depositor DCC
R_{free} test set	2805 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 55213 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3019	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.80	24/1402 (1.7%)	1.45	16/1882 (0.9%)
1	B	1.74	25/1411 (1.8%)	1.45	14/1895 (0.7%)
All	All	1.77	49/2813 (1.7%)	1.45	30/3777 (0.8%)

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	9	LYS	CB-CG	-17.16	1.06	1.52
1	B	166	LYS	CB-CG	-7.84	1.31	1.52
1	A	157	CYS	CB-SG	7.62	1.95	1.82
1	B	147	TYR	CG-CD2	-7.36	1.29	1.39
1	B	80	GLU	CD-OE1	-7.27	1.17	1.25
1	A	91	TYR	CD1-CE1	7.20	1.50	1.39
1	A	20	SER	CB-OG	7.07	1.51	1.42
1	B	50	TYR	CE2-CZ	-7.05	1.29	1.38
1	A	80	GLU	CD-OE2	-7.04	1.18	1.25
1	A	121	TYR	CD1-CE1	6.80	1.49	1.39
1	A	57	LYS	CD-CE	-6.56	1.34	1.51
1	B	26	TYR	CG-CD1	6.34	1.47	1.39
1	B	102	GLU	CD-OE2	6.32	1.32	1.25
1	A	37	GLY	N-CA	6.32	1.55	1.46
1	A	155	GLU	CD-OE1	-6.18	1.18	1.25
1	A	167	PHE	CE1-CZ	6.16	1.49	1.37
1	B	5	ARG	CZ-NH2	6.11	1.41	1.33
1	A	16	PHE	CD2-CE2	6.06	1.51	1.39
1	B	162	ALA	CA-CB	5.98	1.65	1.52
1	B	114	GLU	CD-OE1	5.96	1.32	1.25
1	A	57	LYS	CE-NZ	5.95	1.64	1.49
1	A	132	GLU	CB-CG	5.89	1.63	1.52
1	A	114	GLU	CD-OE1	5.89	1.32	1.25
1	B	147	TYR	CE2-CZ	-5.88	1.30	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	132	GLU	CB-CG	5.87	1.63	1.52
1	B	50	TYR	CD1-CE1	5.86	1.48	1.39
1	B	121	TYR	CE2-CZ	5.82	1.46	1.38
1	B	26	TYR	CG-CD2	5.75	1.46	1.39
1	B	147	TYR	CD2-CE2	5.65	1.47	1.39
1	B	115	LYS	C-O	5.61	1.33	1.23
1	A	85	GLU	CD-OE2	-5.58	1.19	1.25
1	A	102	GLU	CD-OE1	5.54	1.31	1.25
1	A	52	GLU	CB-CG	5.50	1.62	1.52
1	B	157	CYS	CB-SG	5.49	1.91	1.82
1	B	119[A]	GLU	CG-CD	5.48	1.60	1.51
1	B	119[B]	GLU	CG-CD	5.48	1.60	1.51
1	B	4	LYS	CD-CE	5.44	1.64	1.51
1	A	102	GLU	CD-OE2	5.40	1.31	1.25
1	B	71	GLU	CB-CG	5.38	1.62	1.52
1	A	31	GLU	CB-CG	-5.33	1.42	1.52
1	B	168	VAL	CB-CG1	5.33	1.64	1.52
1	B	110	ALA	CA-CB	5.25	1.63	1.52
1	A	88	TYR	CD2-CE2	5.22	1.47	1.39
1	A	83	GLU	CB-CG	5.19	1.62	1.52
1	A	88	TYR	CE2-CZ	5.19	1.45	1.38
1	A	52	GLU	CG-CD	5.14	1.59	1.51
1	B	121	TYR	CD1-CE1	5.08	1.47	1.39
1	A	122	ARG	CG-CD	5.03	1.64	1.51
1	B	72	ASN	N-CA	5.03	1.56	1.46

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	93	LYS	CD-CE-NZ	-9.96	88.79	111.70
1	A	46	ARG	NE-CZ-NH2	-8.96	115.82	120.30
1	A	122	ARG	CG-CD-NE	-8.40	94.15	111.80
1	A	91	TYR	CB-CG-CD2	-7.71	116.38	121.00
1	B	82	PHE	CB-CG-CD1	-7.47	115.57	120.80
1	A	122	ARG	NE-CZ-NH2	-7.28	116.66	120.30
1	B	4	LYS	CD-CE-NZ	7.15	128.14	111.70
1	A	105	ARG	NE-CZ-NH2	-6.91	116.85	120.30
1	B	29	PHE	CB-CG-CD1	-6.83	116.02	120.80
1	A	105	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	B	5	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	B	25	ARG	NE-CZ-NH1	-6.72	116.94	120.30
1	B	156	TYR	CZ-CE2-CD2	6.69	125.82	119.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	TYR	CB-CG-CD1	-6.64	117.01	121.00
1	A	155	GLU	CG-CD-OE1	-6.64	105.03	118.30
1	B	19	GLU	OE1-CD-OE2	-6.60	115.38	123.30
1	A	102	GLU	OE1-CD-OE2	6.42	131.00	123.30
1	B	157	CYS	N-CA-CB	-6.40	99.09	110.60
1	A	12	LEU	CB-CG-CD2	6.31	121.72	111.00
1	B	102	GLU	OE1-CD-OE2	6.13	130.65	123.30
1	A	25	ARG	NE-CZ-NH1	-6.04	117.28	120.30
1	A	155	GLU	OE1-CD-OE2	5.96	130.45	123.30
1	B	151	ASP	CB-CG-OD2	5.93	123.64	118.30
1	A	9	LYS	CB-CA-C	5.84	122.08	110.40
1	B	140	TYR	CB-CG-CD1	-5.69	117.59	121.00
1	A	147	TYR	CB-CG-CD1	-5.50	117.70	121.00
1	A	80	GLU	OE1-CD-OE2	-5.47	116.74	123.30
1	B	46	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	A	60	PHE	CB-CG-CD1	-5.25	117.13	120.80
1	A	9	LYS	CB-CG-CD	-5.21	98.06	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1371	0	1357	20	1
1	B	1380	0	1355	29	1
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	133	0	0	8	0
3	B	129	0	0	18	0
All	All	3019	0	2712	49	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (49) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:9:LYS:HD2	3:B:213:HOH:O	1.13	1.29
1:A:13:GLU:HG2	3:A:175:HOH:O	1.40	1.20
1:B:75:MET:HB2	3:B:266:HOH:O	1.38	1.20
1:B:13:GLU:HG2	3:B:210:HOH:O	1.40	1.18
1:B:4:LYS:HE3	3:B:302:HOH:O	1.55	1.06
1:A:119:GLU:CD	1:A:122:ARG:HH22	1.66	0.97
1:B:5:ARG:CZ	3:B:270:HOH:O	2.13	0.96
1:A:127:LYS:HD2	1:A:134:ILE:HG22	1.43	0.96
1:B:75:MET:HE2	3:B:266:HOH:O	1.67	0.93
1:A:127:LYS:HG2	3:A:268:HOH:O	1.66	0.93
1:A:119:GLU:OE2	1:A:122:ARG:NH2	2.07	0.87
1:B:4:LYS:CE	3:B:302:HOH:O	2.24	0.73
1:A:122:ARG:NH1	3:A:239:HOH:O	2.26	0.69
1:A:138:LYS:HE3	1:A:140:TYR:OH	1.92	0.68
1:A:108:HIS:HD2	3:A:212:HOH:O	1.76	0.68
1:B:43:LYS:NZ	1:B:43:LYS:CB	2.58	0.66
1:A:127:LYS:CG	3:A:268:HOH:O	2.34	0.66
1:A:119:GLU:CD	1:A:122:ARG:NH2	2.45	0.66
1:B:116:ILE:HD11	1:B:147:TYR:CE1	2.33	0.64
1:B:43:LYS:NZ	1:B:43:LYS:HB2	2.14	0.63
1:A:119:GLU:OE1	1:A:122:ARG:NH2	2.30	0.62
1:A:108:HIS:HE1	3:A:253:HOH:O	1.82	0.61
1:B:136:ILE:O	1:B:136:ILE:HG13	2.03	0.58
1:A:75:MET:HE3	3:A:208:HOH:O	2.04	0.58
1:B:123:LYS:HG3	3:B:233:HOH:O	2.03	0.58
1:B:43:LYS:HZ3	1:B:43:LYS:CB	2.18	0.56
1:B:119[B]:GLU:OE2	3:B:233:HOH:O	2.18	0.56
1:B:108:HIS:HD2	3:B:269:HOH:O	1.87	0.56
1:B:83:GLU:OE2	3:B:191:HOH:O	2.17	0.55
1:A:9:LYS:HG3	1:A:10[A]:LYS:N	2.19	0.53
1:B:9:LYS:CD	3:B:213:HOH:O	1.98	0.52
1:B:75:MET:CB	3:B:266:HOH:O	2.20	0.52
1:A:9:LYS:HG3	1:A:10[B]:LYS:N	2.21	0.52
1:B:9:LYS:HB2	3:B:213:HOH:O	2.10	0.51
1:B:123:LYS:CG	3:B:233:HOH:O	2.60	0.51
1:B:116:ILE:CD1	1:B:147:TYR:CE1	2.94	0.50
1:B:108:HIS:CD2	3:B:269:HOH:O	2.65	0.49
1:A:122:ARG:HD3	3:A:247:HOH:O	2.13	0.48
1:B:163:PRO:O	1:B:166:LYS:HG2	2.14	0.48
1:B:43:LYS:HZ3	1:B:43:LYS:HB3	1.78	0.47
1:A:153:ALA:HB3	1:A:164:LYS:HE3	1.96	0.47
1:B:65:LYS:HE3	3:B:247:HOH:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:127:LYS:HD3	1:A:134:ILE:HA	1.99	0.45
1:A:127:LYS:CD	1:A:134:ILE:HA	2.47	0.44
1:B:43:LYS:HZ2	1:B:43:LYS:HB2	1.81	0.43
1:B:130:LYS:HE2	3:B:195:HOH:O	2.18	0.43
1:A:138:LYS:CE	1:A:140:TYR:OH	2.63	0.43
1:B:140:TYR:CE2	1:B:169:VAL:HG22	2.55	0.42
1:B:125:LYS:O	1:B:129:GLU:HG2	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:10[A]:LYS:NZ	1:B:14[A]:GLU:OE2[2.545]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/170 (99%)	167 (99%)	2 (1%)	0	100	100
1	B	170/170 (100%)	167 (98%)	3 (2%)	0	100	100
All	All	339/340 (100%)	334 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/138 (101%)	136 (98%)	3 (2%)	64	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	140/138 (101%)	139 (99%)	1 (1%)	91	80
All	All	279/276 (101%)	275 (99%)	4 (1%)	78	57

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	129	GLU
1	A	152	GLU
1	B	166	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	HIS
1	B	108	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	170/170 (100%)	-0.29	1 (0%) 86 90	10, 15, 30, 40	0
1	B	170/170 (100%)	-0.28	2 (1%) 75 79	10, 15, 31, 43	1 (0%)
All	All	340/340 (100%)	-0.28	3 (0%) 81 85	10, 15, 31, 43	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	136	ILE	4.5
1	B	137	LYS	3.1
1	A	135	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FE2	A	200	1/1	0.07	0.09	12,12,12,12	0
2	FE2	B	202	1/1	0.07	-0.44	10,10,10,10	0
2	FE2	A	201	1/1	0.06	-0.96	11,11,11,11	0
2	FE2	A	202	1/1	0.06	-1.21	10,10,10,10	0
2	FE2	B	200	1/1	0.05	-1.32	12,12,12,12	0
2	FE2	B	201	1/1	0.04	-2.21	12,12,12,12	0

6.5 Other polymers ⓘ

There are no such residues in this entry.