



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 06:28 PM GMT

PDB ID : 1PYI  
Title : CRYSTAL STRUCTURE OF A PPR1-DNA COMPLEX: DNA RECOGNITION BY PROTEINS CONTAINING A ZN2CYS6 BINUCLEAR CLUSTER  
Authors : Marmorstein, R.; Harrison, S.C.  
Deposited on : 1995-01-04  
Resolution : 3.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1824 (3.30-3.10)
Clashscore	79885	1078 (3.26-3.14)
Ramachandran outliers	78287	1059 (3.26-3.14)
Sidechain outliers	78261	1058 (3.26-3.14)
RSRZ outliers	66119	1825 (3.30-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	D	14	
1	E	14	
2	A	96	
2	B	96	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1826 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*TP\*CP\*GP\*GP\*CP\*AP\*AP\*TP\*TP\*GP\*CP\*CP\*GP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	14	Total	C	N	O	P	0	0	0
			284	136	53	82	13			
1	E	14	Total	C	N	O	P	0	0	0
			284	136	53	82	13			

- Molecule 2 is a protein called PROTEIN (PYRIMIDINE PATHWAY REGULATOR 1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	88	Total	C	N	O	S	0	0	0
			697	438	124	127	8			
2	B	70	Total	C	N	O	S	0	0	0
			556	350	100	98	8			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total	O	0	0
			1	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(\*TP\*CP\*GP\*GP\*CP\*AP\*AP\*TP\*TP\*GP\*CP\*CP\*GP\*A)-3')

Chain D: 




- Molecule 1: DNA (5'-D(\*TP\*CP\*GP\*GP\*CP\*AP\*AP\*TP\*TP\*GP\*CP\*CP\*GP\*A)-3')

Chain E: 





- Molecule 2: PROTEIN (PYRIMIDINE PATHWAY REGULATOR 1)

Chain A: 



- Molecule 2: PROTEIN (PYRIMIDINE PATHWAY REGULATOR 1)

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.80Å 92.60Å 174.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 3.20 10.04 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-3.20) 98.3 (10.04-3.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.91 (at 2.99Å)	Xtriage
Refinement program	?	Depositor
R, $R_{free}$	0.245 , 0.330 0.306 , 0.368	Depositor DCC
$R_{free}$ test set	830 reflections (10.63%)	DCC
Wilson B-factor (Å <sup>2</sup> )	63.2	Xtriage
Anisotropy	0.480	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 51.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	1 of 8639 reflections (0.012%)	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	1826	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	D	1.98	10/318 (3.1%)	3.47	41/489 (8.4%)
1	E	1.58	3/318 (0.9%)	2.65	27/489 (5.5%)
2	A	1.11	5/709 (0.7%)	2.12	33/952 (3.5%)
2	B	1.25	5/565 (0.9%)	2.17	27/757 (3.6%)
All	All	1.41	23/1910 (1.2%)	2.53	128/2687 (4.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
2	A	0	2
All	All	0	4

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	73	PRO	N-CD	9.44	1.61	1.47
1	D	9	DT	C1'-N1	9.20	1.61	1.49
1	D	9	DT	C4-C5	-8.62	1.37	1.45
1	E	23	DT	C5-C7	8.61	1.55	1.50
1	D	9	DT	C2-N3	7.68	1.43	1.37

The worst 5 of 128 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	9	DT	C4-C5-C7	-28.72	101.77	119.00
1	D	9	DT	C6-C5-C7	27.81	139.59	122.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	9	DT	C5-C4-O4	-22.06	109.45	124.90
1	D	9	DT	N3-C4-O4	15.63	129.28	119.90
1	D	14	DA	O4'-C1'-N9	13.77	117.64	108.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	104	ASN	Sidechain
2	A	111	ASP	Sidechain
1	D	3	DG	Sidechain
1	D	9	DT	Sidechain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	284	0	159	25	0
1	E	284	0	159	32	0
2	A	697	0	714	112	0
2	B	556	0	575	96	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	E	1	0	0	0	0
All	All	1826	0	1607	223	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 65.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:113:PRO:HB2	2:A:116:LEU:HD11	1.29	1.14
2:A:97:ASP:HB2	2:A:98:PRO:HD2	1.31	1.05
1:D:1:DT:H6	1:D:1:DT:HO5'	1.05	0.97
2:A:101:ILE:HG21	2:B:83:ARG:HE	1.32	0.95

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:E:21:DA:H2"	1:E:22:DT:H5"	1.48	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	86/96 (90%)	59 (69%)	16 (19%)	11 (13%)	0	3
2	B	68/96 (71%)	54 (79%)	10 (15%)	4 (6%)	2	20
All	All	154/192 (80%)	113 (73%)	26 (17%)	15 (10%)	1	7

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	46	GLN
2	A	97	ASP
2	A	113	PRO
2	B	97	ASP
2	A	67	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	80/88 (91%)	61 (76%)	19 (24%)	1	4
2	B	64/88 (73%)	49 (77%)	15 (23%)	1	5
All	All	144/176 (82%)	110 (76%)	34 (24%)	1	4



5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	105	ILE
2	B	32	THR
2	B	83	ARG
2	A	112	GLU
2	A	64	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	46	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	D	14/14 (100%)	0.13	0 100 100	8, 25, 36, 49	0
1	E	14/14 (100%)	0.26	0 100 100	14, 28, 46, 51	0
2	A	88/96 (91%)	0.02	0 100 100	11, 32, 54, 57	0
2	B	70/96 (72%)	-0.14	0 100 100	3, 23, 43, 62	0
All	All	186/220 (84%)	-0.02	0 100 100	3, 29, 51, 62	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	B	321	1/1	0.13	-1.72	12,12,12,12	0
3	ZN	A	22	1/1	0.06	-1.93	26,26,26,26	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	21	1/1	0.11	-2.57	26,26,26,26	0
3	ZN	B	322	1/1	0.08	-3.04	16,16,16,16	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.