



Full wwPDB X-ray Structure Validation Report i

Apr 8, 2014 – 10:32 PM EDT

PDB ID : 4PYZ
Title : Crystal structure of the first two Ubl domains of Deubiquitylase USP7
Authors : Walker, J.R.; Dong, A.; Ong, M.S.; Dhe-Paganon, S.; Kania, J.; Bountra, C.;
Arrowsmith, C.H.; Edwards, A.M.; Tong, Y.; Structural Genomics Consortium
(SGC)
Deposited on : 2014-03-28
Resolution : 2.84 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1439
EDS	:	FAILED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22978

1 Overall quality at a glance

The reported resolution of this entry is 2.84 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2848 (2.88-2.80)
Ramachandran outliers	78287	2786 (2.88-2.80)
Sidechain outliers	78261	2789 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	273	
1	B	273	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4106 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	2	0
			2077	1308	352	404	13			
1	B	261	Total	C	N	O	S	0	1	0
			2015	1271	335	395	14			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	521	SER	-	EXPRESSION TAG	UNP Q93009
A	522	HIS	-	EXPRESSION TAG	UNP Q93009
A	523	HIS	-	EXPRESSION TAG	UNP Q93009
A	524	HIS	-	EXPRESSION TAG	UNP Q93009
A	525	HIS	-	EXPRESSION TAG	UNP Q93009
A	526	HIS	-	EXPRESSION TAG	UNP Q93009
A	527	HIS	-	EXPRESSION TAG	UNP Q93009
A	528	SER	-	EXPRESSION TAG	UNP Q93009
A	529	SER	-	EXPRESSION TAG	UNP Q93009
A	530	GLY	-	EXPRESSION TAG	UNP Q93009
A	531	LEU	-	EXPRESSION TAG	UNP Q93009
A	532	VAL	-	EXPRESSION TAG	UNP Q93009
A	533	PRO	-	EXPRESSION TAG	UNP Q93009
A	534	ARG	-	EXPRESSION TAG	UNP Q93009
A	535	GLY	-	EXPRESSION TAG	UNP Q93009
A	536	SER	-	EXPRESSION TAG	UNP Q93009
B	521	SER	-	EXPRESSION TAG	UNP Q93009
B	522	HIS	-	EXPRESSION TAG	UNP Q93009
B	523	HIS	-	EXPRESSION TAG	UNP Q93009
B	524	HIS	-	EXPRESSION TAG	UNP Q93009
B	525	HIS	-	EXPRESSION TAG	UNP Q93009
B	526	HIS	-	EXPRESSION TAG	UNP Q93009
B	527	HIS	-	EXPRESSION TAG	UNP Q93009
B	528	SER	-	EXPRESSION TAG	UNP Q93009
B	529	SER	-	EXPRESSION TAG	UNP Q93009

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	530	GLY	-	EXPRESSION TAG	UNP Q93009
B	531	LEU	-	EXPRESSION TAG	UNP Q93009
B	532	VAL	-	EXPRESSION TAG	UNP Q93009
B	533	PRO	-	EXPRESSION TAG	UNP Q93009
B	534	ARG	-	EXPRESSION TAG	UNP Q93009
B	535	GLY	-	EXPRESSION TAG	UNP Q93009
B	536	SER	-	EXPRESSION TAG	UNP Q93009

- Molecule 2 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total X 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	6	Total O 6 6	0	0
3	B	6	Total O 6 6	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7

Chain A: 



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7

Chain B: 



4 Data and refinement statistics

EDS failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.33Å 103.75Å 116.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.71 – 2.84	Depositor
% Data completeness (in resolution range)	98.3 (40.71-2.84)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.10.0, COOT 0.7.2	Depositor
R, R_{free}	0.244 , 0.287	Depositor
Wilson B-factor (Å ²)	66.4	Xtriage
Anisotropy	1.097	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	2 of 19770 reflections (0.010%)	Xtriage
Total number of atoms	4106	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2119	0.55	0/2876
1	B	0.38	0/2056	0.55	0/2795
All	All	0.39	0/4175	0.55	0/5671

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2077	0	0	0	0
1	B	2015	0	0	0	0
2	B	2	0	0	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
All	All	4106	0	0	0	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/273 (96%)	259 (98%)	4 (2%)	0	100	100
1	B	260/273 (95%)	253 (97%)	7 (3%)	0	100	100
All	All	523/546 (96%)	512 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/249 (87%)	210 (97%)	6 (3%)	56	88
1	B	203/249 (82%)	200 (98%)	3 (2%)	76	96
All	All	419/498 (84%)	410 (98%)	9 (2%)	64	93

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	619	GLN
1	A	669	LEU
1	A	676	LEU
1	A	705	ILE
1	A	747	GLN
1	A	772	ASP
1	B	534	ARG
1	B	607	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	637	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS failed to run properly - this section will therefore be empty.

6.4 Ligands ⓘ

EDS failed to run properly - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS failed to run properly - this section will therefore be empty.