



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 06:19 AM GMT

PDB ID : 3QBD
Title : 3-Dehydroquinate Synthase (aroB) from Mycobacterium tuberculosis in complex with NAD
Authors : Cheng, W.C.; Chen, T.J.; Wang, W.C.
Deposited on : 2011-01-13
Resolution : 2.47 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3277 (2.50-2.46)
Clashscore	79885	4136 (2.50-2.46)
Ramachandran outliers	78287	4052 (2.50-2.46)
Sidechain outliers	78261	4054 (2.50-2.46)
RSRZ outliers	66119	3279 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	368	
1	B	368	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5260 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-dehydroquinate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	0	0
			2546	1621	451	467	7			
1	B	342	Total	C	N	O	S	0	0	0
			2537	1616	449	465	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP A5U5N7
A	-4	HIS	-	EXPRESSION TAG	UNP A5U5N7
A	-3	HIS	-	EXPRESSION TAG	UNP A5U5N7
A	-2	HIS	-	EXPRESSION TAG	UNP A5U5N7
A	-1	HIS	-	EXPRESSION TAG	UNP A5U5N7
A	0	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	-5	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	-4	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	-3	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	-2	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	-1	HIS	-	EXPRESSION TAG	UNP A5U5N7
B	0	HIS	-	EXPRESSION TAG	UNP A5U5N7

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total	O	0	0
			53	53		
3	B	36	Total	O	0	0
			36	36		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-dehydroquinate synthase



- Molecule 1: 3-dehydroquinate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	142.83Å 142.83Å 37.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.47 28.70 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.47) 99.8 (28.70-2.47)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.196 , 0.258 0.196 , 0.252	Depositor DCC
R_{free} test set	1389 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 15.2	EDS
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 27611 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5260	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 72.50 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.1765e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/2589 (0.1%)	0.92	7/3528 (0.2%)
1	B	1.08	1/2580 (0.0%)	0.89	2/3516 (0.1%)
All	All	1.09	4/5169 (0.1%)	0.91	9/7044 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	307	TYR	CD2-CE2	-5.48	1.31	1.39
1	A	307	TYR	CD1-CE1	-5.36	1.31	1.39
1	A	14	VAL	CB-CG2	-5.31	1.41	1.52
1	B	307	TYR	CD2-CE2	-5.20	1.31	1.39

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	297	ILE	CG1-CB-CG2	-5.90	98.42	111.40
1	B	340	LYS	CD-CE-NZ	-5.76	98.46	111.70
1	A	308	ASP	CB-CG-OD1	5.69	123.42	118.30
1	A	308	ASP	CB-CG-OD2	-5.68	113.19	118.30
1	A	122	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	A	295	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	143	GLY	N-CA-C	5.34	126.44	113.10
1	A	97	ARG	NE-CZ-NH2	-5.11	117.75	120.30
1	A	32	ASP	CB-CG-OD2	-5.04	113.77	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2546	0	2623	43	0
1	B	2537	0	2615	39	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
3	A	53	0	0	2	0
3	B	36	0	0	0	0
All	All	5260	0	5290	79	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (79) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:58:ARG:HH11	1:A:58:ARG:HG2	1.29	0.97
1:A:97:ARG:HH12	1:B:144:LYS:HE2	1.32	0.93
1:A:97:ARG:NH2	1:A:122:ARG:HH11	1.65	0.93
1:A:97:ARG:HH22	1:A:122:ARG:HH11	1.09	0.90
1:A:58:ARG:CG	1:A:58:ARG:HH11	1.85	0.88
1:A:27:LEU:HD13	1:A:58:ARG:NE	1.92	0.85
1:A:97:ARG:HH22	1:A:122:ARG:NH1	1.77	0.83
1:B:48:LEU:HD23	1:B:48:LEU:N	1.94	0.82
1:B:338:LEU:O	1:B:340:LYS:HD2	1.80	0.80
1:A:97:ARG:NH2	1:A:122:ARG:NH1	2.32	0.77
1:A:97:ARG:NH1	1:B:144:LYS:HE2	2.03	0.74
1:B:212:PRO:HA	1:B:217:LEU:HD23	1.72	0.71
1:A:264:ARG:HD2	3:A:415:HOH:O	1.92	0.69
1:A:58:ARG:NH1	1:A:58:ARG:HG2	2.09	0.67
1:B:9:THR:HG22	1:B:21:VAL:HG13	1.78	0.66
1:A:103:SER:O	1:A:129:LEU:HB2	1.95	0.66
1:B:132:THR:O	1:B:136:MET:HG3	1.96	0.65
1:B:28:ASP:OD1	1:B:58:ARG:NH2	2.27	0.65
1:A:58:ARG:CG	1:A:58:ARG:NH1	2.57	0.62
1:B:257:ARG:HD2	1:B:261:TYR:OH	1.99	0.62
1:B:47:GLY:C	1:B:48:LEU:HD23	2.20	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:27:LEU:CD1	1:A:58:ARG:NE	2.62	0.61
1:A:240:GLU:HA	1:A:240:GLU:OE1	1.99	0.61
1:B:48:LEU:CD2	1:B:48:LEU:N	2.65	0.59
1:A:121:LEU:O	1:A:122:ARG:HB2	2.03	0.58
1:B:44:HIS:HD2	1:B:45:GLN:O	1.84	0.58
1:A:129:LEU:HD12	1:A:166:LEU:HB2	1.85	0.57
1:A:259:GLU:O	1:A:262:ARG:HG2	2.05	0.56
1:A:80:LEU:HB3	1:A:81:PRO:HD3	1.88	0.56
1:B:48:LEU:HD12	1:B:104:LEU:HG	1.89	0.54
1:A:53:GLU:O	1:A:57:LYS:HB2	2.08	0.53
1:B:331:ARG:NH2	1:B:343:ARG:HD3	2.24	0.53
1:B:86:ILE:O	1:B:90:LEU:HG	2.08	0.53
1:B:262:ARG:HH11	1:B:262:ARG:HG3	1.73	0.53
1:A:27:LEU:HD13	1:A:58:ARG:HE	1.71	0.52
1:B:280:ALA:HB2	1:B:334:VAL:HB	1.92	0.52
1:A:48:LEU:HD13	1:A:104:LEU:HG	1.93	0.51
1:A:58:ARG:NH1	1:A:58:ARG:HB3	2.25	0.51
1:A:44:HIS:HD2	1:A:45:GLN:O	1.94	0.50
1:A:151:ALA:O	1:A:155:LEU:HD12	2.11	0.50
1:A:58:ARG:CB	1:A:58:ARG:HH11	2.24	0.49
1:B:177:ARG:HG2	1:B:181:ILE:HD12	1.94	0.49
1:A:122:ARG:NH2	1:B:154:ASN:O	2.25	0.48
1:B:85:PHE:O	1:B:89:VAL:HG23	2.13	0.48
1:B:53:GLU:O	1:B:57:LYS:HG2	2.13	0.48
1:B:331:ARG:HH22	1:B:343:ARG:NH1	2.10	0.47
1:B:78:LYS:NZ	1:B:154:ASN:HD21	2.12	0.47
1:A:102:VAL:CG1	1:A:129:LEU:HD22	2.44	0.47
1:B:259:GLU:HG3	1:B:263:TRP:CD1	2.49	0.47
1:A:31:GLU:OE1	1:A:58:ARG:NH1	2.44	0.47
1:B:135:GLY:HA2	1:B:139:ALA:HB3	1.96	0.47
1:A:267:ALA:HB1	1:A:304:PRO:HG3	1.97	0.46
1:A:103:SER:OG	1:A:128:HIS:HA	2.16	0.46
1:B:46:PRO:HD3	1:B:71:ILE:O	2.17	0.44
1:A:203:GLU:OE1	1:A:293:ARG:NH2	2.45	0.44
1:A:259:GLU:O	1:A:260:ARG:C	2.54	0.44
1:B:292:GLN:HG3	1:B:295:ARG:HH21	1.83	0.44
1:A:135:GLY:HA2	1:A:139:ALA:HB3	2.00	0.43
1:A:58:ARG:CB	1:A:58:ARG:NH1	2.82	0.43
1:B:6:ALA:N	1:B:7:PRO:CD	2.81	0.43
1:B:27:LEU:HD13	1:B:58:ARG:HD3	2.01	0.42
1:B:130:PRO:HG3	1:B:140:ALA:CB	2.49	0.42
1:A:27:LEU:CD1	1:A:58:ARG:HE	2.30	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:267:ALA:HB1	1:A:304:PRO:CG	2.50	0.42
1:A:180:MET:CE	1:A:212:PRO:HG3	2.49	0.42
1:A:308:ASP:HA	1:A:309:PRO:HD3	1.91	0.42
1:A:262:ARG:HE	1:A:262:ARG:HB3	1.45	0.42
1:B:45:GLN:HG3	1:B:109:ALA:HB2	2.03	0.41
1:B:80:LEU:N	1:B:81:PRO:HD2	2.35	0.41
1:B:217:LEU:HB3	1:B:218:PRO:HD3	2.01	0.41
1:A:59:LEU:O	1:A:64:VAL:HB	2.21	0.41
1:B:98:LYS:HE2	1:B:98:LYS:HB3	1.94	0.41
1:B:103:SER:OG	1:B:128:HIS:HA	2.21	0.41
1:B:228:LYS:HE3	1:B:245:LEU:HD13	2.02	0.41
1:A:182:CYS:HB2	3:A:415:HOH:O	2.21	0.41
1:A:79:ASP:O	1:A:80:LEU:C	2.59	0.40
1:B:331:ARG:HH22	1:B:343:ARG:HH11	1.68	0.40
1:B:228:LYS:O	1:B:232:VAL:HG13	2.21	0.40
1:B:168:ASP:O	1:B:171:THR:OG1	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/368 (92%)	326 (96%)	11 (3%)	1 (0%)	50	71
1	B	336/368 (91%)	317 (94%)	18 (5%)	1 (0%)	50	71
All	All	674/736 (92%)	643 (95%)	29 (4%)	2 (0%)	50	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	25	GLY
1	A	25	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/280 (92%)	248 (96%)	11 (4%)	40	65
1	B	259/280 (92%)	254 (98%)	5 (2%)	69	90
All	All	518/560 (92%)	502 (97%)	16 (3%)	52	78

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	58	ARG
1	A	97	ARG
1	A	122	ARG
1	A	129	LEU
1	A	166	LEU
1	A	240	GLU
1	A	257	ARG
1	A	262	ARG
1	A	305	VAL
1	A	340	LYS
1	B	27	LEU
1	B	48	LEU
1	B	262	ARG
1	B	296	THR
1	B	340	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	154	ASN
1	B	44	HIS
1	B	154	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	A	400	-	48,48,48	0.92	2 (4%)	73,73,73	1.96	16 (21%)
2	NAD	B	400	-	48,48,48	0.97	1 (2%)	73,73,73	1.78	13 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	400	-	-	0/30/62/62	0/3/5/5
2	NAD	B	400	-	-	0/30/62/62	0/3/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	400	NAD	C4A-N9A	-3.67	1.32	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAD	O4B-C1B	2.52	1.45	1.41
2	A	400	NAD	C4A-N9A	-2.16	1.34	1.37

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	NAD	N3A-C2A-N1A	-7.42	122.51	128.71
2	B	400	NAD	N3A-C2A-N1A	-7.38	122.54	128.71
2	A	400	NAD	O4B-C1B-N9A	-6.65	102.25	108.44
2	A	400	NAD	N3A-C4A-N9A	5.69	135.71	125.43
2	B	400	NAD	O4D-C1D-N1N	4.93	113.00	107.95
2	B	400	NAD	C3N-C7N-N7N	4.53	122.93	117.77
2	B	400	NAD	N3A-C4A-N9A	3.91	132.49	125.43
2	A	400	NAD	O4B-C1B-C2B	-3.79	100.96	106.77
2	B	400	NAD	O7N-C7N-N7N	-3.77	117.14	122.59
2	A	400	NAD	C5A-C4A-N3A	-3.45	118.20	125.70
2	A	400	NAD	C8A-N9A-C4A	3.37	109.47	106.90
2	A	400	NAD	O7N-C7N-C3N	-3.08	116.10	119.58
2	A	400	NAD	C2A-N3A-C4A	3.03	122.64	114.01
2	A	400	NAD	O3B-C3B-C4B	-2.95	102.39	111.08
2	B	400	NAD	O2A-PA-O1A	2.81	127.90	112.21
2	A	400	NAD	C3N-C7N-N7N	2.78	120.93	117.77
2	B	400	NAD	O4B-C1B-N9A	-2.57	106.05	108.44
2	A	400	NAD	O4D-C1D-C2D	-2.49	102.95	106.77
2	A	400	NAD	C5N-C4N-C3N	-2.47	117.11	120.32
2	B	400	NAD	O3-PN-O1N	2.46	114.71	108.83
2	A	400	NAD	C3D-C2D-C1D	2.39	104.64	100.91
2	B	400	NAD	O4D-C1D-C2D	-2.32	103.21	106.77
2	A	400	NAD	C4D-O4D-C1D	-2.28	107.27	109.75
2	B	400	NAD	O3B-C3B-C4B	-2.22	104.54	111.08
2	B	400	NAD	C5A-C4A-N3A	-2.22	120.87	125.70
2	A	400	NAD	N7A-C8A-N9A	-2.20	108.13	114.36
2	B	400	NAD	C2A-N3A-C4A	2.18	120.22	114.01
2	B	400	NAD	N6A-C6A-N1A	2.18	123.64	119.36
2	A	400	NAD	O2N-PN-O1N	2.08	125.00	118.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/368 (93%)	-0.32	5 (1%) 70 72	13, 27, 45, 61	0
1	B	342/368 (92%)	-0.16	2 (0%) 86 88	17, 34, 55, 64	0
All	All	686/736 (93%)	-0.24	7 (1%) 79 80	13, 30, 51, 64	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	97	ARG	3.2
1	A	260	ARG	2.8
1	A	361	ALA	2.6
1	A	314	GLN	2.5
1	B	53	GLU	2.2
1	A	177	ARG	2.2
1	A	240	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAD	A	400	44/44	0.11	-0.31	18,25,32,33	0
2	NAD	B	400	44/44	0.11	-0.63	22,31,40,45	0

6.5 Other polymers ⓘ

There are no such residues in this entry.