



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 06:43 AM GMT

PDB ID : 2QL5  
Title : Crystal Structure of caspase-7 with inhibitor AC-DMQD-CHO  
Authors : Agniswamy, J.; Fang, B.; Weber, I.  
Deposited on : 2007-07-12  
Resolution : 2.34 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	4049 (2.38-2.30)
Clashscore	79885	1094 (2.36-2.32)
Ramachandran outliers	78287	1080 (2.36-2.32)
Sidechain outliers	78261	1081 (2.36-2.32)
RSRZ outliers	66119	4050 (2.38-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	173	
1	C	173	
2	B	97	
2	D	97	
3	E	5	
3	F	5	
4	G	5	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	CIT	D	850	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3963 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	0	0
			1100	691	188	210	11			
1	C	140	Total	C	N	O	S	0	0	0
			1100	691	188	210	11			

- Molecule 2 is a protein called Caspase-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	92	Total	C	N	O	S	0	0	0
			758	487	128	139	4			
2	D	92	Total	C	N	O	S	0	0	0
			758	487	128	139	4			

- Molecule 3 is a protein called inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	5	Total	C	N	O	S	0	0	0
			36	20	5	10	1			
3	F	5	Total	C	N	O	S	0	0	0
			36	20	5	10	1			

- Molecule 4 is a protein called peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	5	Total	C	N	O	0	0	0
			37	20	8	9			

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			13	6	7		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	33	Total	O	0	0
			33	33		
6	B	23	Total	O	0	0
			23	23		
6	C	43	Total	O	0	0
			43	43		
6	D	23	Total	O	0	0
			23	23		
6	E	1	Total	O	0	0
			1	1		
6	F	2	Total	O	0	0
			2	2		




- Molecule 3: inhibitor

Chain F: 



- Molecule 4: peptide

Chain G: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.25Å 87.25Å 187.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.34 39.87 – 2.34	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.34) 85.0 (39.87-2.34)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.78 (at 2.34Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.212 , 0.233 0.216 , 0.239	Depositor DCC
$R_{free}$ test set	1497 reflections (4.94%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.7	Xtriage
Anisotropy	0.963	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.7	EDS
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32067 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3963	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ASJ, ACE, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/1117	0.62	1/1496 (0.1%)
1	C	0.39	0/1117	0.65	2/1496 (0.1%)
2	B	0.36	0/780	0.61	0/1054
2	D	0.38	0/780	0.62	0/1054
3	E	0.58	0/25	0.81	0/32
3	F	0.97	0/25	1.61	1/32 (3.1%)
4	G	0.46	0/37	0.47	0/46
All	All	0.39	0/3881	0.64	4/5210 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	802	ASP	N-CA-CB	-6.76	98.43	110.60
1	C	422	LEU	CA-CB-CG	6.18	129.51	115.30
1	A	152	GLY	N-CA-C	-5.77	98.68	113.10
1	C	452	GLY	N-CA-C	-5.58	99.15	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1100	0	1085	26	0
1	C	1100	0	1085	21	0
2	B	758	0	731	13	0
2	D	758	0	731	16	0
3	E	36	0	29	4	0
3	F	36	0	29	0	0
4	G	37	0	26	1	0
5	D	13	0	6	2	0
6	A	33	0	0	1	0
6	B	23	0	0	0	0
6	C	43	0	0	0	0
6	D	23	0	0	0	0
6	E	1	0	0	0	0
6	F	2	0	0	0	0
All	All	3963	0	3722	73	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (73) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:80:LYS:HD3	1:A:80:LYS:H	1.06	1.09
1:A:57:THR:HG22	1:A:58:TYR:H	1.33	0.93
1:A:57:THR:HG22	1:A:58:TYR:N	1.90	0.86
1:A:80:LYS:HD3	1:A:80:LYS:N	1.91	0.82
1:A:80:LYS:CD	1:A:80:LYS:H	1.87	0.81
1:C:495:ILE:O	1:C:496:GLN:HB2	1.81	0.80
1:C:464:ALA:O	1:C:467:ARG:HG3	1.83	0.79
2:D:527:PRO:HG3	5:D:850:CIT:O4	1.85	0.76
1:A:186:CYS:H	3:E:705:ASJ:H	1.53	0.72
1:C:463:THR:HG21	2:D:521:PHE:HE2	1.53	0.70
2:D:533:ARG:HA	2:D:539:SER:HA	1.71	0.70
1:C:366:LYS:HE2	1:C:433:ASN:ND2	2.07	0.69
2:B:233:ARG:HA	2:B:239:SER:HA	1.75	0.68
2:D:592:VAL:HG11	5:D:850:CIT:H21	1.75	0.67
2:B:266:ASN:OD1	2:B:289:PRO:HB2	1.95	0.67
2:B:212:LYS:HA	1:C:495:ILE:O	1.95	0.65
1:A:97:LEU:HD13	1:A:140:ILE:HG21	1.80	0.63
1:A:124:LYS:O	1:A:128:GLU:HG3	1.99	0.62
1:C:449:VAL:HB	1:C:456:VAL:HG13	1.81	0.61
1:A:175:LEU:HD21	1:C:496:GLN:HG2	1.83	0.60
1:A:147:GLU:HG3	1:A:148:ASN:ND2	2.18	0.59
1:C:420:GLN:NE2	1:C:462:LEU:HD23	2.18	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:186:CYS:N	3:E:705:ASJ:H	2.19	0.58
1:A:163:THR:HG21	2:B:221:PHE:HE1	1.70	0.57
2:D:560:GLN:O	2:D:564:ARG:HG3	2.04	0.57
2:B:261:ILE:O	2:B:265:VAL:HG23	2.05	0.56
1:C:366:LYS:HE2	1:C:433:ASN:HD21	1.73	0.54
1:C:397:LEU:HD13	1:C:440:ILE:HG21	1.90	0.54
1:C:495:ILE:HG13	1:C:496:GLN:N	2.23	0.53
1:A:183:ILE:HD12	1:A:183:ILE:N	2.23	0.53
1:C:374:ASN:HD22	1:C:412:ASN:ND2	2.07	0.53
1:C:374:ASN:HD22	1:C:412:ASN:HD22	1.55	0.53
4:G:810:GLN:OE1	4:G:810:GLN:N	2.42	0.52
2:B:288:ILE:HG23	2:B:288:ILE:O	2.09	0.52
3:E:701:ACE:H2	3:E:703:MET:HE2	1.92	0.52
2:B:257:GLU:OE2	2:B:258:ILE:HG22	2.10	0.52
1:A:136:CYS:HB3	1:A:178:PRO:HG2	1.92	0.51
2:D:557:GLU:OE2	2:D:558:ILE:HG22	2.11	0.50
1:A:84:MET:HB3	1:A:144:HIS:CD2	2.47	0.49
1:A:111:TYR:CG	1:A:122:LEU:HD21	2.46	0.49
2:D:566:ASN:OD1	2:D:589:PRO:HB2	2.11	0.49
1:C:369:LYS:HG3	1:C:370:CYS:N	2.28	0.49
2:B:220:LEU:HD23	2:B:220:LEU:C	2.33	0.48
1:C:415:SER:HB2	1:C:454:ASP:OD2	2.15	0.47
1:A:80:LYS:CD	1:A:80:LYS:N	2.65	0.47
2:D:561:ILE:O	2:D:565:VAL:HG23	2.15	0.47
1:A:70:CYS:HA	1:A:138:ALA:O	2.14	0.46
1:A:191:LEU:HD22	2:B:285:LYS:HG3	1.98	0.46
3:E:702:ASP:O	3:E:703:MET:HE2	2.15	0.46
1:A:136:CYS:CB	1:A:178:PRO:HG2	2.46	0.46
2:D:574:GLU:HG2	2:D:584:GLU:HA	1.98	0.45
1:C:495:ILE:HG13	1:C:496:GLN:H	1.82	0.45
1:C:369:LYS:HD2	1:C:409:ILE:HD12	1.99	0.45
2:B:266:ASN:HA	2:B:289:PRO:HD2	1.98	0.44
1:A:57:THR:CG2	1:A:58:TYR:N	2.62	0.44
2:B:258:ILE:HG23	2:B:259:MET:N	2.33	0.43
2:B:251:GLU:HG2	2:B:252:HIS:CD2	2.52	0.43
1:A:66:LYS:HG2	6:A:206:HOH:O	2.17	0.43
1:A:101:PHE:O	1:A:106:PHE:HB2	2.18	0.43
1:C:397:LEU:HD22	1:C:440:ILE:HD13	2.01	0.42
1:A:84:MET:HE2	1:A:144:HIS:HB3	2.01	0.42
2:D:514:PRO:HB2	2:D:517:ALA:HB2	2.01	0.42
2:D:583:HIS:CD2	2:D:584:GLU:HG3	2.54	0.42
2:B:268:ARG:HH12	2:B:272:HIS:CG	2.38	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:578:ASP:O	2:D:580:PRO:HD3	2.20	0.42
2:D:602:SER:O	2:D:603:GLN:HB3	2.19	0.42
1:A:78:PHE:N	1:A:78:PHE:CD1	2.88	0.41
1:C:443:SER:OG	1:C:444:HIS:N	2.53	0.41
1:A:69:LYS:HE3	1:A:129:GLU:OE2	2.20	0.41
1:C:397:LEU:HD12	1:C:442:LEU:HD11	2.03	0.41
2:D:539:SER:O	2:D:543:GLN:HG3	2.21	0.41
2:D:512:LYS:O	2:D:513:ILE:HD12	2.20	0.41
1:C:489:THR:HG22	2:D:530:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/173 (80%)	131 (95%)	7 (5%)	0	100	100
1	C	138/173 (80%)	134 (97%)	4 (3%)	0	100	100
2	B	90/97 (93%)	90 (100%)	0	0	100	100
2	D	90/97 (93%)	88 (98%)	2 (2%)	0	100	100
3	E	3/5 (60%)	3 (100%)	0	0	100	100
3	F	3/5 (60%)	2 (67%)	1 (33%)	0	100	100
4	G	3/5 (60%)	2 (67%)	1 (33%)	0	100	100
All	All	465/555 (84%)	450 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/152 (79%)	118 (98%)	2 (2%)	73	85
1	C	120/152 (79%)	116 (97%)	4 (3%)	50	64
2	B	84/88 (96%)	80 (95%)	4 (5%)	35	45
2	D	84/88 (96%)	83 (99%)	1 (1%)	82	91
3	E	3/3 (100%)	3 (100%)	0	100	100
3	F	3/3 (100%)	2 (67%)	1 (33%)	0	0
4	G	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	417/489 (85%)	404 (97%)	13 (3%)	52	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	LYS
1	A	108	VAL
2	B	213	ILE
2	B	234	SER
2	B	268	ARG
2	B	278	ASP
1	C	398	PHE
1	C	408	VAL
1	C	422	LEU
1	C	454	ASP
2	D	513	ILE
3	F	804	GLN
4	G	812	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	ASN
2	B	303	GLN
1	C	412	ASN
1	C	448	ASN
3	F	804	GLN
4	G	810	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ASJ	E	705	1,3	7,7,7	0.69	0	8,8,8	1.09	1 (12%)
3	ASJ	F	805	1,3	7,7,7	1.12	0	8,8,8	1.21	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASJ	E	705	1,3	-	0/6/6/6	0/0/0/0
3	ASJ	F	805	1,3	-	0/6/6/6	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	805	ASJ	O-C-CA	2.74	117.95	111.87
3	E	705	ASJ	CB-CA-C	-2.44	107.12	112.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	CIT	D	850	-	12,12,12	1.32	1 (8%)	17,17,17	1.53	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CIT	D	850	-	-	0/16/16/16	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	850	CIT	C2-C3	3.03	1.57	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	850	CIT	O6-C6-C3	4.65	119.66	112.89
5	D	850	CIT	O2-C1-O1	-2.07	118.03	123.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/173 (80%)	0.02	3 (2%) 60 71	49, 63, 79, 91	0
1	C	140/173 (80%)	-0.10	2 (1%) 72 81	44, 56, 72, 100	0
2	B	92/97 (94%)	0.57	8 (8%) 10 17	42, 59, 94, 101	0
2	D	92/97 (94%)	0.56	13 (14%) 3 6	46, 58, 95, 101	0
3	E	5/5 (100%)	0.06	0 100 100	64, 73, 75, 76	0
3	F	5/5 (100%)	0.21	0 100 100	63, 71, 75, 75	0
4	G	5/5 (100%)	2.35	2 (40%) 1 1	110, 111, 119, 119	0
All	All	479/555 (86%)	0.22	28 (5%) 21 32	42, 60, 87, 119	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	813	GLY	4.9
2	B	282	PHE	4.3
2	D	592	VAL	4.1
2	B	290	CYS	3.8
1	C	495	ILE	3.7
1	C	496	GLN	3.7
2	D	591	VAL	3.6
1	A	57	THR	3.6
2	B	291	VAL	3.6
2	B	292	VAL	3.5
2	D	520	LEU	3.5
2	D	521	PHE	3.2
4	G	814	GLU	3.2
2	D	593	SER	3.1
2	B	222	ALA	3.0
2	D	590	CYS	3.0
2	D	603	GLN	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	582	PHE	2.6
2	B	278	ASP	2.6
2	D	519	PHE	2.5
2	D	522	ALA	2.5
2	D	594	MET	2.4
2	D	578	ASP	2.4
2	D	583	HIS	2.4
2	B	293	SER	2.3
1	A	196	GLN	2.1
1	A	195	ILE	2.1
2	B	280	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ASJ	E	705	8/8	0.13	0.52	62,63,65,66	0
3	ASJ	F	805	8/8	0.10	-0.36	60,62,64,64	0

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	CIT	D	850	13/13	0.44	3.04	108,111,113,113	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.