



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 03:31 PM GMT

PDB ID : 2QLR
Title : Crystal structure of human kynurenine aminotransferase II
Authors : Han, Q.; Robinson, R.; Li, J.
Deposited on : 2007-07-13
Resolution : 2.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	425	
1	B	425	
1	C	425	
1	D	425	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	GOL	A	427	-	X
2	GOL	C	427	-	X
2	GOL	C	428	-	X
2	GOL	C	429	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
2	GOL	C	430	-	X
2	GOL	D	427	-	X
2	GOL	D	428	-	X
2	GOL	D	429	-	X

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 14176 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine/alpha-aminoadipateaminotransferase mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	425	Total	C	N	O	P	S	0	0	0
			3347	2147	560	621	1	18			
1	B	425	Total	C	N	O	P	S	0	0	0
			3348	2147	560	622	1	18			
1	C	425	Total	C	N	O	P	S	0	0	0
			3347	2147	560	621	1	18			
1	D	425	Total	C	N	O	P	S	0	0	0
			3348	2147	560	622	1	18			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	166	Total O 166 166	0	0
3	B	199	Total O 199 199	0	0
3	C	166	Total O 166 166	0	0
3	D	183	Total O 183 183	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Kynurenine/alpha-aminoadipateaminotransferase mitochondrial

Chain A:



• Molecule 1: Kynurenine/alpha-aminoadipateaminotransferase mitochondrial


Chain B:






• Molecule 1: Kynurenine/alpha-aminoadipateaminotransferase mitochondrial

Chain C:



• Molecule 1: Kynurenine/alpha-aminoadipateaminotransferase mitochondrial

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.50Å 70.97Å 121.13Å 90.00° 101.10° 90.00°	Depositor
Resolution (Å)	30.11 – 2.30 30.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.3 (30.11-2.30) 96.3 (30.10-2.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.242 , 0.256 0.227 , 0.241	Depositor DCC
R_{free} test set	3883 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.56$, $\langle L^2 \rangle = 0.40$	Xtriage
Outliers	17 of 78400 reflections (0.022%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14176	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.65 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.4387e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.26	6/3404 (0.2%)	1.24	20/4620 (0.4%)
1	B	1.33	14/3405 (0.4%)	1.17	17/4620 (0.4%)
1	C	1.33	8/3404 (0.2%)	1.23	25/4620 (0.5%)
1	D	1.34	17/3405 (0.5%)	1.21	19/4620 (0.4%)
All	All	1.32	45/13618 (0.3%)	1.21	81/18480 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	4
1	C	0	5
1	D	1	4
All	All	1	18

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	195	TYR	CD1-CE1	10.56	1.55	1.39
1	C	112	CYS	CB-SG	-9.87	1.65	1.82
1	D	12	ALA	CA-CB	8.14	1.69	1.52
1	A	86	TRP	CB-CG	8.11	1.64	1.50
1	B	312	GLU	CB-CG	7.82	1.67	1.52

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	ARG	NE-CZ-NH1	15.51	128.06	120.30
1	A	283	ARG	NE-CZ-NH2	-12.25	114.18	120.30
1	C	14	ARG	NE-CZ-NH2	9.75	125.18	120.30
1	D	257	ARG	NE-CZ-NH1	9.66	125.13	120.30
1	C	321	ARG	NE-CZ-NH2	-8.95	115.82	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	188	LYS	CA

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	17	SER	Peptide
1	A	188	LYS	Peptide
1	A	19	ILE	Peptide
1	A	341	LEU	Peptide
1	A	342	THR	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3347	0	3358	206	0
1	B	3348	0	3358	200	0
1	C	3347	0	3358	209	0
1	D	3348	0	3358	205	0
2	A	12	0	16	1	0
2	B	6	0	8	0	0
2	C	30	0	40	6	0
2	D	24	0	32	24	0
3	A	166	0	0	25	0
3	B	199	0	0	27	0
3	C	166	0	0	17	0
3	D	183	0	0	34	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	14176	0	13528	779	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 29.

The worst 5 of 779 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:60:THR:HG22	3:D:555:HOH:O	1.38	1.22
1:D:182:ASP:HB2	3:D:568:HOH:O	1.40	1.22
1:D:137:LEU:C	1:D:137:LEU:HD23	1.61	1.20
1:C:166:ILE:HD11	1:C:216:ILE:CD1	1.71	1.18
1:C:166:ILE:HD11	1:C:216:ILE:HD11	1.24	1.16

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/425 (99%)	381 (90%)	33 (8%)	8 (2%)	12	9
1	B	422/425 (99%)	391 (93%)	26 (6%)	5 (1%)	19	19
1	C	422/425 (99%)	393 (93%)	24 (6%)	5 (1%)	19	19
1	D	422/425 (99%)	381 (90%)	33 (8%)	8 (2%)	12	9
All	All	1688/1700 (99%)	1546 (92%)	116 (7%)	26 (2%)	15	13

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	SER
1	A	28	ARG
1	B	203	PRO
1	C	203	PRO
1	D	19	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/369 (100%)	312 (85%)	57 (15%)	4	3
1	B	369/369 (100%)	315 (85%)	54 (15%)	5	4
1	C	369/369 (100%)	315 (85%)	54 (15%)	5	4
1	D	369/369 (100%)	300 (81%)	69 (19%)	2	2
All	All	1476/1476 (100%)	1242 (84%)	234 (16%)	4	3

5 of 234 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	367	LYS
1	C	137	LEU
1	D	327	SER
1	B	408	GLU
1	C	34	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	201	ASN
1	C	91	GLN
1	D	294	HIS
1	B	237	GLN
1	B	305	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	263	1	24,24,25	4.11	6 (25%)	30,32,34	1.41	6 (20%)
1	LLP	B	263	1	24,24,25	3.52	8 (33%)	30,32,34	1.74	8 (26%)
1	LLP	C	263	1	24,24,25	4.02	10 (41%)	30,32,34	2.24	9 (30%)
1	LLP	D	263	1	24,24,25	4.42	7 (29%)	30,32,34	2.15	11 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	263	1	-	0/15/17/19	0/1/1/1
1	LLP	B	263	1	-	0/15/17/19	0/1/1/1
1	LLP	C	263	1	-	0/15/17/19	0/1/1/1
1	LLP	D	263	1	-	0/15/17/19	0/1/1/1

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	263	LLP	O-C	20.03	1.25	1.11
1	A	263	LLP	O-C	18.51	1.24	1.11
1	C	263	LLP	O-C	16.26	1.22	1.11
1	B	263	LLP	O-C	13.94	1.21	1.11
1	C	263	LLP	O3-C3	-5.96	1.22	1.37

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	263	LLP	C4-C4'-NZ	6.87	124.31	111.52
1	D	263	LLP	C6-C5-C4	5.09	121.96	118.10
1	C	263	LLP	C-CA-N	-4.85	108.98	113.83
1	C	263	LLP	OP2-P-OP4	-4.29	94.80	106.65
1	D	263	LLP	C5'-C5-C6	-3.69	112.29	119.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	426	-	5,5,5	0.23	0	5,5,5	1.00	0
2	GOL	A	427	-	5,5,5	0.32	0	5,5,5	0.61	0
2	GOL	B	426	-	5,5,5	0.45	0	5,5,5	1.05	0
2	GOL	C	426	-	5,5,5	0.42	0	5,5,5	0.75	0
2	GOL	C	427	-	5,5,5	0.47	0	5,5,5	1.13	0
2	GOL	C	428	-	5,5,5	0.44	0	5,5,5	1.36	0
2	GOL	C	429	-	5,5,5	0.42	0	5,5,5	0.63	0
2	GOL	C	430	-	5,5,5	0.51	0	5,5,5	0.90	0
2	GOL	D	426	-	5,5,5	0.63	0	5,5,5	1.00	0
2	GOL	D	427	-	5,5,5	0.30	0	5,5,5	0.58	0
2	GOL	D	428	-	5,5,5	0.83	0	5,5,5	1.70	1 (20%)
2	GOL	D	429	-	5,5,5	1.06	0	5,5,5	2.15	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	426	-	-	0/4/4/4	0/0/0/0
2	GOL	A	427	-	-	0/4/4/4	0/0/0/0
2	GOL	B	426	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	426	-	-	0/4/4/4	0/0/0/0
2	GOL	C	427	-	-	0/4/4/4	0/0/0/0
2	GOL	C	428	-	-	0/4/4/4	0/0/0/0
2	GOL	C	429	-	-	0/4/4/4	0/0/0/0
2	GOL	C	430	-	-	0/4/4/4	0/0/0/0
2	GOL	D	426	-	-	0/4/4/4	0/0/0/0
2	GOL	D	427	-	-	0/4/4/4	0/0/0/0
2	GOL	D	428	-	-	0/4/4/4	0/0/0/0
2	GOL	D	429	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	429	GOL	O1-C1-C2	3.36	126.09	109.71
2	D	429	GOL	O2-C2-C3	-2.52	96.75	108.22
2	D	428	GOL	O3-C3-C2	2.51	121.97	109.71
2	D	429	GOL	O3-C3-C2	2.33	121.10	109.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/425 (100%)	0.31	16 (3%) 38 49	24, 36, 54, 78	0
1	B	425/425 (100%)	0.30	19 (4%) 32 42	23, 34, 51, 92	0
1	C	425/425 (100%)	0.32	24 (5%) 24 33	23, 35, 52, 77	0
1	D	425/425 (100%)	0.27	17 (4%) 36 47	23, 34, 52, 84	0
All	All	1700/1700 (100%)	0.30	76 (4%) 31 42	23, 35, 53, 92	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	27	SER	11.8
1	C	29	GLY	10.0
1	B	26	LEU	8.6
1	B	28	ARG	8.3
1	B	24	ASP	7.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LLP	D	263	24/25	0.18	0.95	23,32,35,35	0
1	LLP	B	263	24/25	0.18	0.49	23,30,33,34	0
1	LLP	A	263	24/25	0.16	0.26	23,33,36,36	0
1	LLP	C	263	24/25	0.15	-0.38	27,30,34,37	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	D	429	6/6	0.46	16.00	23,30,40,43	0
2	GOL	C	430	6/6	0.20	9.61	48,53,56,57	0
2	GOL	C	428	6/6	0.33	8.51	54,56,58,60	0
2	GOL	C	427	6/6	0.34	7.92	34,47,53,54	0
2	GOL	D	427	6/6	0.41	7.26	55,56,59,62	0
2	GOL	D	428	6/6	0.33	6.24	46,51,52,53	0
2	GOL	C	429	6/6	0.52	4.58	63,66,68,69	0
2	GOL	A	427	6/6	0.24	4.52	58,61,62,63	0
2	GOL	D	426	6/6	0.24	0.93	44,52,55,59	0
2	GOL	B	426	6/6	0.33	0.77	63,64,66,66	0
2	GOL	C	426	6/6	0.28	0.25	59,64,65,65	0
2	GOL	A	426	6/6	0.23	-0.29	68,70,71,72	0

6.5 Other polymers ⓘ

There are no such residues in this entry.