



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 12:50 PM GMT

PDB ID : 2QTW
Title : The Crystal Structure of PCSK9 at 1.9 Angstroms Resolution Reveals structural homology to Resistin within the C-terminal domain
Authors : Hampton, E.N.; Knuth, M.W.; Li, J.; Harris, J.L.; Lesley, S.A.; Spraggon, G.
Deposited on : 2007-08-02
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	124	
2	B	546	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4697 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9 Propeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	1	0
			748	479	136	131	2			

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	486	Total	C	N	O	S	0	1	0
			3614	2229	668	685	32			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	693	HIS	-	EXPRESSION TAG	UNP Q8NBP7
B	694	HIS	-	EXPRESSION TAG	UNP Q8NBP7
B	695	HIS	-	EXPRESSION TAG	UNP Q8NBP7
B	696	HIS	-	EXPRESSION TAG	UNP Q8NBP7
B	697	HIS	-	EXPRESSION TAG	UNP Q8NBP7
B	698	HIS	-	EXPRESSION TAG	UNP Q8NBP7

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	66	Total	O	0	0
			66	66		
5	B	254	Total	O	0	0
			254	254		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proprotein convertase subtilisin/kexin type 9 Propeptide

Chain A: 



- Molecule 2: Proprotein convertase subtilisin/kexin type 9

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.43Å 70.15Å 148.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 47.79 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-1.90) 99.8 (47.79-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.239 0.198 , 0.243	Depositor DCC
R_{free} test set	2654 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 52170 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4697	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/768	0.77	0/1037
2	B	0.73	3/3682 (0.1%)	0.75	1/4996 (0.0%)
All	All	0.74	3/4450 (0.1%)	0.75	1/6033 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	654	CYS	CB-SG	-6.35	1.71	1.82
2	B	255	CYS	C-N	-5.62	1.21	1.34
2	B	256	GLN	C-N	-5.29	1.23	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	179	LEU	CA-CB-CG	6.44	130.11	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	244	GLY	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	B	494	LYS	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	748	0	763	4	1
2	B	3614	0	3522	60	1
3	B	14	0	13	1	0
4	B	1	0	0	0	0
5	A	66	0	0	1	0
5	B	254	0	0	7	0
All	All	4697	0	4298	63	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (63) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:201:MET:HE1	2:B:203:THR:HG22	1.44	0.98
2:B:615:ILE:HD12	2:B:619:GLN:HE21	1.34	0.91
2:B:615:ILE:HD12	2:B:619:GLN:NE2	1.95	0.81
2:B:446:PRO:HB2	2:B:448:THR:HG22	1.63	0.78
2:B:185:LEU:HD11	2:B:271:ILE:HD11	1.65	0.77
2:B:186:ASP:OD2	2:B:230:LEU:HD12	1.85	0.76
2:B:302:GLN:HE21	2:B:306:ARG:HH21	1.34	0.76
2:B:529:LEU:HG	2:B:532:ALA:HB2	1.75	0.68
2:B:201:MET:HE1	2:B:203:THR:CG2	2.23	0.68
2:B:435:VAL:HG23	5:B:1061:HOH:O	1.94	0.68
2:B:343:ASP:HB3	2:B:423:VAL:HG12	1.76	0.66
2:B:437:THR:O	5:B:1167:HOH:O	2.15	0.64
2:B:612:GLU:HG2	2:B:675:VAL:HG22	1.81	0.62
2:B:176:GLY:O	2:B:244:GLY:HA3	2.00	0.61
2:B:343:ASP:OD2	2:B:423:VAL:HG11	2.00	0.61
2:B:343:ASP:CB	2:B:423:VAL:HG12	2.31	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:552:CYS:HB3	2:B:557[B]:HIS:CE1	2.39	0.58
2:B:615:ILE:CD1	2:B:619:GLN:HE21	2.13	0.58
2:B:638:LEU:HD13	2:B:675:VAL:HG21	1.86	0.56
2:B:189:ILE:CD1	2:B:200:VAL:HG11	2.37	0.55
2:B:638:LEU:HD22	2:B:673:THR:HG21	1.90	0.54
2:B:469:ARG:HD2	2:B:515:PHE:CD1	2.44	0.53
2:B:549:ARG:HG2	2:B:589:VAL:HG22	1.91	0.53
1:A:97:ARG:CZ	5:A:202:HOH:O	2.57	0.53
2:B:201:MET:CE	2:B:203:THR:HG22	2.29	0.52
2:B:345:PRO:HD3	2:B:424:ILE:HG23	1.91	0.52
1:A:67:CYS:SG	5:B:1256:HOH:O	2.04	0.52
2:B:343:ASP:CB	2:B:423:VAL:CG1	2.88	0.51
2:B:179:LEU:HD22	2:B:401:SER:HA	1.93	0.51
2:B:468:THR:OG1	2:B:471:ALA:HB2	2.11	0.50
2:B:177:GLY:HA2	2:B:401:SER:OG	2.12	0.50
2:B:632:LEU:HD13	2:B:676:ALA:HB1	1.95	0.49
2:B:193:HIS:HD2	2:B:195:GLU:H	1.61	0.49
2:B:531:GLN:O	3:B:1001:NAG:H82	2.13	0.49
2:B:632:LEU:CD1	2:B:676:ALA:HB1	2.43	0.48
2:B:455:LEU:HD22	2:B:606:LEU:HD11	1.96	0.47
2:B:186:ASP:OD1	2:B:226:HIS:ND1	2.36	0.47
2:B:201:MET:CE	2:B:203:THR:CG2	2.91	0.46
2:B:554:GLN:HB2	2:B:557[B]:HIS:CE1	2.51	0.46
2:B:435:VAL:HG12	2:B:461:TRP:CZ2	2.51	0.46
1:A:152:GLN:C	2:B:386:SER:OG	2.54	0.46
2:B:557[A]:HIS:CE1	2:B:602:HIS:HB2	2.52	0.45
2:B:410:GLU:HA	2:B:528:LEU:HD11	1.99	0.44
2:B:493:GLY:CA	2:B:645:LEU:HD12	2.48	0.44
2:B:200:VAL:HG22	2:B:247:MET:HB2	2.00	0.44
2:B:493:GLY:HA3	2:B:645:LEU:HD12	1.99	0.43
2:B:417:HIS:ND1	5:B:1181:HOH:O	2.36	0.42
2:B:495:ARG:CG	5:B:1133:HOH:O	2.66	0.42
2:B:194:ARG:HG3	2:B:377:THR:HG22	2.02	0.42
2:B:557[B]:HIS:HB2	2:B:601:CYS:O	2.20	0.42
2:B:201:MET:HE2	2:B:201:MET:HB3	1.95	0.42
2:B:289:LEU:HD12	2:B:289:LEU:C	2.41	0.42
2:B:302:GLN:O	2:B:306:ARG:HG3	2.19	0.41
2:B:166:TYR:HA	2:B:447:SER:OG	2.20	0.41
2:B:495:ARG:HG2	5:B:1133:HOH:O	2.19	0.41
2:B:359:VAL:O	2:B:438:PRO:HG2	2.20	0.41
1:A:70:ASP:N	1:A:71:PRO:CD	2.83	0.41
2:B:552:CYS:HB2	2:B:586:ASN:HB3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:490:SER:OG	2:B:493:GLY:N	2.51	0.41
2:B:554:GLN:H	2:B:557[B]:HIS:CE1	2.39	0.41
2:B:243:LYS:O	2:B:244:GLY:C	2.58	0.41
2:B:680:ARG:NE	5:B:1185:HOH:O	2.52	0.40
2:B:499:ARG:HD3	2:B:512:HIS:NE2	2.36	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:78:TYR:OH	2:B:405:GLU:OE2[3.655]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/124 (73%)	90 (99%)	1 (1%)	0	100	100
2	B	471/546 (86%)	459 (98%)	9 (2%)	3 (1%)	33	19
All	All	562/670 (84%)	549 (98%)	10 (2%)	3 (0%)	38	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	244	GLY
2	B	545	SER
2	B	280	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/104 (77%)	79 (99%)	1 (1%)	80	77
2	B	389/437 (89%)	378 (97%)	11 (3%)	56	45
All	All	469/541 (87%)	457 (97%)	12 (3%)	59	49

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	ARG
2	B	201	MET
2	B	278	GLN
2	B	375	CYS
2	B	490	SER
2	B	571	LEU
2	B	615	ILE
2	B	623	THR
2	B	638	LEU
2	B	644	VAL
2	B	645	LEU
2	B	659	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	HIS
1	A	116	HIS
2	B	193	HIS
2	B	302	GLN
2	B	449	HIS
2	B	537	HIS
2	B	554	GLN
2	B	602	HIS
2	B	619	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	1001	2	12,14,15	0.65	0	15,19,21	1.05	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1001	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1001	NAG	C3-C4-C5	2.18	114.10	110.20
3	B	1001	NAG	O5-C5-C6	2.16	109.25	106.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/124 (74%)	0.07	4 (4%) 34 34	17, 23, 37, 44	0
2	B	486/546 (89%)	0.59	45 (9%) 9 8	13, 26, 53, 74	0
All	All	578/670 (86%)	0.51	49 (8%) 11 10	13, 25, 52, 74	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	515	PHE	10.7
2	B	571	LEU	6.6
2	B	546	MET	6.1
2	B	642	SER	5.3
2	B	220	ALA	5.2
2	B	212	ASP	4.9
2	B	449	HIS	4.8
2	B	570	ASP	4.7
2	B	568	VAL	4.6
2	B	616	PRO	4.5
2	B	659	ARG	4.4
2	B	615	ILE	4.4
2	B	167	ARG	4.4
2	B	470	MET	3.9
2	B	448	THR	3.9
1	A	61	THR	3.8
2	B	531	GLN	3.7
2	B	671	ALA	3.7
2	B	469	ARG	3.7
2	B	620	GLU	3.7
2	B	447	SER	3.6
2	B	569	GLU	3.5
2	B	453	TRP	3.5
2	B	619	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	584	GLN	3.3
2	B	516	GLY	3.3
2	B	543	GLU	3.2
2	B	672	VAL	3.2
2	B	643	HIS	3.1
2	B	505	GLY	3.0
2	B	194	ARG	2.9
2	B	221	SER	2.8
2	B	517	GLY	2.8
2	B	452	GLY	2.7
1	A	108	LEU	2.6
2	B	514	ALA	2.6
2	B	555	GLN	2.6
2	B	530	PRO	2.6
2	B	658	SER	2.5
2	B	604	PRO	2.5
1	A	104	ARG	2.4
1	A	103	ALA	2.4
2	B	255	CYS	2.3
2	B	237	ARG	2.3
2	B	503	GLN	2.2
2	B	199	ARG	2.1
2	B	591	HIS	2.1
2	B	494	LYS	2.1
2	B	614	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	B	1001	14/15	0.33	1.11	57,62,65,69	0
4	CA	B	1	1/1	0.17	0.79	61,61,61,61	0

6.5 Other polymers ⓘ

There are no such residues in this entry.